


Q A 533

W. H. Meyer

1894

H 22



Digitized by the Internet Archive  
in 2010 with funding from  
Boston Library Consortium Member Libraries



# ELEMENTARY TRIGONOMETRY.

Library of the  
State of New York



THE  
MUSIC

# ELEMENTARY TRIGONOMETRY

BY

H. S. HALL, M.A.,

FORMERLY SCHOLAR OF CHRIST'S COLLEGE, CAMBRIDGE;  
MASTER OF THE MILITARY SIDE, CLIFTON COLLEGE.

AND

S. R. KNIGHT, B.A., M.B., CH.B.,

FORMERLY SCHOLAR OF TRINITY COLLEGE, CAMBRIDGE;  
LATE ASSISTANT MASTER AT MARLBOROUGH COLLEGE.

BOSTON COLLEGE LIBRARY  
CHESTNUT HILL, MASS.

London:

MACMILLAN AND CO.

AND NEW YORK.

1893

[All Rights reserved.]

Cambridge:  
PRINTED BY C. J. CLAY, M.A. AND SONS,  
AT THE UNIVERSITY PRESS.

QA 533  
. H 22

150028

## PREFACE.

THE following pages will be found to comprise all the parts of Elementary Trigonometry which can conveniently be treated without the use of infinite series and imaginary quantities.

The chapters have been subdivided into short sections, and the examples to illustrate each section have been very carefully selected and arranged, the earlier ones being easy enough for any reader to whom the subject is new, while the later ones, and the Miscellaneous Examples scattered throughout the book, will furnish sufficient practice for those who intend to pursue the subject further as part of a mathematical education.

No substantial progress in Trigonometry can be made until the fundamental properties of the Trigonometrical Ratios have been thoroughly mastered. To attain this object very considerable practice in easy Identities and Equations is necessary. We have therefore given special prominence to examples of this kind in the early pages; with the same end in view we have postponed the subject of Radian or Circular Measure to a later stage than is usual, believing that it is in every way more satis-

factory to dwell on the properties of the trigonometrical ratios, and to exemplify their use in easy problems, than to bewilder a beginner with an angular system the use of which he cannot appreciate, and which at this stage furnishes nothing but practice in easy Arithmetic.

The subject of Logarithms and their application has been treated very fully, and illustrated by a selection of carefully graduated Examples. It is hoped that the examples worked out in this section may serve as useful models for the student, and may do something to cure that inaccuracy in logarithmic work which is so often due to clumsy arrangement.

In the experience of most teachers it is found extremely difficult to get boys to handle problems in Heights and Distances with any degree of confidence and skill. Accordingly we have devoted much thought to the exposition of this part of the subject, and by careful classification of the Examples we have endeavoured to make Chapters VI. and XVII. as easy and attractive as possible.

Very little advance can be expected in Trigonometry until the principal formulæ can be quoted readily, but whether it is advisable for learners to have lists of formulæ compiled for them, so as to be easily accessible at all times, is a matter upon which teachers hold different views. In our own opinion it is distinctly mischievous to furnish such lists; it encourages indolent habits, and fosters a spurious confidence which leads to disaster when the student has to rely solely upon his own knowledge.

In the general arrangement and succession of the different parts of the subject we have been mainly guided by our own long experience in the class room; but as the manuscript and proof-sheets have been read by several skilled teachers, and have been frequently tested by pupils

in all stages of proficiency, the hope is entertained that our treatment is such as to enable beginners to take an intelligent interest in the subject from the first, and to acquire a sound elementary knowledge of practical Trigonometry before they encounter the more theoretical difficulties. At the same time, as each chapter is, as far as possible, complete in itself, it will be easy for teachers to adopt a different order of treatment if they prefer it; the full Table of Contents will facilitate the selection of a suitable course of reading, besides furnishing a useful aid to students who are rapidly revising the subject.

We are indebted to several friends for valuable criticism and advice; in particular, we have to thank Mr T. D. Davies of Clifton College for many useful hints, and for some ingenious examples and solutions in Chapters xxiv. and xxv.

H. S. HALL.

S. R. KNIGHT.

*November, 1893.*





## CONTENTS.

### Chapter I. MEASUREMENT OF ANGLES.

	PAGE
Definition of Angle . . . . .	1
Sexagesimal and Centesimal Measures . . . . .	2
Formula $\frac{D}{9} = \frac{G}{10}$ . . . . .	3

### Chapter II. TRIGONOMETRICAL RATIOS.

Definitions of Ratio and Commensurable Quantities . . . . .	5
Definitions of the Trigonometrical Ratios . . . . .	6
Sine and cosine are less than unity, secant and cosecant are greater than unity, tangent and cotangent are unrestricted . . . . .	7
The trigonometrical ratios are independent of the lengths of the lines which include the angle . . . . .	9
Definition of <i>function</i> . . . . .	10

### Chapter III. RELATIONS BETWEEN THE RATIOS.

The reciprocal relations . . . . .	12
Tangent and cotangent in terms of sine and cosine . . . . .	13
Sine-cosine, tangent-secant, cotangent-cosecant formulæ . . . . .	14
Easy Identities . . . . .	16
Each ratio can be expressed in terms of any of the others . . . . .	21

### Chapter IV. TRIGONOMETRICAL RATIOS OF CERTAIN ANGLES.

Trigonometrical Ratios of $45^\circ$ , $60^\circ$ , $30^\circ$ . . . . .	24
Definition of complementary angles . . . . .	27
Each function of an angle is equal to the corresponding co-function of its complement . . . . .	27
Easy Trigonometrical Equations . . . . .	30
Miscellaneous Examples. A. . . . .	32

**Chapter V. SOLUTION OF RIGHT-ANGLED TRIANGLES.**

	PAGE
CASE I. When two sides are given . . . . .	35
CASE II. When one side and one acute angle are given . . . . .	36
Case of triangle considered as sum or difference of two right-angled triangles . . . . .	38

**Chapter VI. EASY PROBLEMS.**

Angles of Elevation and Depression . . . . .	41
The Mariner's Compass . . . . .	45

**Chapter VII. RADIAN OR CIRCULAR MEASURE.**

Definition of Radian . . . . .	49
Circumference of circle = $2\pi$ (radius) . . . . .	50
All radians are equal . . . . .	51
$\pi$ radians = 2 right angles = 180 degrees . . . . .	52
Radian contains 57.2958 degrees . . . . .	52
Formula $\frac{D}{180} = \frac{\theta}{\pi}$ . . . . .	53
Values of the functions of $\frac{\pi}{4}, \frac{\pi}{3}, \frac{\pi}{6}$ . . . . .	55
Ratios of the complementary angle $\frac{\pi}{2} - \theta$ . . . . .	56
Radian measure of angles of a regular polygon . . . . .	56
Radian measure of an angle = $\frac{\text{subtending arc}}{\text{radius}}$ . . . . .	58
Radian and Circular Measures are equivalent . . . . .	58
Miscellaneous Examples. B. . . . .	61

**Chapter VIII. RATIOS OF ANGLES OF ANY MAGNITUDE.**

Convention of Signs (1) for line, (2) for plane surface, (3) for angles . . . . .	64
Definitions of the trigonometrical ratios of any angle . . . . .	66
Signs of the trigonometrical ratios in the four quadrants . . . . .	68
Definition of Coterminal Angles . . . . .	68
The fundamental formulæ of Chap. III. are true for all values of the angle . . . . .	70
The ambiguity of sign in $\cos A = \pm \sqrt{1 - \sin^2 A}$ can be removed when $A$ is known . . . . .	71

**Chapter IX. VARIATIONS OF THE FUNCTIONS.**

	PAGE
Definition of <i>limit</i> . . . . .	74
Functions of $0^\circ$ and $90^\circ$ . . . . .	74
Changes in the sign and magnitude of $\sin A$ . . . . .	76
Changes in the sign and magnitude of $\tan A$ . . . . .	78
Definition of Circular Functions . . . . .	79
Note on the old definitions . . . . .	80
Miscellaneous Examples. C. . . . .	81

**Chapter X. CIRCULAR FUNCTIONS OF ALLIED ANGLES.**

Circular functions of $180^\circ - A$ . . . . .	82
Definition of supplementary angles . . . . .	83
Circular functions of $180^\circ + A$ . . . . .	84
Circular functions of $90^\circ + A$ . . . . .	85
Circular functions of $-A$ . . . . .	86
Definition of even and odd functions . . . . .	87
Circular functions of $90^\circ - A$ for any value of $A$ . . . . .	88
Circular functions of $n \cdot 360^\circ \pm A$ . . . . .	89
The functions of any angle can be expressed as the same functions of some acute angle . . . . .	90
The number of angles which have the same trigonometrical ratio is unlimited . . . . .	91

**Chapter XI. FUNCTIONS OF COMPOUND ANGLES.**

Expansions of the sine and cosine of $A+B$ and $A-B$ . . . . .	94
$\sin(A+B) \sin(A-B) = \sin^2 A - \sin^2 B$ . . . . .	96
Expansions of $\tan(A+B)$ and $\cot(A+B)$ . . . . .	98
Expansions of $\sin(A+B+C)$ and $\tan(A+B+C)$ . . . . .	99
Converse use of the Addition Formulæ . . . . .	100
Functions of $2A$ . . . . .	102
Functions of $3A$ . . . . .	105
Value of $\sin 18^\circ$ . . . . .	106

**Chapter XII. TRANSFORMATION OF PRODUCTS AND SUMS.**

Transformation of products into sums or differences . . . . .	110
Transformation of sums or differences into products . . . . .	112
Relations when $A+B+C=180^\circ$ . . . . .	118

# Chapter XIII. RELATIONS BETWEEN THE SIDES AND ANGLES OF A TRIANGLE.

	PAGE
$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ . . . . .	123
$a^2 = b^2 + c^2 - 2bc \cos A$ , and $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$ . . . . .	124
$a = b \cos C + c \cos B$ . . . . .	124
The above sets of formulæ are not independent . . . . .	125
Solution of Triangles without logarithms . . . . .	126
CASE I. When the three sides are given . . . . .	126
CASE II. When two sides and the included angle are given . . . . .	127
CASE III. When two angles and a side are given . . . . .	128
CASE IV. When two sides and an angle opposite to one of them are given. Sometimes this case is <i>ambiguous</i> . . . . .	130
The Ambiguous Case discussed geometrically . . . . .	131
The Ambiguous Case discussed by first finding the third side . . . . .	135
Miscellaneous Examples. D. . . . .	138

# Chapter XIV. LOGARITHMS.

$a^x = N$ and $x = \log_a N$ are equivalent . . . . .	139
$a^{\log_a N} = N$ is identically true . . . . .	139
Logarithm of a product, quotient, power, root . . . . .	140
The characteristic of a common logarithm may be written down by inspection . . . . .	142
The logarithms of all numbers which have the same significant digits have the same mantissa . . . . .	143
$\log_b N = \frac{1}{\log_a b} \times \log_a N$ , and $\log_b a \times \log_a b = 1$ . . . . .	146
Exponential Equations . . . . .	148
Miscellaneous Examples. E. . . . .	150

# Chapter XV. THE USE OF LOGARITHMIC TABLES.

Rule of Proportional Parts . . . . .	152
Use of Tables of Common Logarithms . . . . .	152
Use of Tables of Natural and Logarithmic Functions . . . . .	156

**Chapter XVI.** SOLUTION OF TRIANGLES WITH LOGARITHMS.

	PAGE
Functions of the half-angles in terms of the sides . . . . .	164
Sin $A$ in terms of the sides . . . . .	166
Solution when the three sides are given . . . . .	167
Solution when two sides and the included angle are given . . . . .	170
Solution when two angles and a side are given . . . . .	174
Solution when two sides and the angle opposite to one of them are given . . . . .	175
Adaptation of $a^2 + b^2$ to logarithmic work . . . . .	177
Adaptation of $a^2 + b^2 - 2ab \cos C$ to logarithmic work . . . . .	178

**Chapter XVII.** HEIGHTS AND DISTANCES.

Measurements in one plane . . . . .	184
Problems dependent on Geometry . . . . .	189
Measurements in more than one plane . . . . .	193

**Chapter XVIII.** PROPERTIES OF TRIANGLES AND POLYGONS.

Area of a triangle . . . . .	198
Radius of the circum-circle of a triangle . . . . .	200
Radius of the in-circle of a triangle . . . . .	201
Radii of the ex-circles of a triangle . . . . .	202
Some important relations established by Geometry . . . . .	204
Inscribed and circumscribed polygons . . . . .	208
Area of a circle and sector of a circle . . . . .	210
The Ex-central Triangle . . . . .	212
The Pedal Triangle . . . . .	214
Distances of in-centre and ex-centres from circum-centre . . . . .	216
Distance of orthocentre from circum-centre . . . . .	218
Area of any quadrilateral . . . . .	220
Diagonals and circum-radius of a cyclic quadrilateral . . . . .	222
Miscellaneous Examples. F. . . . .	228

**Chapter XIX.** GENERAL VALUES AND INVERSE FUNCTIONS.

Formula for all angles which have a given sine . . . . .	232
Formula for all angles which have a given cosine . . . . .	233
Formula for all angles which have a given tangent . . . . .	234
Formula for angles both equi-sinal and equi-cosinal . . . . .	234

	PAGE
General solution of equations . . . . .	236
Inverse Circular Functions . . . . .	238
Solution of equations expressed in inverse notation . . . . .	244
Miscellaneous Examples. G. . . . .	246

## Chapter XX. FUNCTIONS OF SUBMULTIPLE ANGLES.

Trigonometrical Ratios of $\frac{\pi}{8}$ . . . . .	247
Given $\cos A$ to find $\sin \frac{A}{2}$ and $\cos \frac{A}{2}$ . . . . .	248
To express $\sin \frac{A}{2}$ and $\cos \frac{A}{2}$ in terms of $\sin A$ . . . . .	250
Variation in sign and magnitude of $\cos \theta - \sin \theta$ . . . . .	254
Sine and cosine of $9^\circ$ . . . . .	254
To find $\tan \frac{A}{2}$ when $\tan A$ is given . . . . .	256
Given a function of $A$ to find the functions of $\frac{A}{2}$ . . . . .	258
Given $\cos A$ to find $\cos \frac{A}{3}$ . . . . .	259

## Chapter XXI. LIMITS AND APPROXIMATIONS.

If $\theta < \frac{\pi}{2}$ , $\sin \theta$ , $\theta$ , $\tan \theta$ are in ascending order of magnitude . . . . .	261
When $\theta = 0$ , $\frac{\sin \theta}{\theta} = 1$ , and $\frac{\tan \theta}{\theta} = 1$ . . . . .	262
$\cos \theta > 1 - \frac{\theta^2}{2}$ , and $\sin \theta > \theta - \frac{\theta^3}{4}$ . . . . .	265
Value of $\sin 10''$ . . . . .	266
$\cos \frac{\theta}{2} \cos \frac{\theta}{4} \cos \frac{\theta}{8} \cos \frac{\theta}{16} \dots = \frac{\sin \theta}{\theta}$ . . . . .	266
$\frac{\sin \theta}{\theta}$ decreases from 1 to $\frac{2}{\pi}$ as $\theta$ increases from 0 to $\frac{\pi}{2}$ . . . . .	267
Distance and Dip of the Visible Horizon . . . . .	269

## Chapter XXII. GEOMETRICAL PROOFS.

Expansion of $\tan(A+B)$ . . . . .	273
Formulae for transformation of sums into products . . . . .	274



	PAGE
Proof of the 2A formulæ . . . . .	276
Value of $\sin 18^\circ$ . . . . .	277
Proofs by Projection . . . . .	278
General analytical proof of the Addition Formulæ . . . . .	282
Miscellaneous Examples. H. . . . .	283
Graphs of $\sin \theta$ , $\tan \theta$ , $\sec \theta$ . . . . .	285

### Chapter XXIII. SUMMATION OF FINITE SERIES.

If $u_r = v_{r+1} - v_r$ , then $S = v_{n+1} - v_1$ . . . . .	288
Sum of the sines and cosines of a series of $n$ angles in A. P. . . . .	289
When the common difference is $\frac{2k\pi}{n}$ , the sum is zero . . . . .	290
Sum of the squares and cubes of the sines and cosines of a series of angles in A. P. . . . .	293

### Chapter XXIV. MISCELLANEOUS TRANSFORMATIONS AND IDENTITIES.

Symmetrical Expressions. $\Sigma$ and $\Pi$ notation . . . . .	296
Alternating Expressions . . . . .	303
Allied formulæ in Algebra and Trigonometry . . . . .	306
Identities derived by substitution . . . . .	308

### Chapter XXV. MISCELLANEOUS THEOREMS AND EXAMPLES.

Inequalities. Maxima and Minima . . . . .	313
Elimination . . . . .	319
Application of Trigonometry to Theory of Equations . . . . .	326
Application of Theory of Equations to Trigonometry . . . . .	328
Miscellaneous Examples. I. . . . .	336





# ELEMENTARY TRIGONOMETRY.


## CHAPTER I.

### MEASUREMENT OF ANGLES.

1. THE word Trigonometry in its primary sense signifies the measurement of triangles. From an early date the science also included the establishment of the relations which subsist between the sides, angles, and area of a triangle; but now it has a much wider scope and embraces all manner of geometrical and algebraical investigations carried on through the medium of certain quantities called trigonometrical ratios, which will be defined in Chap. II. In every branch of Higher Mathematics, whether Pure or Applied, a knowledge of Trigonometry is of the greatest value.

2. **Definition of Angle.** Suppose that the straight line  $OP$  in the figure is capable of revolving about the point  $O$ , and suppose that in this way it has passed successively from the position  $OA$  to the positions occupied by  $OB$ ,  $OC$ ,  $OD$ , ..., then the angle between  $OA$  and any position such as  $OC$  is measured by the *amount of revolution* which the line  $OP$  has undergone in passing from its initial position  $OA$  into its final position  $OC$ .

Moreover the line  $OP$  may make any number of complete revolutions through the original position  $OA$  before taking up its final position.



It will thus be seen that in Trigonometry angles are not restricted as in Euclid, but may be of any magnitude.

The point  $O$  is called the *origin*, and  $OA$  the *initial line*; the revolving line  $OP$  is known as the *generating line* or the *radius vector*.

**3. Measurement of Angles.** We must first select some *fixed* unit. The natural unit would be a right angle, but as in practice this is inconveniently large, two systems of measurement have been established, in each of which the unit is a certain fraction of a right angle.

**4. Sexagesimal Measure.** A right angle is divided into 90 equal parts called *degrees*, a degree into 60 equal parts called *minutes*, a minute into 60 equal parts called *seconds*. An angle is measured by stating the number of degrees, minutes, and seconds which it contains.

For shortness, each of these three divisions, degrees, minutes, seconds, is denoted by a symbol; thus the angle which contains 53 degrees 37 minutes 2.53 seconds is expressed symbolically in the form  $53^{\circ} 37' 2.53''$ .

**5. Centesimal Measure.** A right angle is divided into 100 equal parts called *grades*, a grade into 100 equal parts called *minutes*, a minute into 100 equal parts called *seconds*. In this system the angle which contains 53 grades 37 minutes 2.53 seconds is expressed symbolically in the form  $53^{\circ} 37' 2.53''$ .

It will be noticed that different accents are used to denote sexagesimal and centesimal minutes and seconds; for though they have the same names, a centesimal minute and second are not the same as a sexagesimal minute and second. Thus a right angle contains  $90 \times 60$  sexagesimal minutes, whereas it contains  $100 \times 100$  centesimal minutes.

Sexagesimal Measure is sometimes called the English System, and Centesimal Measure the French System.

**6.** In *numerical* calculations the sexagesimal measure is always used. The centesimal method was proposed at the time of the French Revolution as part of a general system of decimal measurement, but has never been adopted even in France, as it would have made necessary the alteration of Geographical, Nautical, Astronomical, and other tables prepared according to the sexagesimal method. Beyond giving a few examples in transformation from one system to the other which afford exercise in easy Arithmetic, we shall after this rarely allude to centesimal measure.

In *theoretical* work it is convenient to use another method of measurement, where the unit is the angle subtended at the centre of a circle by an arc whose length is equal to the radius. This system is known as **Circular** or **Radian Measure**, and will be fully explained in Chapter VII.

An angle is usually represented by a single letter, different letters  $A, B, C, \dots, \alpha, \beta, \gamma, \dots, \theta, \phi, \psi, \dots$ , being used to distinguish different angles. For angles estimated in sexagesimal or centesimal measure these letters are used indifferently, but we shall always denote angles in circular measure by letters taken from the Greek alphabet.

7. If the number of degrees and grades contained in an angle be  $D$  and  $G$  respectively, to prove that  $\frac{D}{9} = \frac{G}{10}$ .

In sexagesimal measure, the given angle when expressed as the fraction of a right angle is denoted by  $\frac{D}{90}$ . In centesimal measure, the same fraction is denoted by  $\frac{G}{100}$ ;

$$\therefore \frac{D}{90} = \frac{G}{100}; \text{ that is, } \frac{D}{9} = \frac{G}{10}.$$

8. To pass from one system to the other it is advisable first to express the given angle in terms of a right angle.

In centesimal measure any number of grades, minutes, and seconds may be immediately expressed as the decimal of a right angle. Thus

$$23 \text{ grades} = \frac{23}{100} \text{ of a right angle} = \cdot 23 \text{ of a right angle};$$

$$15 \text{ minutes} = \frac{15}{100} \text{ of a grade} = \cdot 15 \text{ of a grade} = \cdot 0015 \text{ of a right angle};$$

$$\therefore 23^{\circ} 15' = \cdot 2315 \text{ of a right angle.}$$

$$\text{Similarly, } 15^{\circ} 7' 53\cdot 4'' = \cdot 1507534 \text{ of a right angle.}$$

Conversely, any decimal of a right angle can be at once expressed in grades, minutes, and seconds. Thus

$$\begin{aligned} \cdot 2173025 \text{ of a right angle} &= 21\cdot 73025^{\circ} \\ &= 21^{\circ} 73\cdot 025' \\ &= 21^{\circ} 73' 2\cdot 5''. \end{aligned}$$

In practice the intermediate steps are omitted.

*Example 1.* Reduce  $2^{\circ} 13' 4.5''$  to sexagesimal measure.

This angle =  $\cdot 0213045$  of a right angle  
 $= 1^{\circ} 55' 2.658''$ .

$$\begin{array}{r} \cdot 0213045 \text{ of a right angle} \\ \hline 90 \\ \hline 1.917405 \text{ degrees} \\ \hline 60 \\ \hline 55.0443 \text{ minutes} \\ \hline 60 \\ \hline 2.658 \text{ seconds.} \end{array}$$

OBS. In the Answers we shall express the angles to the nearest tenth of a second, so that the above result would be written  $1^{\circ} 55' 2.7''$ .

*Example 2.* Reduce  $12^{\circ} 13' 14.3''$  to centesimal measure.

This angle =  $\cdot 13578487$ ... of a right angle  
 $= 13^{\circ} 57' 84.9''$ .

$$\begin{array}{r} 60 ) 14.3 \text{ seconds} \\ \hline 60 ) 13.238333 \dots \text{minutes} \\ \hline 90 ) 12.2206388 \dots \text{degrees} \\ \hline \cdot 13578487 \dots \text{of a right angle.} \end{array}$$

### EXAMPLES. I.

Express as the decimal of a right angle

- |                           |                          |                           |
|---------------------------|--------------------------|---------------------------|
| 1. $67^{\circ} 30'$ .     | 2. $11^{\circ} 15'$ .    | 3. $37^{\circ} 50'$ .     |
| 4. $2^{\circ} 10' 12''$ . | 5. $8^{\circ} 0' 36''$ . | 6. $2^{\circ} 4' 4.5''$ . |

Reduce to centesimal measure

- |                               |                              |                              |
|-------------------------------|------------------------------|------------------------------|
| 7. $69^{\circ} 13' 30''$ .    | 8. $19^{\circ} 0' 45''$ .    | 9. $50^{\circ} 37' 5.7''$ .  |
| 10. $43^{\circ} 52' 38.1''$ . | 11. $11^{\circ} 0' 38.4''$ . | 12. $142^{\circ} 15' 45''$ . |
| 13. $12' 9''$ .               | 14. $3' 26.3''$ .            |                              |

Reduce to sexagesimal measure

- |                             |                            |                              |
|-----------------------------|----------------------------|------------------------------|
| 15. $56^{\circ} 87' 50''$ . | 16. $39^{\circ} 6' 25''$ . | 17. $40^{\circ} 1' 25.4''$ . |
| 18. $1^{\circ} 2' 3''$ .    | 19. $3^{\circ} 2' 5''$ .   | 20. $8^{\circ} 10' 6.5''$ .  |
| 21. $6' 25''$ .             | 22. $37' 5''$ .            |                              |

23. The sum of two angles is  $80^{\circ}$  and their difference is  $18^{\circ}$ ; find the angles in degrees.

24. The number of degrees in a certain angle added to the number of grades in the angle is 152: what is the angle?

25. If the same angle contains in English measure  $x$  minutes, and in French measure  $y$  minutes, prove that  $50x = 27y$ .

26. If  $s$  and  $t$  respectively denote the numbers of sexagesimal and centesimal seconds in any angle, prove that

$$250s = 81t.$$

## CHAPTER II.

### TRIGONOMETRICAL RATIOS.

9. DEFINITION. **Ratio** is the relation which one quantity bears to another of the *same* kind, the comparison being made by considering what multiple, part or parts, one quantity is of the other.

To find what multiple or part  $A$  is of  $B$  we divide  $A$  by  $B$ ; hence the ratio of  $A$  to  $B$  may be measured by the fraction  $\frac{A}{B}$ .

In order to compare two quantities they must be expressed in terms of the same unit. Thus the ratio of 2 yards to 27 inches is measured by the fraction  $\frac{2 \times 3 \times 12}{27}$  or  $\frac{8}{3}$ .

OBS. Since a ratio expresses the *number* of times that one quantity contains another, *every ratio is a numerical quantity*.

10. DEFINITION. If the ratio of any two quantities can be expressed exactly by the ratio of two integers the quantities are said to be **commensurable**; otherwise, they are said to be **incommensurable**. For instance, the quantities  $8\frac{1}{3}$  and  $5\frac{1}{3}$  are commensurable, while the quantities  $\sqrt{2}$  and 3 are incommensurable. But by finding the numerical value of  $\sqrt{2}$  we may express the value of the ratio  $\sqrt{2} : 3$  by the ratio of two commensurable quantities to any required degree of approximation. Thus to 5 decimal places  $\sqrt{2}=1.41421$ , and therefore to the same degree of approximation


$$\sqrt{2} : 3 = 1.41421 : 3 = 141421 : 300000.$$

Similarly, for the ratio of any two incommensurable quantities.



## Trigonometrical Ratios.

11. Let  $PAQ$  be any acute angle; in  $AP$  one of the boundary lines take a point  $B$  and draw  $BC$  perpendicular to  $AQ$ . Thus a right-angled triangle  $BAC$  is formed.



With reference to the angle  $A$  the following definitions are employed.

The ratio  $\frac{BC}{AB}$  or  $\frac{\text{opposite side}}{\text{hypotenuse}}$  is called the **sine of A**.

The ratio  $\frac{AC}{AB}$  or  $\frac{\text{adjacent side}}{\text{hypotenuse}}$  is called the **cosine of A**.

The ratio  $\frac{BC}{AC}$  or  $\frac{\text{opposite side}}{\text{adjacent side}}$  is called the **tangent of A**.

The ratio  $\frac{AC}{BC}$  or  $\frac{\text{adjacent side}}{\text{opposite side}}$  is called the **cotangent of A**.

The ratio  $\frac{AB}{AC}$  or  $\frac{\text{hypotenuse}}{\text{adjacent side}}$  is called the **secant of A**.

The ratio  $\frac{AB}{BC}$  or  $\frac{\text{hypotenuse}}{\text{opposite side}}$  is called the **cosecant of A**.

These six ratios are known as the **trigonometrical ratios**. It will be shewn later that as long as the angle remains the same the trigonometrical ratios remain the same. [Art. 19.]

12. Instead of writing in full the words *sine*, *cosine*, *tangent*, *cotangent*, *secant*, *cosecant*, abbreviations are adopted. Thus the above definitions may be more conveniently expressed and arranged as follows :

$$\sin A = \frac{BC}{AB}, \quad \text{cosec } A = \frac{AB}{BC},$$

$$\cos A = \frac{AC}{AB}, \quad \sec A = \frac{AB}{AC},$$

$$\tan A = \frac{BC}{AC}, \quad \cot A = \frac{AC}{BC}.$$

In addition to these six ratios, two others, the *versed sine* and *coversed sine* are sometimes used; they are written *vers A* and *covers A* and are thus defined:


$$\text{vers } A = 1 - \cos A, \quad \text{covers } A = 1 - \sin A.$$

13. In Chapter VIII. the definitions of the trigonometrical ratios will be extended to the case of angles of any magnitude, but for the present we confine our attention to the consideration of acute angles.

14. Although the verbal form of the definitions of the trigonometrical ratios given in Art. 11 may be helpful to the student at first, he will gain no freedom in their use until he is able to write down from the figure any ratio at sight.

In the adjoining figure,  $PQR$  is a right-angled triangle in which  $PQ=13$ ,  $PR=5$ ,  $QR=12$ .

Since  $PQ$  is the greatest side,  $R$  is the right angle. The trigonometrical ratios of the angles  $P$  and  $Q$  may be written down at once; for example,



$$\sin Q = \frac{PR}{PQ} = \frac{5}{13}, \quad \cos Q = \frac{QR}{PQ} = \frac{12}{13},$$

$$\tan P = \frac{QR}{PR} = \frac{12}{5}, \quad \operatorname{cosec} P = \frac{PQ}{QR} = \frac{13}{12}.$$

15. It is important to observe that *the trigonometrical ratios of an angle are numerical quantities*. Each one of them represents the *ratio of one length to another*, and they must themselves never be regarded as lengths.

16. In every right-angled triangle the hypotenuse is the greatest side; hence from the definitions of Art. 11 it will be seen that those ratios which have the hypotenuse in the *denominator* can never be greater than unity, while those which have the hypotenuse in the *numerator* can never be less than unity. Those ratios which do not involve the hypotenuse are not thus restricted in value, for either of the two sides which subtend the acute angles may be the greater. Hence

*the sine and cosine of an angle can never be greater than 1;*  
*the cosecant and secant of an angle can never be less than 1;*  
*the tangent and cotangent may have any numerical value.*

17. Let  $ABC$  be a right-angled triangle having the right angle at  $A$ ; then by Euc. I. 47,


the sq. on  $BC$

=sum of sqq. on  $AC$  and  $AB$ ,

or, more briefly,

$$BC^2 = AC^2 + AB^2.$$

When we use this latter mode of expression it is understood that the sides  $AB$ ,  $AC$ ,  $BC$  are expressed in terms of some common unit, and the above statement may be regarded as a *numerical relation* connecting the numbers of units of length in the three sides of a right-angled triangle.



It is usual to denote the numbers of units of length in the sides opposite the angles  $A$ ,  $B$ ,  $C$  by the letters  $a$ ,  $b$ ,  $c$  respectively. Thus in the above figure we have  $a^2 = b^2 + c^2$ , so that if the lengths of two sides of a right-angled triangle are known, this equation will give the length of the third side.


*Example 1.*  $ABC$  is a right-angled triangle of which  $C$  is the right angle; if  $a=3$ ,  $b=4$ , find  $c$ , and also  $\sin A$  and  $\cot B$ .

$$\text{Here } c^2 = a^2 + b^2 = (3)^2 + (4)^2 = 9 + 16 = 25;$$

$$\therefore c = 5.$$

$$\text{Also } \sin A = \frac{BC}{AB} = \frac{3}{5};$$

$$\cot B = \frac{BC}{AC} = \frac{3}{4}.$$



*Example 2.* A ladder 17 ft. long is placed with its foot at a distance of 8 ft. from the wall of a house and just reaches a window-sill. Find the height of the window-sill, and the sine and tangent of the angle which the ladder makes with the wall.

Let  $AC$  be the ladder, and  $BC$  the wall.


Let  $x$  be the number of feet in  $BC$ ;

$$\text{then } x^2 = (17)^2 - (8)^2 = (17+8)(17-8) = 25 \times 9;$$

$$\therefore x = 5 \times 3 = 15.$$

$$\text{Also } \sin C = \frac{AB}{AC} = \frac{8}{17};$$

$$\tan C = \frac{AB}{BC} = \frac{8}{15}.$$






18. The following important proposition depends upon the property of similar triangles proved in *Euc. vi. 4*. The student who has not read the sixth Book of Euclid should not fail to notice the result arrived at, even if he is unable at this stage to understand the proof.

19. *To prove that the trigonometrical ratios remain unaltered so long as the angle remains the same.*

Let  $\angle AOP$  be any acute angle. In  $OP$  take any points  $B$  and



$D$ , and draw  $BC$  and  $DE$  perpendicular to  $OA$ . Also take any point  $F$  in  $OP$  and draw  $FG$  at right angles to  $OP$ .

$$\text{From the triangle } BOC, \quad \sin POA = \frac{BC}{OB};$$

$$\text{from the triangle } DOE, \quad \sin POA = \frac{DE}{OD};$$

$$\text{from the triangle } FOG, \quad \sin POA = \frac{FG}{OG}.$$

But the triangles  $BOC$ ,  $DOE$ ,  $FOG$  are equiangular ;

$$\therefore \frac{BC}{OB} = \frac{DE}{OD} = \frac{FG}{OG}. \quad [\text{Euc. vi. 4.}]$$

Thus the sine of the angle  $POA$  is the same whether it is obtained from the triangle  $BOC$ , or from the triangle  $DOE$ , or from the triangle  $FOG$ .

A similar proof holds for each of the other trigonometrical ratios. These ratios are therefore independent of the length of the revolving line and depend only on the magnitude of the angle.

20. If  $A$  denote any acute angle, we have proved that all the trigonometrical ratios of  $A$  depend only on the magnitude of the angle  $A$  and not upon the lengths of the lines which bound the angle. It may easily be seen that a change made in the value of  $A$  will produce a consequent change in the values of all the trigonometrical ratios of  $A$ . This point will be discussed more fully in Chap. IX.

DEFINITION. Any expression which involves a variable quantity  $x$ , and whose value is dependent on that of  $x$  is called a **function of  $x$** .

Hence the trigonometrical *ratios* may also be defined as trigonometrical *functions*; for the present we shall chiefly employ the term *ratio*, but in a later part of the subject the idea of ratio is gradually lost and the term *function* becomes more appropriate.

21. The use of the principle proved in Art. 19 is well shewn in the following example, where the trigonometrical ratios are employed as a connecting link between the lines and angles.

*Example.*  $ABC$  is a right-angled triangle of which  $A$  is the right angle.  $BD$  is drawn perpendicular to  $BC$  and meets  $CA$  produced in  $D$ : if  $AB=12$ ,  $AC=16$ ,  $BC=20$ , find  $BD$  and  $CD$ .

From the right-angled triangle  $CBD$ ,

$$\frac{BD}{BC} = \tan C;$$

from the right-angled triangle  $ABC$ ,


$$\frac{AB}{AC} = \tan C;$$

$$\therefore \frac{BD}{BC} = \frac{AB}{AC};$$

$$\therefore \frac{BD}{20} = \frac{12}{16}; \text{ whence } BD = 15.$$

Again, 
$$\frac{CD}{CB} = \sec C = \frac{BC}{CA};$$

$$\therefore \frac{CD}{20} = \frac{20}{16}; \text{ whence } CD = 25.$$



The same results can be obtained by the help of Euc. VI. 8.

## EXAMPLES. II.

1. The sides  $AB$ ,  $BC$ ,  $CA$  of a right-angled triangle are 17, 15, 8 respectively; write down the values of  $\sin A$ ,  $\sec A$ ,  $\tan B$ ,  $\sec B$ .

2. The sides  $PQ$ ,  $QR$ ,  $RP$  of a right-angled triangle are 13, 5, 12 respectively: write down the values of  $\cot P$ ,  $\operatorname{cosec} Q$ ,  $\cos Q$ ,  $\cos P$ .

3.  $ABC$  is a triangle in which  $A$  is a right angle; if  $b=15$ ,  $c=20$ , find  $\alpha$ ,  $\sin C$ ,  $\cos B$ ,  $\cot C$ ,  $\sec C$ .

4.  $ABC$  is a triangle in which  $B$  is a right angle; if  $a=24$ ,  $b=25$ , find  $c$ ,  $\sin C$ ,  $\tan A$ ,  $\operatorname{cosec} A$ .

5. The sides  $ED$ ,  $EF$ ,  $DF$  of a right-angled triangle are 35, 37, 12 respectively: write down the values of  $\sec E$ ,  $\sec F$ ,  $\cot E$ ,  $\sin F$ .

6. The hypotenuse of a right-angled triangle is 15 inches, and one of the sides is 9 inches: find the third side and the sine, cosine and tangent of the angle opposite to it.

7. Find the hypotenuse  $AB$  of a right-angled triangle in which  $AC=7$ ,  $BC=24$ . Write down the sine and cosine of  $A$ , and shew that the sum of their squares is equal to 1.

8. A ladder 41 ft. long is placed with its foot at a distance of 9 ft. from the wall of a house and just reaches a window-sill. Find the height of the window-sill, and the sine and cotangent of the angle which the ladder makes with the ground.

9. A ladder is 29 ft. long; how far must its foot be placed from a wall so that the ladder may just reach the top of the wall which is 21 ft. from the ground? Write down all the trigonometrical ratios of the angle between the ladder and the wall.

10.  $ABCD$  is a square;  $C$  is joined to  $E$ , the middle point of  $AD$ : find all the trigonometrical ratios of the angle  $ECD$ .

11.  $ABCD$  is a quadrilateral in which the diagonal  $AC$  is at right angles to each of the sides  $AB$ ,  $CD$ : if  $AB=15$ ,  $AC=36$ ,  $AD=85$ , find  $\sin ABC$ ,  $\sec ACB$ ,  $\cos CDA$ ,  $\operatorname{cosec} DAC$ .

12.  $PQRS$  is a quadrilateral in which the angle  $PSR$  is a right angle. If the diagonal  $PR$  is at right angles to  $RQ$ , and  $RP=20$ ,  $RQ=21$ ,  $RS=16$ , find  $\sin PRS$ ,  $\tan RPS$ ,  $\cos RPQ$ ,  $\operatorname{cosec} PQR$ .

## CHAPTER III.

### RELATIONS BETWEEN THE TRIGONOMETRICAL RATIOS.

#### 22. Reciprocal relations between certain ratios.

(1) Let  $ABC$  be a triangle, right-angled at  $C$ ;

then  $\sin A = \frac{BC}{AB} = \frac{a}{c},$


and  $\operatorname{cosec} A = \frac{AB}{BC} = \frac{c}{a};$

$$\therefore \sin A \times \operatorname{cosec} A = \frac{a}{c} \times \frac{c}{a} = 1.$$

Thus  $\sin A$  and  $\operatorname{cosec} A$  are reciprocals ;

$$\therefore \sin A = \frac{1}{\operatorname{cosec} A},$$

and  $\operatorname{cosec} A = \frac{1}{\sin A}.$



(2) Again,

$$\cos A = \frac{AC}{AB} = \frac{b}{c}, \text{ and } \sec A = \frac{AB}{AC} = \frac{c}{b};$$

$$\therefore \cos A \times \sec A = \frac{b}{c} \times \frac{c}{b} = 1;$$

$$\therefore \cos A = \frac{1}{\sec A}, \text{ and } \sec A = \frac{1}{\cos A}.$$

(3) Also

$$\tan A = \frac{BC}{AC} = \frac{a}{b}, \text{ and } \cot A = \frac{AC}{BC} = \frac{b}{a};$$

$$\therefore \tan A \times \cot A = \frac{a}{b} \times \frac{b}{a} = 1;$$

$$\therefore \tan A = \frac{1}{\cot A}, \text{ and } \cot A = \frac{1}{\tan A}.$$

23. To express  $\tan A$  and  $\cot A$  in terms of  $\sin A$  and  $\cos A$ .


From the adjoining figure we have

$$\begin{aligned}\tan A &= \frac{BC}{AC} = \frac{a}{b} = \frac{a}{c} \div \frac{b}{c} \\ &= \sin A \div \cos A ;\end{aligned}$$

$$\therefore \tan A = \frac{\sin A}{\cos A}.$$

$$\begin{aligned}\text{Again, } \cot A &= \frac{AC}{BC} = \frac{b}{a} = \frac{b}{c} \div \frac{a}{c} \\ &= \cos A \div \sin A ;\end{aligned}$$

$$\therefore \cot A = \frac{\cos A}{\sin A} ;$$



which is also evident from the reciprocal relation  $\cot A = \frac{1}{\tan A}$ .

*Example.* Prove that  $\operatorname{cosec} A \tan A = \sec A$ .

$$\begin{aligned}\operatorname{cosec} A \tan A &= \frac{1}{\sin A} \times \frac{\sin A}{\cos A} = \frac{1}{\cos A} \\ &= \sec A.\end{aligned}$$

24. We frequently meet with expressions which involve the square and other powers of the trigonometrical ratios, such as  $(\sin A)^2$ ,  $(\tan A)^3$ , ... It is usual to write these in the shorter forms  $\sin^2 A$ ,  $\tan^3 A$ , ...

$$\begin{aligned}\text{Thus } \tan^2 A &= (\tan A)^2 = \left( \frac{\sin A}{\cos A} \right)^2 \\ &= \frac{(\sin A)^2}{(\cos A)^2} = \frac{\sin^2 A}{\cos^2 A}.\end{aligned}$$


*Example.* Shew that  $\sin^2 A \sec A \cot^2 A = \cos A$ .

$$\begin{aligned}\sin^2 A \sec A \cot^2 A &= \sin^2 A \times \frac{1}{\cos A} \times \left( \frac{\cos A}{\sin A} \right)^2 \\ &= \sin^2 A \times \frac{1}{\cos A} \times \frac{\cos^2 A}{\sin^2 A} \\ &= \cos A,\end{aligned}$$

by cancelling factors common to numerator and denominator.

25. To prove that  $\sin^2 A + \cos^2 A = 1$ .

Let  $BAC$  be any acute angle; draw  $BC$  perpendicular to



$AC$ , and denote the sides of the right-angled triangle  $ABC$  by  $a, b, c$ .

By definition,  $\sin A = \frac{BC}{AB} = \frac{a}{c}$ ;

and  $\cos A = \frac{AC}{AB} = \frac{b}{c}$ ;

$$\begin{aligned}\therefore \sin^2 A + \cos^2 A &= \frac{a^2}{c^2} + \frac{b^2}{c^2} = \frac{a^2 + b^2}{c^2} \\ &= \frac{c^2}{c^2} \\ &= 1.\end{aligned}$$

COR.  $\sin^2 A = 1 - \cos^2 A$ ,  $\sin A = \sqrt{1 - \cos^2 A}$ ;  
 $\cos^2 A = 1 - \sin^2 A$ ,  $\cos A = \sqrt{1 - \sin^2 A}$ .

*Example 1.* Prove that  $\cos^4 A - \sin^4 A = \cos^2 A - \sin^2 A$ .

$$\begin{aligned}\cos^4 A - \sin^4 A &= (\cos^2 A + \sin^2 A)(\cos^2 A - \sin^2 A) \\ &= \cos^2 A - \sin^2 A,\end{aligned}$$

since the first factor is equal to 1.

*Example 2.* Prove that  $\cot a \sqrt{1 - \cos^2 a} = \cos a$ .

$$\begin{aligned}\cot a \sqrt{1 - \cos^2 a} &= \cot a \times \sin a \\ &= \frac{\cos a}{\sin a} \times \sin a = \cos a.\end{aligned}$$

26. To prove that  $\sec^2 A = 1 + \tan^2 A$ .

With the figure of the previous article, we have

$$\sec A = \frac{AB}{AC} = \frac{c}{b};$$

$$\begin{aligned}\therefore \sec^2 A &= \frac{c^2}{b^2} = \frac{b^2 + a^2}{b^2} \\ &= 1 + \frac{a^2}{b^2} \\ &= 1 + \tan^2 A.\end{aligned}$$

$$\begin{aligned}\text{COR.} \quad \sec^2 A - \tan^2 A &= 1, & \sec A &= \sqrt{1 + \tan^2 A}, \\ \tan^2 A &= \sec^2 A - 1, & \tan A &= \sqrt{\sec^2 A - 1}.\end{aligned}$$

*Example.* Prove that  $\cos A \sqrt{\sec^2 A - 1} = \sin A$ .

$$\begin{aligned}\cos A \sqrt{\sec^2 A - 1} &= \cos A \times \tan A \\ &= \cos A \times \frac{\sin A}{\cos A} \\ &= \sin A.\end{aligned}$$

27. To prove that  $\operatorname{cosec}^2 A = 1 + \cot^2 A$ .

With the figure of Art. 25, we have

$$\operatorname{cosec} A = \frac{AB}{BC} = \frac{c}{a};$$

$$\begin{aligned}\therefore \operatorname{cosec}^2 A &= \frac{c^2}{a^2} = \frac{a^2 + b^2}{a^2} \\ &= 1 + \frac{b^2}{a^2} \\ &= 1 + \cot^2 A.\end{aligned}$$

$$\begin{aligned}\text{COR.} \quad \operatorname{cosec}^2 A - \cot^2 A &= 1, & \operatorname{cosec} A &= \sqrt{1 + \cot^2 A}, \\ \cot^2 A &= \operatorname{cosec}^2 A - 1, & \cot A &= \sqrt{\operatorname{cosec}^2 A - 1}.\end{aligned}$$

*Example.* Prove that  $\cot^4 a - 1 = \operatorname{cosec}^4 a - 2 \operatorname{cosec}^2 a$ .

$$\begin{aligned}\cot^4 a - 1 &= (\cot^2 a + 1)(\cot^2 a - 1) \\ &= \operatorname{cosec}^2 a (\operatorname{cosec}^2 a - 1 - 1) \\ &= \operatorname{cosec}^2 a (\operatorname{cosec}^2 a - 2) \\ &= \operatorname{cosec}^4 a - 2 \operatorname{cosec}^2 a.\end{aligned}$$



28. The formulæ proved in the last three articles are not independent, for they are merely different ways of expressing in trigonometrical symbols the property of a right-angled triangle proved in Euc. I. 47.

29. It will be useful here to collect the formulæ proved in this chapter.

$$\text{I. } \operatorname{cosec} A \times \sin A = 1, \quad \operatorname{cosec} A = \frac{1}{\sin A}, \quad \sin A = \frac{1}{\operatorname{cosec} A};$$

$$\sec A \times \cos A = 1, \quad \sec A = \frac{1}{\cos A}, \quad \cos A = \frac{1}{\sec A};$$

$$\cot A \times \tan A = 1, \quad \cot A = \frac{1}{\tan A}, \quad \tan A = \frac{1}{\cot A}.$$

$$\text{II. } \tan A = \frac{\sin A}{\cos A}, \quad \cot A = \frac{\cos A}{\sin A}.$$

$$\begin{aligned} \text{III. } \quad \sin^2 A + \cos^2 A &= 1, \\ \sec^2 A &= 1 + \tan^2 A, \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A. \end{aligned}$$

### Easy Identities.

30. We shall now exemplify the use of these fundamental formulæ in proving *identities*. An identity asserts that two expressions are always equal, and the proof of this equality is called "proving the identity." Some easy illustrations have already been given in this chapter. The general method of procedure is to choose one of the expressions given (usually the more complicated of the two) and to shew by successive transformations that it can be made to assume the form of the other.

*Example 1.* Prove that  $\sin^3 A \cot^3 A + \cos^2 A \tan^2 A = 1$ .

Here it will be found convenient to express all the trigonometrical ratios in terms of the sine and cosine.

$$\begin{aligned} \text{The first side} &= \sin^2 A \cdot \frac{\cos^2 A}{\sin^2 A} + \cos^2 A \cdot \frac{\sin^2 A}{\cos^2 A} \\ &= \cos^2 A + \sin^2 A \\ &= 1. \end{aligned}$$



*Example 2.* Prove that  $\sec^4 \theta - \sec^2 \theta = \tan^2 \theta + \tan^4 \theta$ .

The form of this identity at once suggests that we should use the secant-tangent formula of Art. 26; hence

$$\begin{aligned}\text{the first side} &= \sec^2 \theta (\sec^2 \theta - 1) \\ &= (1 + \tan^2 \theta) \tan^2 \theta \\ &= \tan^2 \theta + \tan^4 \theta.\end{aligned}$$

### EXAMPLES. III. a.

Prove the following identities :

1.  $\sin A \cot A = \cos A$ .
2.  $\cos A \tan A = \sin A$ .
3.  $\cot A \sec A = \operatorname{cosec} A$ .
4.  $\sin A \sec A = \tan A$ .
5.  $\cos A \operatorname{cosec} A = \cot A$ .
6.  $\cot A \sec A \sin A = 1$ .
7.  $(1 - \cos^2 A) \operatorname{cosec}^2 A = 1$ .
8.  $(1 - \sin^2 A) \sec^2 A = 1$ .
9.  $\cot^2 \theta (1 - \cos^2 \theta) = \cos^2 \theta$ .
10.  $(1 - \cos^2 \theta) \sec^2 \theta = \tan^2 \theta$ .
11.  $\tan a \sqrt{1 - \sin^2 a} = \sin a$ .
12.  $\operatorname{cosec} a \sqrt{1 - \sin^2 a} = \cot a$ .
13.  $(1 + \tan^2 A) \cos^2 A = 1$ .
14.  $(\sec^2 A - 1) \cot^2 A = 1$ .
15.  $(1 - \cos^2 \theta) (1 + \tan^2 \theta) = \tan^2 \theta$ .
16.  $\cos a \operatorname{cosec} a \sqrt{\sec^2 a - 1} = 1$ .
17.  $\sin^2 A (1 + \cot^2 A) = 1$ .
18.  $(\operatorname{cosec}^2 A - 1) \tan^2 A = 1$ .
19.  $(1 - \cos^2 A) (1 + \cot^2 A) = 1$ .
20.  $\sin a \sec a \sqrt{\operatorname{cosec}^2 a - 1} = 1$ .
21.  $\cos a \sqrt{\cot^2 a + 1} = \sqrt{\operatorname{cosec}^2 a - 1}$ .
22.  $\sin^2 \theta \cot^2 \theta + \sin^2 \theta = 1$ .
23.  $(1 + \tan^2 \theta) (1 - \sin^2 \theta) = 1$ .
24.  $\sin^2 \theta \sec^2 \theta = \sec^2 \theta - 1$ .
25.  $\operatorname{cosec}^2 \theta \tan^2 \theta - 1 = \tan^2 \theta$ .

Prove the following identities :

$$26. \quad \frac{1}{\sec^2 A} + \frac{1}{\operatorname{cosec}^2 A} = 1.$$

$$27. \quad \frac{1}{\cos^2 A} - \frac{1}{\cot^2 A} = 1.$$

$$28. \quad \frac{\sin A}{\operatorname{cosec} A} + \frac{\cos A}{\sec A} = 1.$$

$$29. \quad \frac{\sec A}{\cos A} - \frac{\tan A}{\cot A} = 1.$$

$$30. \quad \sin^4 a - \cos^4 a = 2 \sin^2 a - 1 = 1 - 2 \cos^2 a.$$

$$31. \quad \sec^4 a - 1 = 2 \tan^2 a + \tan^4 a.$$

$$32. \quad \operatorname{cosec}^4 a - 1 = 2 \cot^2 a + \cot^4 a.$$

$$33. \quad (\tan a \operatorname{cosec} a)^2 - (\sin a \sec a)^2 = 1.$$

$$34. \quad (\sec \theta \cot \theta)^2 - (\cos \theta \operatorname{cosec} \theta)^2 = 1.$$

$$35. \quad \tan^2 \theta - \cot^2 \theta = \sec^2 \theta - \operatorname{cosec}^2 \theta.$$

31. The foregoing examples have required little more than a direct application of the fundamental formulæ; we shall now give some identities offering a greater variety of treatment.

*Example 1.* Prove that  $\sec^2 A + \operatorname{cosec}^2 A = \sec^2 A \operatorname{cosec}^2 A$ .

$$\begin{aligned} \text{The first side} &= \frac{1}{\cos^2 A} + \frac{1}{\sin^2 A} = \frac{\sin^2 A + \cos^2 A}{\cos^2 A \sin^2 A} \\ &= \frac{1}{\cos^2 A \sin^2 A} = \sec^2 A \operatorname{cosec}^2 A. \end{aligned}$$

Occasionally it is found convenient to prove the equality of the two expressions by reducing each to the same form.

*Example 2.* Prove that

$$\sin^2 A \tan A + \cos^2 A \cot A + 2 \sin A \cos A = \tan A + \cot A.$$

$$\begin{aligned} \text{The first side} &= \sin^2 A \cdot \frac{\sin A}{\cos A} + \cos^2 A \cdot \frac{\cos A}{\sin A} + 2 \sin A \cos A \\ &= \frac{\sin^4 A + \cos^4 A + 2 \sin^2 A \cos^2 A}{\sin A \cos A} \\ &= \frac{(\sin^2 A + \cos^2 A)^2}{\sin A \cos A} = \frac{1}{\sin A \cos A}. \end{aligned}$$

$$\begin{aligned} \text{The second side} &= \frac{\sin A}{\cos A} + \frac{\cos A}{\sin A} = \frac{\sin^2 A + \cos^2 A}{\cos A \sin A} \\ &= \frac{1}{\sin A \cos A}. \end{aligned}$$

$$\text{Thus each side of the identity} = \frac{1}{\sin A \cos A}.$$

*Example 3.* Prove that  $\frac{\tan \alpha - \cot \beta}{\tan \beta - \cot \alpha} = \tan \alpha \cot \beta$ .

$$\begin{aligned} \text{The first side} &= \frac{\tan \alpha - \cot \beta}{\frac{1}{\cot \beta} - \frac{1}{\tan \alpha}} = \frac{\tan \alpha - \cot \beta}{\frac{\tan \alpha - \cot \beta}{\tan \alpha \cot \beta}} \\ &= \frac{\tan \alpha - \cot \beta}{1} \times \frac{\tan \alpha \cot \beta}{\tan \alpha - \cot \beta} \\ &= \tan \alpha \cot \beta. \end{aligned}$$

The transformations in the successive steps are usually suggested by the form into which we wish to bring the result. For instance, in this last example we might have proved the identity by substituting for the tangent and cotangent in terms of the sine and cosine. This however is not the best method, for the form in which the right-hand side is given suggests that we should retain  $\tan \alpha$  and  $\cot \beta$  unchanged throughout the work.

### EXAMPLES. III. b.

Prove the following identities :

1.  $\frac{\sin a \cot^2 a}{\cos a} = \frac{1}{\tan a}.$
2.  $\frac{\sec^2 a \cot a}{\operatorname{cosec}^2 a} = \tan a.$
3.  $1 - \operatorname{vers} \theta = \sin \theta \cot \theta.$
4.  $\operatorname{vers} \theta \sec \theta = \sec \theta - 1.$
5.  $\sec \theta - \tan \theta \sin \theta = \cos \theta.$
6.  $\tan \theta + \cot \theta = \sec \theta \operatorname{cosec} \theta.$
7.  $\sqrt{1 + \cot^2 A} \cdot \sqrt{\sec^2 A - 1} \cdot \sqrt{1 - \sin^2 A} = 1.$
8.  $(\cos \theta + \sin \theta)^2 + (\cos \theta - \sin \theta)^2 = 2.$
9.  $(1 + \tan \theta)^2 + (1 - \tan \theta)^2 = 2 \sec^2 \theta.$
10.  $(\cot \theta - 1)^2 + (\cot \theta + 1)^2 = 2 \operatorname{cosec}^2 \theta.$
11.  $\sin^2 A (1 + \cot^2 A) + \cos^2 A (1 + \tan^2 A) = 2.$
12.  $\cos^2 A (\sec^2 A - \tan^2 A) + \sin^2 A (\operatorname{cosec}^2 A - \cot^2 A) = 1.$
13.  $\cot^2 a + \cot^4 a = \operatorname{cosec}^4 a - \operatorname{cosec}^2 a.$
14.  $\frac{\tan^2 a}{1 + \tan^2 a} \cdot \frac{1 + \cot^2 a}{\cot^2 a} = \sin^2 a \sec^2 a.$
15.  $\frac{1}{1 - \sin a} + \frac{1}{1 + \sin a} = 2 \sec^2 a.$

Prove the following identities :

$$16. \quad \frac{\tan a}{\sec a - 1} + \frac{\tan a}{\sec a + 1} = 2 \operatorname{cosec} a.$$

$$17. \quad \frac{1}{1 + \sin^2 a} + \frac{1}{1 + \operatorname{cosec}^2 a} = 1.$$

$$18. \quad (\sec \theta + \operatorname{cosec} \theta)(\sin \theta + \cos \theta) = \sec \theta \operatorname{cosec} \theta + 2.$$

$$19. \quad (\cos \theta - \sin \theta)(\operatorname{cosec} \theta - \sec \theta) = \sec \theta \operatorname{cosec} \theta - 2.$$

$$20. \quad (1 + \cot \theta + \operatorname{cosec} \theta)(1 + \cot \theta - \operatorname{cosec} \theta) = 2 \cot \theta.$$

$$21. \quad (\sec \theta + \tan \theta - 1)(\sec \theta - \tan \theta + 1) = 2 \tan \theta.$$

$$22. \quad (\sin A + \operatorname{cosec} A)^2 + (\cos A + \sec A)^2 = \tan^2 A + \cot^2 A + 7.$$

$$23. \quad (\sec^2 A + \tan^2 A)(\operatorname{cosec}^2 A + \cot^2 A) = 1 + 2 \sec^2 A \operatorname{cosec}^2 A.$$

$$24. \quad (1 - \sin A + \cos A)^2 = 2(1 - \sin A)(1 + \cos A).$$

$$25. \quad \sin A(1 + \tan A) + \cos A(1 + \cot A) = \sec A + \operatorname{cosec} A.$$

$$26. \quad \cos \theta(\tan \theta + 2)(2 \tan \theta + 1) = 2 \sec \theta + 5 \sin \theta.$$

$$27. \quad (\tan \theta + \sec \theta)^2 = \frac{1 + \sin \theta}{1 - \sin \theta}.$$

$$28. \quad \frac{2 \sin \theta \cos \theta - \cos \theta}{1 - \sin \theta + \sin^2 \theta - \cos^2 \theta} = \cot \theta.$$

$$29. \quad \cot^2 \theta \cdot \frac{\sec \theta - 1}{1 + \sin \theta} + \sec^2 \theta \cdot \frac{\sin \theta - 1}{1 + \sec \theta} = 0.$$

[The following examples contain functions of two angles; in each case the two angles are quite independent of each other.]

$$30. \quad \tan^2 a + \sec^2 \beta = \sec^2 a + \tan^2 \beta.$$

$$31. \quad \frac{\tan a + \cot \beta}{\cot a + \tan \beta} = \frac{\tan a}{\tan \beta}. \quad 32. \quad \frac{\tan a - \cot \beta}{\cot a - \tan \beta} = -\frac{\cot \beta}{\cot a}.$$

$$33. \quad \cot a \tan \beta (\tan a + \cot \beta) = \cot a + \tan \beta.$$

$$34. \quad \sin^2 a \cos^2 \beta - \cos^2 a \sin^2 \beta = \sin^2 a - \sin^2 \beta.$$

$$35. \quad \sec^2 a \tan^2 \beta - \tan^2 a \sec^2 \beta = \tan^2 \beta - \tan^2 a.$$

$$36. \quad (\sin a \cos \beta + \cos a \sin \beta)^2 + (\cos a \cos \beta - \sin a \sin \beta)^2 = 1.$$

32. By means of the relations collected together in Art. 29, all the trigonometrical ratios can be expressed in terms of any one.

*Example 1.* Express all the trigonometrical ratios of  $A$  in terms of  $\tan A$ .

$$\text{We have } \cot A = \frac{1}{\tan A};$$

$$\sec A = \sqrt{1 + \tan^2 A};$$

$$\cos A = \frac{1}{\sec A} = \frac{1}{\sqrt{1 + \tan^2 A}};$$

$$\sin A = \frac{\sin A}{\cos A} \cos A = \tan A \cos A = \frac{\tan A}{\sqrt{1 + \tan^2 A}};$$

$$\operatorname{cosec} A = \frac{1}{\sin A} = \frac{\sqrt{1 + \tan^2 A}}{\tan A}.$$

Obs. In writing down the ratios we choose the simplest and most natural order. For instance,  $\cot A$  is obtained at once by the *reciprocal relation* connecting the tangent and cotangent:  $\sec A$  comes immediately from the tangent-secant formula; the remaining three ratios now readily follow.

*Example 2.* Given  $\cos A = \frac{5}{13}$ , find  $\operatorname{cosec} A$  and  $\cot A$ .

$$\begin{aligned} \operatorname{cosec} A &= \frac{1}{\sin A} = \frac{1}{\sqrt{1 - \cos^2 A}} \\ &= \frac{1}{\sqrt{1 - \left(\frac{5}{13}\right)^2}} = \frac{1}{\sqrt{1 - \frac{25}{169}}} = \frac{1}{\sqrt{\frac{144}{169}}} = \frac{1}{\frac{12}{13}} = \frac{13}{12}. \end{aligned}$$

$$\begin{aligned} \cot A &= \frac{\cos A}{\sin A} = \cos A \times \operatorname{cosec} A \\ &= \frac{5}{13} \times \frac{13}{12} = \frac{5}{12}. \end{aligned}$$

33. *It is always possible to describe a right-angled triangle when two sides are given:* for the third side can be found by Euc. I. 47, and the construction can then be effected by Euc. I. 22. We can thus readily obtain all the trigonometrical ratios when one is given, or express all in terms of any one.

*Example 1.* Given  $\cos A = \frac{5}{13}$ , find  $\operatorname{cosec} A$  and  $\cot A$ .

Take a right-angled triangle  $PQR$ , of which  $Q$  is the right angle, having the hypotenuse  $PR = 13$  units, and  $PQ = 5$  units.

Let  $QR = x$  units; then


$$\begin{aligned} x^2 &= (13)^2 - (5)^2 = (13 + 5)(13 - 5) \\ &= 18 \times 8 = 9 \times 2 \times 8; \\ \therefore x &= 3 \times 4 = 12. \end{aligned}$$

Now  $\cos RPQ = \frac{PQ}{PR} = \frac{5}{13},$

so that  $\angle RPQ = A.$

Hence  $\operatorname{cosec} A = \frac{PR}{QR} = \frac{13}{12},$

and  $\cot A = \frac{PQ}{QR} = \frac{5}{12}. \quad [\text{Compare Art. 32, Ex. 2.}]$




*Example 2.* Find  $\tan A$  and  $\cos A$  in terms of  $\operatorname{cosec} A$ .

Take a triangle  $PQR$  right-angled at  $Q$ , and having  $\angle RPQ = A$ . For shortness, denote  $\operatorname{cosec} A$  by  $c$ .

Then  $\operatorname{cosec} A = c = \frac{c}{1};$

but  $\operatorname{cosec} A = \frac{PR}{QR};$   
 $\therefore \frac{PR}{QR} = \frac{c}{1}.$



Let  $QR$  be taken as the unit of measurement;

then  $QR = 1$ , and therefore  $PR = c$ .

Let  $PQ$  contain  $x$  units; then

$$x^2 = c^2 - 1, \text{ so that } x = \sqrt{c^2 - 1}.$$

Hence  $\tan A = \frac{QR}{PQ} = \frac{1}{\sqrt{c^2 - 1}} = \frac{1}{\sqrt{\operatorname{cosec}^2 A - 1}},$

and  $\cos A = \frac{PQ}{PR} = \frac{\sqrt{c^2 - 1}}{c} = \frac{\sqrt{\operatorname{cosec}^2 A - 1}}{\operatorname{cosec} A}.$

## EXAMPLES. III. c.

1. Given  $\sin A = \frac{1}{2}$ , find  $\sec A$  and  $\cot A$ .
2. Given  $\tan A = \frac{4}{3}$ , find  $\sin A$  and  $\cos A$ .
3. Find  $\cot \theta$  and  $\sin \theta$  when  $\sec \theta = 4$ .
4. If  $\tan a = \frac{1}{2}$ , find  $\sec a$  and  $\operatorname{cosec} a$ .
5. Find the sine and cotangent of an angle whose secant is 7.
6. If  $25 \sin A = 7$ , find  $\tan A$  and  $\sec A$ .
7. Express  $\sin A$  and  $\tan A$  in terms of  $\cos A$ .
8. Express  $\operatorname{cosec} a$  and  $\cos a$  in terms of  $\cot a$ .
9. Find  $\sin \theta$  and  $\cot \theta$  in terms of  $\sec \theta$ .
10. Express all the trigonometrical ratios of  $A$  in terms of  $\sin A$ .
11. Given  $\sin A - \cos A = 0$ , find  $\operatorname{cosec} A$ .
12. If  $\sin A = \frac{m}{n}$ , prove that  $\sqrt{n^2 - m^2} \cdot \tan A = m$ .
13. If  $p \cot \theta = \sqrt{q^2 - p^2}$ , find  $\sin \theta$ .
14. When  $\sec A = \frac{m^2 + 1}{2m}$ , find  $\tan A$  and  $\sin A$ .
15. Given  $\tan A = \frac{2pq}{p^2 - q^2}$ , find  $\cos A$  and  $\operatorname{cosec} A$ .
16. If  $\sec a = \frac{13}{5}$ , find the value of  $\frac{2 \sin a - 3 \cos a}{4 \sin a - 9 \cos a}$ .
17. If  $\cot \theta = \frac{p}{q}$ , find the value of  $\frac{p \cos \theta - q \sin \theta}{p \cos \theta + q \sin \theta}$ .




## CHAPTER IV.

### TRIGONOMETRICAL RATIOS OF CERTAIN ANGLES.

#### 34. Trigonometrical Ratios of $45^\circ$ .

Let  $BAC$  be a right-angled isosceles triangle, with the right angle at  $C$ ; so that  $B = A = 45^\circ$ .



Let each of the equal sides contain  $l$  units,  
then  $AC = BC = l$ .

Also  $AB^2 = l^2 + l^2 = 2l^2$ ;

$$\therefore AB = l\sqrt{2}.$$

$$\therefore \sin 45^\circ = \frac{BC}{AB} = \frac{l}{l\sqrt{2}} = \frac{1}{\sqrt{2}};$$

$$\cos 45^\circ = \frac{AC}{AB} = \frac{l}{l\sqrt{2}} = \frac{1}{\sqrt{2}};$$

$$\tan 45^\circ = \frac{BC}{AC} = \frac{l}{l} = 1.$$


The other three ratios are the reciprocals of these; thus

$$\operatorname{cosec} 45^\circ = \sqrt{2}, \quad \sec 45^\circ = \sqrt{2}, \quad \cot 45^\circ = 1;$$

or they may be read off from the figure.

### 35. Trigonometrical Ratios of $60^\circ$ and $30^\circ$ .

Let  $ABC$  be an equilateral triangle; thus each of its angles is  $60^\circ$ .



Bisect  $\angle BAC$  by  $AD$  meeting  $BC$  at  $D$ ; then  $\angle BAD = 30^\circ$ .

By Euc. I. 4, the triangles  $ABD$ ,  $ACD$  are equal in all respects; therefore  $BD = CD$ , and the angles at  $D$  are right angles.

In the right-angled triangle  $ADB$ , let  $BD = l$ ; then

$$AB = BC = 2l;$$

$$\therefore AD^2 = 4l^2 - l^2 = 3l^2;$$

$$\therefore AD = l\sqrt{3}.$$

$$\therefore \sin 60^\circ = \frac{AD}{AB} = \frac{l\sqrt{3}}{2l} = \frac{\sqrt{3}}{2};$$

$$\cos 60^\circ = \frac{BD}{AB} = \frac{l}{2l} = \frac{1}{2};$$

$$\tan 60^\circ = \frac{AD}{BD} = \frac{l\sqrt{3}}{l} = \sqrt{3}.$$

Again,

$$\sin 30^\circ = \frac{BD}{AB} = \frac{l}{2l} = \frac{1}{2};$$


$$\cos 30^\circ = \frac{AD}{AB} = \frac{l\sqrt{3}}{2l} = \frac{\sqrt{3}}{2};$$

$$\tan 30^\circ = \frac{BD}{AD} = \frac{l}{l\sqrt{3}} = \frac{1}{\sqrt{3}}.$$

The other ratios may be read off from the figure.

36. The trigonometrical ratios of  $45^\circ$ ,  $60^\circ$ ,  $30^\circ$  occur very frequently; it is therefore important that the student should be able to quote readily their numerical values. The exercise which follows will furnish useful practice.

At first it will probably be found safer to make use of the accompanying diagrams than to trust to the memory.



The trigonometrical ratios of  $45^\circ$  can be read off from Fig. 1; those of  $60^\circ$  and  $30^\circ$  from Fig. 2.

*Example 1.* Find the values of  $\sec^3 45^\circ$  and  $\sin 60^\circ \cot 30^\circ \tan 45^\circ$ .

$$\sec^3 45^\circ = (\sec 45^\circ)^3 = (\sqrt{2})^3 = \sqrt{2} \times \sqrt{2} \times \sqrt{2} = 2\sqrt{2}.$$

$$\sin 60^\circ \cot 30^\circ \tan 45^\circ = \frac{\sqrt{3}}{2} \times \sqrt{3} \times 1 = \frac{3}{2}.$$

*Example 2.* Find the value of

$$2 \cot 45^\circ + \cos^3 60^\circ - 2 \sin^4 60^\circ + \frac{3}{4} \tan^2 30^\circ.$$

$$\begin{aligned} \text{The value} &= (2 \times 1) + \left(\frac{1}{2}\right)^3 - 2 \left(\frac{\sqrt{3}}{2}\right)^4 + \frac{3}{4} \left(\frac{1}{\sqrt{3}}\right)^2 \\ &= 2 + \frac{1}{8} - 2 \left(\frac{3}{4}\right)^2 + \frac{3}{4} \left(\frac{1}{3}\right) \\ &= 2 + \frac{1}{8} - \frac{9}{8} + \frac{1}{4} = 1\frac{1}{4}. \end{aligned}$$

### EXAMPLES. IV. a.

Find the numerical value of

- |  |  |
|--|--|
| 1. $\tan^2 60^\circ + 2 \tan^2 45^\circ$ .                   | 2. $\tan^3 45^\circ + 4 \cos^3 60^\circ$ .           |
| 3. $2 \operatorname{cosec}^2 45^\circ - 3 \sec^2 30^\circ$ . | 4. $\cot 60^\circ \tan 30^\circ + \sec^2 45^\circ$ . |

5.  $2 \sin 30^\circ \cos 30^\circ \cot 60^\circ$ .
6.  $\tan^2 45^\circ \sin 60^\circ \tan 30^\circ \tan^2 60^\circ$ .
7.  $\tan^2 60^\circ + 4 \cos^2 45^\circ + 3 \sec^2 30^\circ$ .
8.  $\frac{1}{2} \operatorname{cosec}^2 60^\circ + \sec^2 45^\circ - 2 \cot^2 60^\circ$ .
9.  $\tan^2 30^\circ + 2 \sin 60^\circ + \tan 45^\circ - \tan 60^\circ + \cos^2 30^\circ$ .
10.  $\cot^2 45^\circ + \cos 60^\circ - \sin^2 60^\circ - \frac{3}{4} \cot^2 60^\circ$ .
11.  $3 \tan^2 30^\circ + \frac{4}{3} \cos^2 30^\circ - \frac{1}{2} \sec^2 45^\circ - \frac{1}{3} \sin^2 60^\circ$ .
12.  $\cos 60^\circ - \tan^2 45^\circ + \frac{3}{4} \tan^2 30^\circ + \cos^2 30^\circ - \sin 30^\circ$ .
13.  $\frac{1}{3} \sin^2 60^\circ - \frac{1}{2} \sec 60^\circ \tan^2 30^\circ + \frac{4}{3} \sin^2 45^\circ \tan^2 60^\circ$ .
14. If  $\tan^2 45^\circ - \cos^2 60^\circ = x \sin 45^\circ \cos 45^\circ \tan 60^\circ$ , find  $x$ .
15. Find  $x$  from the equation

$$x \sin 30^\circ \cos^2 45^\circ = \frac{\cot^2 30^\circ \sec 60^\circ \tan 45^\circ}{\operatorname{cosec}^2 45^\circ \operatorname{cosec} 30^\circ}.$$

37. DEFINITION. The **complement** of an angle is its *defect* from a right angle.

Two angles are said to be **complementary** when their sum is a right angle.

Thus in every right-angled triangle, each acute angle is the complement of the other. For in the figure of the next article, if  $B$  is the right angle, the sum of  $A$  and  $C$  is  $90^\circ$ .


$$\therefore C = 90^\circ - A, \text{ and } A = 90^\circ - C.$$

### Trigonometrical Ratios of Complementary Angles.

38. Let  $ABC$  be a right-angled triangle, of which  $B$  is the right angle; then the angles at  $A$  and  $C$  are complementary; so that  $C = 90^\circ - A$ .

$$\therefore \sin(90^\circ - A) = \sin C = \frac{AB}{AC} = \cos A;$$

$$\text{and } \cos(90^\circ - A) = \cos C = \frac{BC}{AC} = \sin A.$$



Similarly, it may be proved that

$$\left. \begin{array}{l} \tan(90^\circ - A) = \cot A, \\ \cot(90^\circ - A) = \tan A; \end{array} \right\} \text{ and } \left. \begin{array}{l} \sec(90^\circ - A) = \operatorname{cosec} A, \\ \operatorname{cosec}(90^\circ - A) = \sec A. \end{array} \right\}$$

39. If we define the **co-sine**, **co-tangent**, **co-secant**, as the **co-functions** of the angle, the foregoing results may be embodied in a single statement:

*each function of an angle is equal to the corresponding co-function of its complement.*

As an illustration of this we may refer to Art. 35, from which it will be seen that

$$\sin 60^\circ = \cos 30^\circ = \frac{\sqrt{3}}{2};$$

$$\sin 30^\circ = \cos 60^\circ = \frac{1}{2};$$

$$\tan 60^\circ = \cot 30^\circ = \sqrt{3}.$$

*Example 1.* Find a value of  $A$  when  $\cos 2A = \sin 3A$ .

Since  $\cos 2A = \sin (90^\circ - 2A)$ ,

the equation becomes  $\sin (90^\circ - 2A) = \sin 3A$ ;

$$\therefore 90^\circ - 2A = 3A;$$

whence

$$A = 18^\circ.$$

Thus *one* value of  $A$  which satisfies the equation is  $A = 18^\circ$ . In a later chapter we shall be able to solve the equation more completely, and shew that there are other values of  $A$  which satisfy it.

*Example 2.* Prove that  $\sec A \sec (90^\circ - A) = \tan A + \tan (90^\circ - A)$ .

Here it will be found easier to begin with the expression on the right side of the identity.

The second side  $= \tan A + \cot A$

$$= \frac{\sin A}{\cos A} + \frac{\cos A}{\sin A} = \frac{\sin^2 A + \cos^2 A}{\cos A \sin A}$$

$$= \frac{1}{\cos A \sin A}$$

$$= \sec A \operatorname{cosec} A = \sec A \sec (90^\circ - A).$$

### EXAMPLES. IV. b.

Find the complements of the following angles:

- |                     |                      |                        |
|---------------------|----------------------|------------------------|
| 1. $67^\circ 30'$ . | 2. $25^\circ 30''$ . | 3. $10^\circ 1' 3''$ . |
| 4. $45^\circ - A$ . | 5. $45^\circ + B$ .  | 6. $30^\circ - B$ .    |

7. In a triangle  $C$  is  $50^\circ$  and  $A$  is the complement of  $10^\circ$ ; find  $B$ .

8. In a triangle  $A$  is the complement of  $40^\circ$ ; and  $B$  is the complement of  $20^\circ$ ; find  $C$ .

Find a value of  $A$  in each of the following equations :

- |                          |  |
|--------------------------|--|
| 9. $\sin A = \cos 4A$ .  | 10. $\cos 3A = \sin 7A$ .                |
| 11. $\tan A = \cot 3A$ . | 12. $\cot A = \tan A$ .                  |
| 13. $\cot A = \tan 2A$ . | 14. $\sec 5A = \operatorname{cosec} A$ . |

Prove the following identities :

15.  $\sin(90^\circ - A) \cot(90^\circ - A) = \sin A$ .
16.  $\sin A \tan(90^\circ - A) \sec(90^\circ - A) = \cot A$ .
17.  $\cos A \tan A \tan(90^\circ - A) \operatorname{cosec}(90^\circ - A) = 1$ .
18.  $\sin A \cos(90^\circ - A) + \cos A \sin(90^\circ - A) = 1$ .
19.  $\cos(90^\circ - A) \operatorname{cosec}(90^\circ - A) = \tan A$ .
20.  $\operatorname{cosec}^2(90^\circ - A) = 1 + \sin^2 A \operatorname{cosec}^2(90^\circ - A)$ .
21.  $\sin A \cot A \cot(90^\circ - A) \sec(90^\circ - A) = 1$ .
22.  $\sec(90^\circ - A) - \cot A \cos(90^\circ - A) \tan(90^\circ - A) = \sin A$ .
23.  $\tan^2 A \sec^2(90^\circ - A) - \sin^2 A \operatorname{cosec}^2(90^\circ - A) = 1$ .
24.  $\tan(90^\circ - A) + \cot(90^\circ - A) = \operatorname{cosec} A \operatorname{cosec}(90^\circ - A)$ .
25.  $\frac{\sin(90^\circ - A)}{\sec(90^\circ - A)} \cdot \frac{\tan(90^\circ - A)}{\cos A} = \cos A$ .
26.  $\frac{\operatorname{cosec}^2 A \tan^2 A}{\cot(90^\circ - A)} \cdot \frac{\cot A}{\sec^2 A} = \sec^2(90^\circ - A) - 1$ .
27.  $\frac{\cot(90^\circ - A)}{\operatorname{cosec}^2 A} \cdot \frac{\sec A \cot^3 A}{\sin^2(90^\circ - A)} = \sqrt{\tan^2 A + 1}$ .
28.  $\frac{\cos^2(90^\circ - A)}{\operatorname{vers} A} = 1 + \sin(90^\circ - A)$ .
29.  $\frac{\cot^2 A \sin^2(90^\circ - A)}{\cot A + \cos A} = \tan(90^\circ - A) - \cos A$ .
30. If  $x \sin(90^\circ - A) \cot(90^\circ - A) = \cos(90^\circ - A)$ , find  $x$ .
31. Find the value of  $x$  which will satisfy  
 $\sec A \operatorname{cosec}(90^\circ - A) - x \cot(90^\circ - A) = 1$ .

### Easy Trigonometrical Equations.

40. As a further exercise in using the formulæ of Art. 29 and the numerical values of the functions of  $45^\circ$ ,  $60^\circ$ ,  $30^\circ$ , we shall now give some examples in trigonometrical equations.

*Example 1.* Solve  $4 \cos A = 3 \sec A$ .

By expressing the secant in terms of the cosine, we have

$$4 \cos A = \frac{3}{\cos A},$$

$$4 \cos^2 A = 3,$$

$$\cos A = \pm \frac{\sqrt{3}}{2}.$$

$$\therefore \cos A = \frac{\sqrt{3}}{2} \dots\dots\dots(1),$$

or  $\cos A = -\frac{\sqrt{3}}{2} \dots\dots\dots(2).$

Since  $\cos 30^\circ = \frac{\sqrt{3}}{2}$ , we see from (1) that  $A = 30^\circ$ .

The student will be able to understand the meaning of the negative result in (2) after he has read Chap. VIII.

*Example 2.* Solve  $3 \sec^2 \theta = 8 \tan \theta - 2$ .

Since  $\sec^2 \theta = 1 + \tan^2 \theta$ ,

we have  $3(1 + \tan^2 \theta) = 8 \tan \theta - 2$ ,

or  $3 \tan^2 \theta - 8 \tan \theta + 5 = 0$ .

This is a quadratic equation in which  $\tan \theta$  is the unknown quantity, and it may be solved by any of the rules for solving quadratic equations.

Thus  $(\tan \theta - 1)(3 \tan \theta - 5) = 0$ ,

therefore *either*  $\tan \theta - 1 = 0 \dots\dots\dots(1),$

or  $3 \tan \theta - 5 = 0 \dots\dots\dots(2).$

From (1),  $\tan \theta = 1$ , so that  $\theta = 45^\circ$ .

From (2),  $\tan \theta = \frac{5}{3}$ , a result which we cannot interpret at present.

41. When an equation involves more than two functions, it will usually be best to express each function in terms of the sine and cosine.



*Example.* Solve  $3 \tan \theta + \cot \theta = 5 \operatorname{cosec} \theta$ .

We have 
$$\frac{3 \sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = \frac{5}{\sin \theta},$$

$$3 \sin^2 \theta + \cos^2 \theta = 5 \cos \theta,$$

$$3(1 - \cos^2 \theta) + \cos^2 \theta = 5 \cos \theta,$$

$$2 \cos^2 \theta + 5 \cos \theta - 3 = 0,$$

$$(2 \cos \theta - 1)(\cos \theta + 3) = 0;$$

therefore *either*  $2 \cos \theta - 1 = 0 \dots\dots\dots(1),$

*or*  $\cos \theta + 3 = 0 \dots\dots\dots(2).$

From (1),  $\cos \theta = \frac{1}{2}$ , so that  $\theta = 60^\circ$ .

From (2),  $\cos \theta = -3$ , a result which must be rejected as *impossible*, because the numerical value of the cosine of an angle can never be greater than unity. [Art. 16.]

### EXAMPLES. IV. c.

Find a solution of each of the following equations:

1.  $2 \sin \theta = \operatorname{cosec} \theta.$
2.  $\tan \theta = 3 \cot \theta.$
3.  $\sec \theta = 4 \cos \theta.$
4.  $\sec \theta - \operatorname{cosec} \theta = 0.$
5.  $4 \sin \theta = 3 \operatorname{cosec} \theta.$
6.  $\operatorname{cosec}^2 \theta = 4.$
7.  $\sqrt{2} \cos \theta = \cot \theta.$
8.  $\tan \theta = 2 \sin \theta.$
9.  $\sec^2 \theta = 2 \tan^2 \theta.$
10.  $\operatorname{cosec}^2 \theta = 4 \cot^2 \theta.$
11.  $\sec^2 \theta = 3 \tan^2 \theta - 1.$
12.  $\sec^2 \theta + \tan^2 \theta = 7.$
13.  $\cot^2 \theta + \operatorname{cosec}^2 \theta = 3.$
14.  $2(\cos^2 \theta - \sin^2 \theta) = 1.$
15.  $2 \cos^2 \theta + 4 \sin^2 \theta = 3.$
16.  $6 \cos^2 \theta = 1 + \cos \theta.$
17.  $4 \sin \theta = 12 \sin^2 \theta - 1.$
18.  $2 \sin^2 \theta = 3 \cos \theta.$
19.  $\tan \theta = 4 - 3 \cot \theta.$
20.  $\cos^2 \theta - \sin^2 \theta = 2 - 5 \cos \theta.$
21.  $\cot \theta + \tan \theta = 2 \sec \theta.$
22.  $4 \operatorname{cosec} \theta + 2 \sin \theta = 9.$
23.  $\tan \theta - \cot \theta = \operatorname{cosec} \theta.$
24.  $2 \cos \theta + 2\sqrt{2} = 3 \sec \theta.$
25.  $2 \sin \theta \tan \theta + 1 = \tan \theta + 2 \sin \theta.$
26.  $6 \tan \theta - 5\sqrt{3} \sec \theta + 12 \cot \theta = 0.$
27. If  $\tan \theta + 3 \cot \theta = 4$ , prove that  $\tan \theta = 1$  or  $3$ .
28. Find  $\cot \theta$  from the equation  

$$\operatorname{cosec}^2 \theta + \cot^2 \theta = 3 \cot \theta.$$

### MISCELLANEOUS EXAMPLES. A.

1. Express as the decimal of a right angle

$$(1) 25^{\circ} 37' 6.4''; \quad (2) 63^{\circ} 21' 36''.$$

2. Shew that

$$\sin A \cos A \tan A + \cos A \sin A \cot A = 1.$$

3. A ladder 29 ft. long just reaches a window at a height of 21 ft. from the ground : find the cosine and cosecant of the angle made by the ladder with the ground.

4. If  $\operatorname{cosec} A = \frac{17}{15}$ , find  $\tan A$  and  $\sec A$ .

5. Shew that  $\operatorname{cosec}^2 A - \cot A \cos A \operatorname{cosec} A - 1 = 0$ .
- 

6. Reduce to sexagesimal measure

$$(1) 17^{\circ} 18' 75''; \quad (2) .0003 \text{ of a right angle.}$$

7.  $ABC$  is a triangle in which  $B$  is a right angle ; if  $c=9$ ,  $a=40$ , find  $b$ ,  $\cot A$ ,  $\sec A$ ,  $\sec C$ .

8. Which of the following statements are possible and which impossible ?

$$(1) 4 \sin \theta = 1; \quad (2) 2 \sec \theta = 1; \quad (3) 7 \tan \theta = 40.$$

9. Prove that  $\cos \theta \operatorname{vers} \theta (\sec \theta + 1) = \sin^2 \theta$ .

10. Express  $\sec a$  and  $\operatorname{cosec} a$  in terms of  $\cot a$ .
- 

11. Find the numerical value of

$$3 \tan^2 30^{\circ} + \frac{1}{4} \sec 60^{\circ} + 5 \cot^2 45^{\circ} - \frac{2}{3} \sin^2 60^{\circ}.$$

12. If  $\tan a = \frac{m}{n}$ , find  $\sin a$  and  $\sec a$ .

13. If  $m$  sexagesimal minutes are equivalent to  $n$  centesimal minutes, prove that  $m = .54n$ .

14. If  $\sin A = \frac{4}{5}$ , prove that  $\tan A + \sec A = 3$ , when  $A$  is an acute angle.

15. Shew that

$$\cot(90^\circ - A) \cot A \cos(90^\circ - A) \tan(90^\circ - A) = \cos A.$$


---

16.  $PQR$  is a triangle in which  $P$  is a right angle; if  $PQ = 21$ ,  $PR = 20$ , find  $\tan Q$  and  $\operatorname{cosec} Q$ .

17. Shew that  $(\tan a - \cot a) \sin a \cos a = 1 - 2 \cos^2 a$ .

18. Find a value of  $\theta$  which satisfies the equation

$$\sec 6\theta = \operatorname{cosec} 3\theta.$$

19. Prove that

$$\tan^2 60^\circ - 2 \tan^2 45^\circ = \cot^2 30^\circ - 2 \sin^2 30^\circ - \frac{3}{4} \operatorname{cosec}^2 45^\circ.$$

20. Solve the equations :

$$(1) \ 3 \sin \theta = 2 \cos^2 \theta; \quad (2) \ 5 \cot \theta - \operatorname{cosec}^2 \theta = 3.$$


---

21. Prove that  $1 + 2 \sec^2 A \tan^2 A - \sec^4 A - \tan^4 A = 0$ .

22. In the equation

$$8 \sin^2 \theta - 10 \sin \theta + 3 = 0,$$

shew that one value of  $\theta$  is impossible, and find the other value.

23. In a triangle  $ABC$  right-angled at  $C$ , prove that

$$\tan A + \tan B = \frac{c^2}{ab}.$$


24. If  $\cot A = c$ , shew that  $c + c^{-1} = \sec A \operatorname{cosec} A$ .

25. Shew that  $\frac{\sqrt{2 \operatorname{vers} A - \operatorname{vers}^2 A}}{1 - \operatorname{vers} A} = \tan A$ .

## CHAPTER V.

### SOLUTION OF RIGHT-ANGLED TRIANGLES.

42. EVERY triangle has six *parts*, namely, three sides and three angles. In Trigonometry it is usual to denote the three angles by the capital letters  $A, B, C$ , and the lengths of the sides respectively opposite to these angles by the letters  $a, b, c$ . It must be understood that  $a, b, c$  are *numerical quantities* expressing the number of units of length contained in the three sides.



43. We know from Geometry that it is always possible to construct a triangle when any three parts are given, provided that one at least of the parts is a side. Similarly, if the values of suitable parts of a triangle be given, we can by Trigonometry find the remaining parts. The process by which this is effected is called the **Solution of the triangle**.

The *general* solution of triangles will be discussed at a later stage; in this chapter we shall confine our attention to right-angled triangles.

44. From Euc. I. 47, we know that when a triangle is right-angled, if any two sides are given the third can be found. Thus in the figure of the next article, where  $ABC$  is a triangle right-angled at  $A$ , we have  $a^2 = b^2 + c^2$ ; whence if any two of the three quantities  $a, b, c$  are given, the third may be determined.


Again, the two acute angles are complementary, so that if one is given the other is also known.

Hence in the solution of right-angled triangles there are really only two cases to be considered :

- I. when *any two sides* are given ;
- II. when *one side* and *one acute angle* are given.

45. CASE I. *To solve a right-angled triangle when two sides are given.*

Let  $ABC$  be a right-angled triangle, of which  $A$  is the right angle, and suppose that any two sides are given;



then the third side may be found from the equation

$$a^2 = b^2 + c^2.$$

Also

$$\cos C = \frac{b}{a}, \text{ and } B = 90^\circ - C;$$

whence  $C$  and  $B$  may be obtained.


*Example.* Given  $B = 90^\circ$ ,  $a = 20$ ,  $b = 40$ , solve the triangle.

$$\begin{aligned} \text{Here } c^2 &= b^2 - a^2 \\ &= 1600 - 400 = 1200; \\ \therefore c &= 20\sqrt{3}. \end{aligned}$$

$$\text{Also } \sin A = \frac{a}{b} = \frac{20}{40} = \frac{1}{2};$$

$$\therefore A = 30^\circ.$$

$$\text{And } C = 90^\circ - A = 90^\circ - 30^\circ = 60^\circ.$$



The solution of a trigonometrical problem may often be obtained in more than one way. In the present case the triangle can be solved without making use of Euc. I. 47.

Another solution may be given as follows:

$$\cos C = \frac{a}{b} = \frac{20}{40} = \frac{1}{2};$$

$$\therefore C = 60^\circ.$$

$$\text{And } A = 90^\circ - C = 90^\circ - 60^\circ = 30^\circ.$$

$$\text{Also } \frac{c}{40} = \cos A = \cos 30^\circ = \frac{\sqrt{3}}{2};$$

$$\therefore c = 20\sqrt{3}.$$

46. CASE II. *To solve a right-angled triangle when one side and one acute angle are given.*

Let  $ABC$  be a right-angled triangle of which  $A$  is the right angle, and suppose one side  $b$  and one acute angle  $C$  are given; then

$$B = 90^\circ - C, \quad \frac{a}{b} = \sec C, \quad \frac{c}{b} = \tan C;$$

whence  $B, a, c$  may be determined.

*Example 1.* Given  $B = 90^\circ, A = 30^\circ, c = 5$ , solve the triangle.



We have  $C = 90^\circ - A = 90^\circ - 30^\circ = 60^\circ$ .

Also 
$$\frac{a}{5} = \tan 30^\circ;$$

$$\begin{aligned} \therefore a &= 5 \tan 30^\circ = \frac{5}{\sqrt{3}} \\ &= \frac{5}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{5\sqrt{3}}{3}. \end{aligned}$$

Again, 
$$\frac{b}{5} = \sec 30^\circ;$$

$$\therefore b = 5 \sec 30^\circ = 5 \times \frac{2}{\sqrt{3}} = \frac{10}{\sqrt{3}} = \frac{10\sqrt{3}}{3}.$$



NOTE. The student should observe that in each case we write down a ratio which connects *the side we are finding with that whose value is given*, and a knowledge of the ratios of the given angle enables us to complete the solution.

*Example 2.* If  $C = 90^\circ, B = 25^\circ 43'$ , and  $c = 100$ , solve the triangle, having given  $\tan 25^\circ 43' = .482$  and  $\cos 25^\circ 43' = .901$ .

Here  $A = 90^\circ - B$   
 $= 90^\circ - 25^\circ 43' = 64^\circ 17'.$


Now 
$$\frac{a}{c} = \cos B;$$

that is, 
$$\frac{a}{100} = \cos 25^\circ 43';$$

$$\begin{aligned} \therefore a &= 100 \cos 25^\circ 43' \\ &= 100 \times .901 = 90.1. \end{aligned}$$

Also 
$$\frac{b}{a} = \tan B, \text{ or } b = a \tan B;$$

$$\therefore b = 90.1 \times \tan 25^\circ 43' = 90.1 \times .482 = 43.4282.$$





## EXAMPLES. V. a.

Solve the triangles in which the following parts are given :

1.  $A=90^\circ$ ,  $a=4$ ,  $b=2\sqrt{3}$ .
2.  $c=6$ ,  $b=12$ ,  $B=90^\circ$ .
3.  $C=90^\circ$ ,  $b=12$ ,  $a=4\sqrt{3}$ .
4.  $a=60$ ,  $b=30$ ,  $A=90^\circ$ .
5.  $a=20$ ,  $c=20$ ,  $B=90^\circ$ .
6.  $a=5\sqrt{3}$ ,  $b=15$ ,  $C=90^\circ$ .
7.  $b=c=2$ ,  $A=90^\circ$ .
8.  $2c=b=6\sqrt{3}$ ,  $B=90^\circ$ .
9.  $C=90^\circ$ ,  $a=9\sqrt{3}$ ,  $A=30^\circ$ .
10.  $A=90^\circ$ ,  $B=30^\circ$ ,  $a=4$ .
11.  $A=60^\circ$ ,  $c=8$ ,  $C=90^\circ$ .
12.  $A=60^\circ$ ,  $C=30^\circ$ ,  $b=6$ .
13.  $B=90^\circ$ ,  $C=60^\circ$ ,  $b=100$ .
14.  $A=30^\circ$ ,  $B=60^\circ$ ,  $b=20\sqrt{3}$ .
15.  $B=C=45^\circ$ ,  $c=4$ .
16.  $2B=C=60^\circ$ ,  $a=8$ .
17. If  $C=90^\circ$ ,  $\cot A=.07$ ,  $b=49$ , find  $a$ .
18. If  $C=90^\circ$ ,  $A=38^\circ 19'$ ,  $c=50$ , find  $a$ ;  
given  $\sin 38^\circ 19'=.62$ .
19. If  $a=100$ ,  $B=90^\circ$ ,  $C=40^\circ 51'$ , find  $c$ ;  
given  $\tan 40^\circ 51'=.8647$ .
20. If  $b=20$ ,  $A=90^\circ$ ,  $C=78^\circ 12'$ , find  $a$ ;  
given  $\sec 78^\circ 12'=4.89$ .
21. If  $B=90^\circ$ ,  $A=36^\circ$ ,  $c=100$ , solve the triangle;  
given  $\tan 36^\circ=.73$ ,  $\sec 36^\circ=1.24$ .
22. If  $A=90^\circ$ ,  $c=37$ ,  $a=100$ , solve the triangle;  
given  $\sin 21^\circ 43'=.37$ ,  $\cos 21^\circ 43'=.93$ .
23. If  $A=90^\circ$ ,  $B=39^\circ 24'$ ,  $b=25$ , solve the triangle;  
given  $\cot 39^\circ 24'=1.2174$ ,  $\operatorname{cosec} 39^\circ 24'=1.5755$ .
24. If  $C=90^\circ$ ,  $a=225$ ,  $b=272$ , solve the triangle;  
given  $\tan 50^\circ 24'=1.209$ .

47. It will be found that all the varieties of the solution of right-angled triangles which can arise are either included in the two cases of Arts. 45 and 46, or in some modification of them. Sometimes the solution of a problem may be obtained by solving *two* right-angled triangles. The two examples we give as illustrations will in various forms be frequently met with in subsequent chapters.



*Example 1.* In the triangle  $ABC$ , the angles  $A$  and  $B$  are equal to  $30^\circ$  and  $135^\circ$  respectively, and the side  $AB$  is 100 feet; find the length of the perpendicular from  $C$  upon  $AB$  produced.


Draw  $CD$  perpendicular to  $AB$  produced, and let  $CD = x$ .

Then  $\angle CBD = 180^\circ - 135^\circ = 45^\circ$ ;

$$\therefore BD = CD = x.$$

Now in the right-angled triangle  $ADC$ ,

$$\frac{CD}{AD} = \tan DAC = \tan 30^\circ;$$



that is,

$$\frac{x}{x+100} = \frac{1}{\sqrt{3}};$$

$$\therefore x\sqrt{3} = x+100;$$

$$x(\sqrt{3}-1) = 100,$$

$$x = \frac{100}{\sqrt{3}-1} = \frac{100(\sqrt{3}+1)}{3-1};$$

$$\therefore x = 50(\sqrt{3}+1).$$

Thus the distance required is  $50(\sqrt{3}+1)$  feet.

*Example 2.* In the triangle  $ABC$ ,  $AD$  is drawn perpendicular to  $BC$ ; solve the triangle, having given

$$AD = 5, \angle ABD = 60^\circ, \angle ACD = 45^\circ.$$

In the right-angled triangle  $ABD$ ,

$$\frac{AB}{AD} = \operatorname{cosec} ABD;$$


$$\therefore AB = AD \operatorname{cosec} ABD = 5 \operatorname{cosec} 60^\circ$$

$$= 5 \times \frac{2}{\sqrt{3}} = \frac{10}{\sqrt{3}} = \frac{10\sqrt{3}}{3}.$$

$$\text{Also } \frac{BD}{AD} = \cot ABD;$$

$$\therefore BD = AD \cot ABD$$

$$= 5 \cot 60^\circ = \frac{5}{\sqrt{3}} = \frac{5\sqrt{3}}{3}.$$



In the right-angled triangle  $ADC$ ,

$$\angle DAC = 45^\circ = \angle DCA;$$

$$\therefore DC = DA = 5.$$

Thus  $BC = CD + BD = 5 + \frac{5\sqrt{3}}{3} = \frac{15 + 5\sqrt{3}}{3}.$

And  $\frac{AC}{AD} = \operatorname{cosec} ACD;$

$$\therefore AC = AD \operatorname{cosec} ACD = 5 \operatorname{cosec} 45^\circ = 5\sqrt{2}.$$

Finally,  $\angle BAC = 180^\circ - 60^\circ - 45^\circ = 75^\circ.$

Thus  $a = \frac{15 + 5\sqrt{3}}{3}, \quad b = 5\sqrt{2}, \quad c = \frac{10\sqrt{3}}{3}, \quad A = 75^\circ.$

### EXAMPLES. V. b.

1.  $ABC$  is a triangle, and  $BD$  is perpendicular to  $AC$  produced: find  $BD$ , given

$$A = 30^\circ, \quad C = 120^\circ, \quad AC = 20.$$

2. If  $BD$  is perpendicular to the base  $AC$  of a triangle  $ABC$ , find  $a$  and  $c$ , given

$$A = 30^\circ, \quad C = 45^\circ, \quad BD = 10.$$

3. In the triangle  $ABC$ ,  $AD$  is drawn perpendicular to  $BC$  making  $BD$  equal to 15 ft.: find the lengths of  $AB$ ,  $AC$ , and  $AD$ , given that  $B$  and  $C$  are equal to  $30^\circ$  and  $60^\circ$  respectively.

4. In a right-angled triangle  $PQR$ , find the segments of the hypotenuse  $PR$  made by the perpendicular from  $Q$ ; given

$$QR = 8, \quad \angle QRP = 60^\circ, \quad \angle QPR = 30^\circ.$$

5. If  $PQ$  is drawn perpendicular to the straight line  $QRS$ , find  $RS$ , given

$$PQ = 36, \quad \angle RPQ = 30^\circ, \quad \angle SPQ = 60^\circ.$$

6. If  $PQ$  is drawn perpendicular to the straight line  $QRS$ , find  $RS$ , given

$$PQ = 20, \quad \angle PRS = 135^\circ, \quad \angle PSR = 30^\circ.$$

7. In the triangle  $ABC$ , the angles  $B$  and  $C$  are equal to  $45^\circ$  and  $120^\circ$  respectively; if  $a = 40$  find the length of the perpendicular from  $A$  on  $BC$  produced.

8. If  $CD$  is drawn perpendicular to the straight line  $DBA$ , find  $DC$  and  $BD$ , given

$$AB = 59, \quad \angle CBD = 45^\circ, \quad \angle CAB = 32^\circ 50', \quad \cot 32^\circ 50' = 1.59.$$


## CHAPTER VI.

### EASY PROBLEMS.

48. THE principles explained in the previous chapters may now be applied to the solution of problems in heights and distances. It will be assumed that by the use of suitable instruments the necessary lines and angles can be measured with sufficient accuracy for the purposes required.

After the practice afforded by the examples in the last chapter, the student should be able to write down at once any side of a right-angled triangle in terms of another through the medium of the functions of either acute angle. In the present and subsequent chapters it is of great importance to acquire readiness in this respect.


For instance, from the adjoining figure, we have



$$\begin{aligned} a &= c \sin A, & a &= c \cos B, & a &= b \cot B, \\ a &= b \tan A, & c &= a \sec B, & b &= a \tan B. \end{aligned}$$

These relations are not to be committed to memory but in each case should be read off from the figure. There are several other similar relations connecting the parts of the above triangle, and the student should practise himself in obtaining them quickly.

*Example.*  $Q, R, T$  are three points in a straight line, and  $TP$  is drawn perpendicular to  $QT$ . If  $PT=a$ ,  $\angle PQT=\beta$ ,  $\angle PRT=2\beta$ , express the lengths of all the lines of the figure in terms of  $a$  and  $\beta$ .



By Euc. I. 32,

$$\begin{aligned}\angle QPR &= \angle PRT - \angle PQR; \\ \therefore \angle QPR &= 2\beta - \beta = \beta = \angle PQR; \\ \therefore QR &= PR.\end{aligned}$$

In the right-angled triangle  $PRT$ ,

$$\begin{aligned}PR &= a \operatorname{cosec} 2\beta; \\ \therefore QR &= a \operatorname{cosec} 2\beta.\end{aligned}$$



Also

$$TR = a \cot 2\beta.$$

Lastly, in the right-angled triangle  $PQT$ ,

$$\begin{aligned}QT &= a \cot \beta, \\ PQ &= a \operatorname{cosec} \beta.\end{aligned}$$

**49. Angles of elevation and depression.** Let  $OP$  be a horizontal line in the same vertical plane as an object  $Q$ , and let  $OQ$  be joined.



In Fig. 1, where the object  $Q$  is *above* the horizontal line  $OP$ , the angle  $POQ$  is called the **angle of elevation** of the object  $Q$  as seen from the point  $O$ .


In Fig. 2, where the object  $Q$  is *below* the horizontal line  $OP$ , the angle  $POQ$  is called the **angle of depression** of the object  $Q$  as seen from the point  $O$ .

**Example I.** A flagstaff stands on a horizontal plane, and from a point on the ground at a distance of 30 ft. its angle of elevation is  $60^\circ$ : find its height.

Let  $AB$  be the flagstaff,  $C$  the point of observation; then

$$\begin{aligned} AB &= BC \tan 60^\circ = 30 \sqrt{3} \\ &= 30 \times 1.732 = 51.96. \end{aligned}$$

Thus the height is 51.96 ft.



### EXAMPLES. VI. a.

[The results should be expressed in a form free from surds by using the approximations  $\sqrt{2}=1.414$ ,  $\sqrt{3}=1.732$ .]

1. The angle of elevation of the top of a chimney at a distance of 300 feet is  $30^\circ$ : find its height.

2. From a ship's masthead 160 feet high the angle of depression of a boat is observed to be  $30^\circ$ : find its distance from the ship.

3. Find the angle of elevation of the sun when the shadow of a pole 6 feet high is  $2\sqrt{3}$  feet long.

4. At a distance 86.6 feet from the foot of a tower the angle of elevation of the top is  $30^\circ$ . Find the height of the tower and the observer's distance from the top.

5. A ladder 45 feet long just reaches the top of a wall. If the ladder makes an angle of  $60^\circ$  with the wall, find the height of the wall, and the distance of the foot of the ladder from the wall.

6. Two masts are 60 feet and 40 feet high, and the line joining their tops makes an angle of  $33^\circ 41'$  with the horizon: find their distance apart, given  $\cot 33^\circ 41' = 1.5$ .

7. Find the distance of the observer from the top of a cliff which is 132 yards high, given that the angle of elevation is  $41^\circ 18'$ , and that  $\sin 41^\circ 18' = .66$ .

8. One chimney is 30 yards higher than another. A person standing at a distance of 100 yards from the lower observes their tops to be in a line inclined at an angle of  $27^\circ 2'$  to the horizon: find their heights, given  $\tan 27^\circ 2' = .51$ .

**Example II.** From the foot of a tower the angle of elevation of the top of a column is  $60^\circ$ , and from the top of the tower, which is 50 ft. high, the angle of elevation is  $30^\circ$ : find the height of the column.

Let  $AB$  denote the column and  $CD$  the tower; draw  $CE$  parallel to  $DB$ .

Let  $AB = x$ ;

then  $AE = AB - BE = x - 50$ .

Let  $DB = CE = y$ .

From the right-angled triangle  $ADB$ ,

$$y = x \cot 60^\circ = \frac{x}{\sqrt{3}}.$$

From the right-angled triangle  $ACE$ ,

$$y = (x - 50) \cot 30^\circ = \sqrt{3} (x - 50).$$

$$\therefore \frac{x}{\sqrt{3}} = \sqrt{3} (x - 50),$$

$$x = 3 (x - 50);$$

$$x = 75.$$

whence

Thus the column is 75 ft. high.


9. The angle of elevation of the top of a tower is  $30^\circ$ ; on walking 100 yards nearer the elevation is found to be  $60^\circ$ : find the height of the tower.

10. A flagstaff stands upon the top of a building; at a distance of 40 feet the angles of elevation of the tops of the flagstaff and building are  $60^\circ$  and  $30^\circ$ : find the length of the flagstaff.

11. The angles of elevation of a spire at two places due east of it and 200 feet apart are  $45^\circ$  and  $30^\circ$ : find the height of the spire.

12. From the foot of a post the elevation of the top of a steeple is  $45^\circ$ , and from the top of the post, which is 30 feet high, the elevation is  $30^\circ$ : find the height and distance of the steeple.


13. The height of a hill is 3300 feet above the level of a horizontal plane. From a point  $A$  on this plane the angular elevation of the top of the hill is  $60^\circ$ . A balloon rises from  $A$  and ascends vertically upwards at a uniform rate; after 5 minutes the angular elevation of the top of the hill to an observer in the balloon is  $30^\circ$ : find the rate of the balloon's ascent in miles per hour.





**Example III.** From the top of a cliff 150 ft. high the angles of depression of two boats which are due South of the observer are  $15^\circ$  and  $75^\circ$ : find their distance apart, having given

$$\cot 15^\circ = 2 + \sqrt{3} \text{ and } \cot 75^\circ = 2 - \sqrt{3}.$$



Let  $OA$  represent the cliff,  $B$  and  $C$  the boats. Let  $OP$  be a horizontal line through  $O$ ; then

$$\angle POC = 15^\circ \text{ and } \angle POB = 75^\circ;$$

$$\therefore \angle OCA = 15^\circ \text{ and } \angle OBA = 75^\circ.$$

Let  $CB = x$ ,  $AB = y$ ; then  $CA = x + y$ .

From the right-angled triangle  $OBA$ ,

$$y = 150 \cot 75^\circ = 150 (2 - \sqrt{3}) = 300 - 150\sqrt{3}.$$

From the right-angled triangle  $OCA$ ,

$$x + y = 150 \cot 15^\circ = 150 (2 + \sqrt{3}) = 300 + 150\sqrt{3}.$$

By subtraction,  $x = 300\sqrt{3} = 519.6$ .

Thus the distance between the boats is 519.6 ft.

14. From the top of a monument 100 feet high, the angles of depression of two objects on the ground due west of the monument are  $45^\circ$  and  $30^\circ$ : find the distance between them.

15. The angles of depression of the top and foot of a tower seen from a monument 96 feet high are  $30^\circ$  and  $60^\circ$ : find the height of the tower.


16. From the top of a cliff 150 feet high the angles of depression of two boats at sea, each due north of the observer, are  $30^\circ$  and  $15^\circ$ : how far are the boats apart?

17. From the top of a hill the angles of depression of two consecutive milestones on a level road running due south from the observer are  $45^\circ$  and  $22^\circ$  respectively. If  $\cot 22^\circ = 2.475$  find the height of the hill in yards.

18. From the top of a lighthouse 80 yards above the horizon the angles of depression of two rocks due west of the observer are  $75^\circ$  and  $15^\circ$ : find their distance apart, given  $\cot 75^\circ = .268$  and  $\cot 15^\circ = 3.732$ .



50. Trigonometrical Problems sometimes require a knowledge of the **Points of the Mariner's Compass**, which we shall now explain.



In the above figure, it will be seen that 32 points are taken at equal distances on the circumference of a circle, so that the arc between any two consecutive points subtends at the centre of the circle an angle equal to  $3\frac{6}{2}^{\circ}$ , that is to  $11\frac{1}{4}^{\circ}$ .

The points North, South, East, West are called the **Cardinal Points**, and with reference to them the other *points* receive their names. The student will have no difficulty in learning these if he will carefully notice the arrangement in any one of the principal quadrants.

51. Sometimes a slightly different notation is used; thus N.  $11\frac{1}{4}^{\circ}$  E. means a direction  $11\frac{1}{4}^{\circ}$  east of north, and is therefore the same as N. by E. Again S.W. by S. is 3 *points* from south and may be expressed by S.  $33\frac{3}{4}^{\circ}$  W., or since it is 5 *points* from west it can also be expressed by W.  $56\frac{1}{4}^{\circ}$  S. In each of these cases it will be seen that the angular measurement is made from the direction which is first mentioned.

52. The angle between the directions of any two points is obtained by multiplying  $11\frac{1}{4}^\circ$  by the number of intervals between the points. Thus between S. by W. and W.S.W. there are 5 intervals and the angle is  $56\frac{1}{4}^\circ$ ; between N.E. by E. and S.E. there are 7 intervals and the angle is  $78\frac{3}{4}^\circ$ .

53. If  $B$  lies in a certain direction with respect to  $A$ , it is said to *bear* in that direction from  $A$ ; thus Birmingham *bears* N.W. of London, and from Birmingham the *bearing* of London is S.E.

*Example 1.* From a lighthouse  $L$  two ships  $A$  and  $B$  are observed in directions S.W. and  $15^\circ$  East of South respectively. At the same time  $B$  is observed from  $A$  in a S.E. direction. If  $LA$  is 4 miles find the distance between the ships.

Draw  $LS'$  due South; then from the bearings of the two ships,

$$\angle ALS' = 45^\circ, \quad \angle BLS' = 15^\circ,$$

so that

$$\angle ALB = 60^\circ.$$

Through  $A$  draw a line  $NS$  pointing North and South; then

$$\angle NAL = \angle ALS' = 45^\circ,$$

and  $\angle BAS = 45^\circ$ , since  $B$  bears S.E. from  $A$ ;


$$\text{hence } \angle BAL = 180^\circ - 45^\circ - 45^\circ = 90^\circ.$$

In the right-angled triangle  $ABL$ ,

$$AB = AL \tan \angle ALB = 4 \tan 60^\circ$$

$$= 4\sqrt{3} = 6.928.$$

Thus the distance between the ships is 6.928 miles.



*Example 2.* At 9 A.M. a ship which is sailing in a direction E.  $40^\circ$  S. at the rate of 8 miles an hour observes a fort in a direction  $50^\circ$  North of East. At 11 A.M. the fort is observed to bear N.  $20^\circ$  W.: find the distance of the fort from the ship at each observation.


Let  $A$  and  $C$  be the first and second positions of the ship;  $B$  the fort.

Through  $A$  draw lines towards the cardinal points of the compass.

From the observations made

$$\angle EAC = 40^\circ, \quad \angle EAB = 50^\circ, \quad \text{so that } \angle BAC = 90^\circ.$$

Through  $C$  draw  $CN'$  towards the North; then  $\angle BCN' = 20^\circ$ , for the bearing of the fort from  $C$  is N.  $20^\circ$  W.



Also  $\angle ACN' = \angle CAS = 90^\circ - 40^\circ = 50^\circ$ ;  
 $\therefore \angle ACB = \angle ACN' - \angle BCN' = 50^\circ - 20^\circ = 30^\circ$ .

In the right-angled triangle  $ACB$ ,

$$AB = AC \tan ACB = 16 \tan 30^\circ = \frac{16}{\sqrt{3}} = \frac{16\sqrt{3}}{3} = 9.237 \text{ nearly};$$

$$\text{and } BC = AC \sec ACB = 16 \sec 30^\circ = 16 \times \frac{2}{\sqrt{3}} = \frac{32\sqrt{3}}{3} = 18.475 \text{ nearly.}$$

Thus the distances are 9.237 and 18.475 miles nearly.

### EXAMPLES. VI. b.

1. A person walking due E. observes two objects both in the N.E. direction. After walking 800 yards one of the objects is due N. of him, and the other lies N.W.: how far was he from the objects at first?

2. Sailing due E. I observe two ships lying at anchor due S.; after sailing 3 miles the ships bear  $60^\circ$  and  $30^\circ$  S. of W.; how far are they now distant from me?

3. Two vessels leave harbour at noon in directions W.  $28^\circ$  S. and E.  $62^\circ$  S. at the rates 10 and  $10\frac{1}{2}$  miles per hour respectively. Find their distance apart at 2 p.m.

4. A lighthouse facing N. sends out a fan-shaped beam extending from N.E. to N.W. A steamer sailing due W. first sees the light when 5 miles away from the lighthouse and continues to see it for  $30\sqrt{2}$  minutes. What is the speed of the steamer?

5. A ship sailing due S. observes two lighthouses in a line exactly W. After sailing 10 miles they are respectively N.W. and W.N.W.; find their distances from the position of the ship at the first observation.

6. Two vessels sail from port in directions N.  $35^\circ$  W. and S.  $55^\circ$  W. at the rates of 8 and  $8\sqrt{3}$  miles per hour respectively. Find their distance apart at the end of an hour, and the bearing of the second vessel as observed from the first.

7. A vessel sailing S.S.W. is observed at noon to be E.S.E. from a lighthouse 4 miles away. At 1 p.m. the vessel is due S. of the lighthouse: find the rate at which the vessel is sailing. Given  $\tan 67\frac{1}{2}^\circ = 2.414$ .

8.  $A, B, C$  are three places such that from  $A$  the bearing of  $C$  is N.  $10^\circ$  W., and the bearing of  $B$  is N.  $50^\circ$  E.; from  $B$  the bearing of  $C$  is N.  $40^\circ$  W. If the distance between  $B$  and  $C$  is 10 miles, find the distances of  $B$  and  $C$  from  $A$ .

9. A ship steaming due E. sights at noon a lighthouse bearing N.E., 15 miles distant; at 1.30 p.m. the lighthouse bears N.W. How many knots per day is the ship making? Given 60 knots = 69 miles.

10. At 10 o'clock forenoon a coaster is observed from a lighthouse to bear 9 miles away to N.E. and to be holding a south-easterly course; at 1 p.m. the bearing of the coaster is  $15^\circ$  S. of E. Find the rate of the coaster's sailing and its distance from the lighthouse at the time of the second observation.

11. The distance between two lighthouses,  $A$  and  $B$ , is 12 miles and the line joining them bears E.  $15^\circ$  N. At midnight a vessel which is sailing S.  $15^\circ$  E. at the rate of 10 miles per hour is N.E. of  $A$  and N.W. of  $B$ : find to the nearest minute when the vessel crosses the line joining the lighthouses.


12. From  $A$  to  $B$ , two stations of a railway, the line runs W.S.W. At  $A$  a person observes that two spires, whose distance apart is 1.5 miles, are in the same line which bears N.N.W. At  $B$  their bearings are N.  $7\frac{1}{2}^\circ$  E. and N.  $37\frac{1}{2}^\circ$  E. Find the rate of a train which runs from  $A$  to  $B$  in 2 minutes.

## CHAPTER VII.

### RADIAN OR CIRCULAR MEASURE.

54. WE shall now return to the system of measuring angles which was briefly referred to in Art. 6. In this system angles are not measured in terms of a submultiple of the right angle, as in the sexagesimal and centesimal methods, but a certain angle known as a *radian* is taken as the standard unit, in terms of which all other angles are measured.

55. DEFINITION. A **radian** is the angle subtended at the centre of any circle by an arc equal in length to the radius of the circle.




In the above figure,  $ABC$  is a circle, and  $O$  its centre. If on the circumference we measure an arc  $AB$  equal to the radius and join  $OA$ ,  $OB$ , the angle  $AOB$  is a radian.

56. In any system of measurement it is essential that the unit should be always the same. In order to shew that a radian, constructed according to the above definition, is of constant magnitude, we must first establish an important property of the circle.



57. *The circumferences of circles are to one another as their radii.*

Take *any* two circles whose radii are  $r_1$  and  $r_2$ , and in each circle let a regular polygon of  $n$  sides be described.



Let  $A_1B_1$  be a side of the first,  $A_2B_2$  a side of the second polygon, and let their lengths be denoted by  $a_1, a_2$ . Join their extremities to  $O_1$  and  $O_2$  the centres of the circles. We thus obtain two isosceles triangles whose vertical angles are equal, each being  $\frac{1}{n}$  of four right angles.

Hence the triangles are equiangular, and therefore we have by Euc. VI. 4,

$$\frac{A_1B_1}{O_1A_1} = \frac{A_2B_2}{O_2A_2};$$

that is,

$$\frac{a_1}{r_1} = \frac{a_2}{r_2};$$

$$\therefore \frac{na_1}{r_1} = \frac{na_2}{r_2};$$

that is,

$$\frac{p_1}{r_1} = \frac{p_2}{r_2},$$

where  $p_1$  and  $p_2$  are the perimeters of the polygons. This is true whatever be the number of sides in the polygons. By taking  $n$  sufficiently large we can make the perimeters of the two polygons differ from the circumferences of the corresponding circles by as small a quantity as we please; so that ultimately

$$\frac{c_1}{r_1} = \frac{c_2}{r_2},$$

where  $c_1$  and  $c_2$  are the circumferences of the two circles.

58. It thus appears that *the ratio of the circumference of a circle to its radius is the same whatever be the size of the circle; that is,*

*in all circles  $\frac{\text{circumference}}{\text{diameter}}$  is a constant quantity.*

This constant is incommensurable and is always denoted by the Greek letter  $\pi$ . Though its numerical value cannot be found exactly, it is shewn in a later part of the subject that it can be obtained to any degree of approximation. To ten decimal places its value is 3.1415926536. In many cases  $\pi = \frac{22}{7}$ , which is true to two decimal places, is a sufficiently close approximation; where greater accuracy is required the value 3.1416 may be used.

59. If  $c$  denote the circumference of the circle whose radius is  $r$ , we have

$$\frac{\text{circumference}}{\text{diameter}} = \pi;$$

$$\therefore \frac{c}{2r} = \pi,$$

or

$$c = 2\pi r.$$


60. *To prove that all radians are equal.*

Draw *any* circle; let  $O$  be its centre and  $OA$  a radius. Let the arc  $AB$  be measured equal in length to  $OA$ . Join  $OB$ ; then  $\angle AOB$  is a radian. Produce  $AO$  to meet the circumference in  $C$ . By Euc. VI. 33, angles at the centre of a circle are proportional to the arcs on which they stand; hence

$$\frac{\angle AOB}{\text{two right angles}} = \frac{\text{arc } AB}{\text{arc } ABC}$$

$$= \frac{\text{radius}}{\text{semi-circumference}} = \frac{r}{\pi r} = \frac{1}{\pi},$$

which is constant; that is, *a radian always bears the same ratio to two right angles, and therefore is a constant angle.*






61. Since a radian is constant it is taken as a standard unit, and the *number of radians* contained in any angle is spoken of as its **radian measure** or **circular measure**. [See Art. 71.] In this system, an angle is usually denoted by a *mere number*, the unit being implied. Thus when we speak of an angle 2·5, it is understood that its radian measure is 2·5, or, in other words, that the angle contains  $2\frac{1}{2}$  radians.

Where it is desirable to refer to the unit expressly, a radian may be denoted by the letter  $\rho$ , thus  $3\rho$  represents an angle which contains three radians.

62. *To find the radian measure of a right angle.*

Let  $\angle AOC$  be a right angle at the centre of a circle, and  $AOB$  a radian; then the radian measure of  $\angle AOC$

$$\begin{aligned} &= \frac{\angle AOC}{\angle AOB} = \frac{\text{arc } AC}{\text{arc } AB} \\ &= \frac{\frac{1}{4} (\text{circumference})}{\text{radius}} = \frac{\frac{1}{4} (2\pi r)}{r} \\ &= \frac{\pi}{2}; \end{aligned}$$



that is, a right angle contains  $\frac{\pi}{2}$  radians.

63. *To find the number of degrees in a radian.*

From the last article it follows that

$$\pi \text{ radians} = 2 \text{ right angles} = 180 \text{ degrees.}$$

$$\therefore \text{a radian} = \frac{180}{\pi} \text{ degrees.}$$

By division we find that  $\frac{1}{\pi} = \cdot 31831$  nearly;

hence approximately, a radian  $= 180 \times \cdot 31831 = 57\cdot 2958$  degrees.

64. The formula

$$\pi \text{ radians} = 180 \text{ degrees}$$

connecting the sexagesimal and radian measures of an angle, is a useful result which enables us to pass readily from one system to the other.

*Example.* Express  $75^\circ$  in radian measure, and  $\frac{\pi}{54}$  in sexagesimal measure.

(1) Since  $180 \text{ degrees} = \pi \text{ radians}$ ,

$$75 \text{ degrees} = \frac{75}{180} \pi \text{ radians} = \frac{5\pi}{12} \text{ radians.}$$

Thus the radian measure is  $\frac{5\pi}{12}$ .

(2) Since  $\pi \text{ radians} = 180 \text{ degrees}$ ,

$$\frac{\pi}{54} \text{ radians} = \frac{180}{54} \text{ degrees.}$$

Thus the angle  $= \frac{10}{3} \text{ degrees} = 3^\circ 20'$ .

65. It may be well to remind the student that the symbol  $\pi$  always denotes a *number*, viz.  $3.14159\dots$  When the symbol stands alone, without reference to any angle, there can be no ambiguity; but even when  $\pi$  is used to denote an angle, it must still be remembered that  $\pi$  is a *number*, namely, the number of radians in two right angles.

NOTE. It is not uncommon for beginners to make statements such as " $\pi = 180$ " or " $\frac{\pi}{2} = 90$ ." Without some modification this mode of expression is quite incorrect. It is true that  $\pi$  *radians* are equal to  $180$  *degrees*, but the statement ' $\pi = 180$ ' is no more correct than the statement " $20 = 1$ " to denote the equivalence of 20 shillings and 1 sovereign.

66. *If the number of degrees and radians in an angle be represented by  $D$  and  $\theta$  respectively, to prove that*

$$\frac{D}{180} = \frac{\theta}{\pi}.$$

In sexagesimal measure, the ratio of the given angle to two right angles is expressed by  $\frac{D}{180}$ .

In radian measure, the ratio of these same two angles is expressed by  $\frac{\theta}{\pi}$ .

$$\therefore \frac{D}{180} = \frac{\theta}{\pi}.$$

*Example 1.* What is the radian measure of  $45^\circ 13' 30''$ ?

If  $D$  be the number of degrees in the angle,  
we have  $D = 45.225$ .

$$\begin{array}{r} 60 \ ) \ 30 \\ 60 \ ) \ 18.5 \\ \quad .225 \end{array}$$

Let  $\theta$  be the number of radians in the given angle, then

$$\frac{\theta}{\pi} = \frac{45.225}{180} = \frac{1.005}{4};$$

$$\therefore \theta = \frac{\pi}{4} \times 1.005 = \frac{3.1416}{4} \times 1.005$$

$$= .7854 \times 1.005 = .789327.$$

Thus the radian measure is .789327.

*Example 2.* Express in sexagesimal measure the angle whose radian measure is 1.309.

Let  $D$  be the number of degrees; then

$$\frac{D}{180} = \frac{1.309}{\pi};$$

$$\therefore D = \frac{180 \times 1.309}{3.1416} = \frac{180 \times 1309 \times 10}{31416}$$

$$= \frac{180 \times 10}{24} = 75.$$

Thus the angle is  $75^\circ$ .

## EXAMPLES. VII. a.

[Unless otherwise stated  $\pi = 3.1416$ .

*It should be noticed that  $31416 = 8 \times 3 \times 7 \times 11 \times 17$ .]*

Express in radian measure as fractions of  $\pi$  :

- |                 |                     |                     |                     |
|-----------------|---------------------|---------------------|---------------------|
| 1. $45^\circ$ . | 2. $30^\circ$ .     | 3. $105^\circ$ .    | 4. $22^\circ 30'$ . |
| 5. $18^\circ$ . | 6. $57^\circ 30'$ . | 7. $14^\circ 24'$ . | 8. $78^\circ 45'$ . |

Find numerically the radian measure of the following angles :

- |                      |                       |                      |
|----------------------|-----------------------|----------------------|
| 9. $25^\circ 50'$ .  | 10. $37^\circ 30'$ .  | 11. $82^\circ 30'$ . |
| 12. $68^\circ 45'$ . | 13. $157^\circ 30'$ . | 14. $52^\circ 30'$ . |

Express in sexagesimal measure :

15.  $\frac{3\pi}{4}$ .      16.  $\frac{7\pi}{45}$ .      17.  $\frac{5\pi}{27}$ .      18.  $\frac{5\pi}{24}$ .  
 19.  $\cdot 3927\rho$ .    20.  $\cdot 5236\rho$ .    21.  $\cdot 6545\rho$ .    22.  $2\cdot 8798\rho$ .

Taking  $\pi = \frac{22}{7}$ , find the radian measure of :

23.  $36^\circ 32' 24''$ .      24.  $70^\circ 33' 36''$ .  
 25.  $116^\circ 2' 45\cdot 6''$ .    26.  $171^\circ 41' 50\cdot 4''$ .

27. Taking  $\frac{1}{\pi} = \cdot 31831$ , shew that a radian contains 206265 seconds approximately.

28. Shew that a second is approximately equal to  $\cdot 000048$  of a radian.

67. The angles  $\frac{\pi}{4}$ ,  $\frac{\pi}{3}$ ,  $\frac{\pi}{6}$  are the equivalents in radian measure of the angles  $45^\circ$ ,  $60^\circ$ ,  $30^\circ$  respectively.

Hence the results of Arts. 34 and 35 may be written as follows :

$$\begin{aligned} \sin \frac{\pi}{4} &= \frac{1}{\sqrt{2}}, & \cos \frac{\pi}{4} &= \frac{1}{\sqrt{2}}, & \tan \frac{\pi}{4} &= 1; \\ \sin \frac{\pi}{3} &= \frac{\sqrt{3}}{2}, & \cos \frac{\pi}{3} &= \frac{1}{2}, & \tan \frac{\pi}{3} &= \sqrt{3}; \\ \sin \frac{\pi}{6} &= \frac{1}{2}, & \cos \frac{\pi}{6} &= \frac{\sqrt{3}}{2}, & \tan \frac{\pi}{6} &= \frac{1}{\sqrt{3}}. \end{aligned}$$

*Example.* Find the value of

$$3 \tan^2 \frac{\pi}{6} + \frac{4}{3} \cos^2 \frac{\pi}{6} - \frac{1}{2} \cot^3 \frac{\pi}{4} - \frac{2}{3} \sin^2 \frac{\pi}{3} + \frac{1}{8} \sec^4 \frac{\pi}{3}.$$

$$\begin{aligned} \text{The value} &= 3 \left( \frac{1}{\sqrt{3}} \right)^2 + \frac{4}{3} \left( \frac{\sqrt{3}}{2} \right)^2 - \frac{1}{2} (1)^3 - \frac{2}{3} \left( \frac{\sqrt{3}}{2} \right)^2 + \frac{1}{8} (2)^4 \\ &= \left( 3 \times \frac{1}{3} \right) + \left( \frac{4}{3} \times \frac{3}{4} \right) - \frac{1}{2} - \left( \frac{2}{3} \times \frac{3}{4} \right) + \left( \frac{1}{8} \times 16 \right) \\ &= 1 + 1 - \frac{1}{2} - \frac{1}{2} + 2 = 3. \end{aligned}$$

68. When expressed in radian measure the complement of  $\theta$  is  $\frac{\pi}{2} - \theta$ , and corresponding to the formulæ of Art. 38 we now have relations of the form

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta, \quad \tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta.$$

*Example.* Prove that

$$(\cot \theta + \tan \theta) \cot\left(\frac{\pi}{2} - \theta\right) = \operatorname{cosec}^2\left(\frac{\pi}{2} - \theta\right).$$

$$\begin{aligned} \text{The first side} &= (\cot \theta + \tan \theta) \tan \theta \\ &= \cot \theta \tan \theta + \tan^2 \theta \\ &= 1 + \tan^2 \theta = \sec^2 \theta \\ &= \operatorname{cosec}^2\left(\frac{\pi}{2} - \theta\right). \end{aligned}$$

69. By means of Euc. I. 32, it is easy to find the number of radians in each angle of a regular polygon.

*Example.* Express in radians the interior angle of a regular polygon which has  $n$  sides.

The sum of the *exterior* angles = 4 right angles. [Euc. I. 32 Cor.]

Let  $\theta$  be the number of radians in an exterior angle; then

$$n\theta = 2\pi, \text{ and therefore } \theta = \frac{2\pi}{n}.$$

But interior angle = two right angles - exterior angle

$$= \pi - \theta = \pi - \frac{2\pi}{n}.$$

Thus each interior angle =  $\frac{(n-2)\pi}{n}$ .

### EXAMPLES. VII. b.

Find the numerical value of

- |   |  |
|---|--|
| 1. $\sin \frac{\pi}{3} \cos \frac{\pi}{6} \cot \frac{\pi}{4}.$              | 2. $\tan \frac{\pi}{6} \cot \frac{\pi}{3} \cos \frac{\pi}{4}.$ |
| 3. $\frac{1}{2} \cos \frac{\pi}{3} + 2 \operatorname{cosec} \frac{\pi}{6}.$ | 4. $2 \sin \frac{\pi}{4} + \frac{1}{2} \sec \frac{\pi}{4}.$    |

Find the numerical value of

$$5. \cot^2 \frac{\pi}{6} + 4 \cos^2 \frac{\pi}{4} + 3 \sec^2 \frac{\pi}{6}.$$

$$6. 3 \tan^2 \frac{\pi}{6} - \frac{1}{3} \sin^2 \frac{\pi}{3} - \frac{1}{2} \operatorname{cosec}^2 \frac{\pi}{4} + \frac{4}{3} \cos^2 \frac{\pi}{6}.$$

$$7. \left( \sin \frac{\pi}{6} + \cos \frac{\pi}{6} \right) \left( \sin \frac{\pi}{3} - \cos \frac{\pi}{3} \right) \sec \frac{\pi}{3}.$$

Prove the following identities :

$$8. \sin \theta \sec \left( \frac{\pi}{2} - \theta \right) - \cot \theta \cot \left( \frac{\pi}{2} - \theta \right) = 0.$$

$$9. \sin^2 \left( \frac{\pi}{2} - \theta \right) \operatorname{cosec} \theta - \tan^2 \left( \frac{\pi}{2} - \theta \right) \sin \theta = 0.$$

$$10. \frac{\sin^2 \left( \frac{\pi}{2} - \theta \right)}{\operatorname{cosec} \theta} \cdot \frac{\sec \theta}{\cot \left( \frac{\pi}{2} - \theta \right)} = \cos^2 \theta.$$

$$11. \tan \theta + \tan \left( \frac{\pi}{2} - \theta \right) = \sec \theta \sec \left( \frac{\pi}{2} - \theta \right).$$

$$12. \sec^2 \theta + \sec^2 \left( \frac{\pi}{2} - \theta \right) = (1 + \tan^2 \theta) \sec^2 \left( \frac{\pi}{2} - \theta \right).$$

13. Find the number of radians in each exterior angle of  
(1) a regular octagon, (2) a regular quindecagon.

14. Find the number of radians in each interior angle of  
(1) a regular dodecagon, (2) a regular heptagon.

15. Shew that

$$\tan^2 \frac{\pi}{3} - \cot^2 \frac{\pi}{3} = \frac{\cos^2 \frac{\pi}{6} - \cos^2 \frac{\pi}{3}}{\cos^2 \frac{\pi}{3} \cos^2 \frac{\pi}{6}}.$$

16. Shew that the sum of the squares of

$$\sin \theta + \sin \left( \frac{\pi}{2} - \theta \right) \text{ and } \cos \theta - \cos \left( \frac{\pi}{2} - \theta \right)$$

is equal to 2.

70. To prove that the radian measure of any angle at the centre of a circle is expressed by the fraction  $\frac{\text{subtending arc}}{\text{radius}}$ .


Let  $\angle AOC$  be any angle at the centre of a circle, and  $\angle AOB$  a radian; then

radian measure of  $\angle AOC$

$$= \frac{\angle AOC}{\angle AOB}$$

$$= \frac{\text{arc } AC}{\text{arc } AB}$$

$$= \frac{\text{arc } AC}{\text{radius}},$$



since  $\text{arc } AB = \text{radius}$ ;

that is, the radian measure of  $\angle AOC = \frac{\text{subtending arc}}{\text{radius}}$ .

71. If  $a$  be the length of the arc which subtends an angle of  $\theta$  radians at the centre of a circle of radius  $r$ , we have seen in the preceding article that

$$\theta = \frac{a}{r}, \text{ and therefore } a = r\theta.$$

The fraction  $\frac{\text{arc}}{\text{radius}}$  is usually called the *circular measure* of the angle at the centre of the circle subtended by the arc.

The *circular measure* of an angle is therefore equal to its *radian measure*, each denoting the *number of radians* contained in the angle. We have preferred to use the term *radian measure* exclusively, in order to keep prominently in view the unit of measurement, namely the radian.

NOTE. The term *circular measure* is a survival from the times when Mathematicians spoke of the trigonometrical functions of the *arc*. [See page 80.]

*Example 1.* Find the angle subtended by an arc of 7.5 feet at the centre of a circle whose radius is 5 yards.

Let the angle contain  $\theta$  radians; then

$$\theta = \frac{\text{arc}}{\text{radius}} = \frac{7.5}{15} = \frac{1}{2}.$$

Thus the angle is half a radian.



*Example 2.* In running a race at a uniform speed on a circular course, a man in each minute traverses an arc of a circle which subtends  $2\frac{2}{7}$  radians at the centre of the course. If each lap is 792 yards, how long does he take to run a mile?  $\left[\pi = \frac{22}{7}\right]$ .

Let  $r$  yards be the radius of the circle; then

$$2\pi r = \text{circumference} = 792;$$

$$\therefore r = \frac{792}{2\pi} = \frac{792 \times 7}{2 \times 22} = 126.$$

Let  $a$  yards be the length of the arc traversed in each minute; then from the formula  $a = r\theta$ ,

$$a = 126 \times 2\frac{2}{7} = \frac{126 \times 20}{7} = 360;$$

that is, the man runs 360 yds. in each minute.

$$\therefore \text{the time} = \frac{1760}{360} \text{ or } \frac{44}{9} \text{ minutes.}$$

Thus the time is 4 min.  $53\frac{1}{3}$  sec.

*Example 3.* Find the radius of a globe such that the distance measured along its surface between two places on the same meridian whose latitudes differ by  $1^\circ 10'$  may be 1 inch, reckoning that  $\pi = \frac{22}{7}$ .

Let the adjoining figure represent a section of the globe through the meridian on which the two places  $P$  and  $Q$  lie. Let  $O$  be the centre, and denote the radius by  $r$  inches.

$$\text{Now } \frac{\text{arc } PQ}{\text{radius}} = \text{number of radians in } \angle POQ;$$

but  $\text{arc } PQ = 1$  inch, and  $\angle POQ = 1^\circ 10'$ ;


$$\therefore \frac{1}{r} = \text{number of radians in } 1\frac{1}{6}^\circ$$

$$= 1\frac{1}{6} \times \frac{\pi}{180} = \frac{7}{6} \times \frac{22}{7} \times \frac{1}{180} = \frac{11}{540};$$

whence

$$r = \frac{540}{11} = 49\frac{1}{11}.$$

Thus the radius is  $49\frac{1}{11}$  inches.



## EXAMPLES. VII. c.

1. Find the radian measure of the angle subtended by an arc of 1.6 yards at the centre of a circle whose radius is 24 feet.

2. An angle whose circular measure is .73 subtends at the centre of a circle an arc of 219 feet; find the radius of the circle.

3. An angle at the centre of a circle whose radius is 2.5 yards is subtended by an arc of 7.5 feet; what is the angle?

4. What is the length of the arc which subtends an angle of 1.625 radians at the centre of a circle whose radius is 3.6 yards?

5. An arc of 17 yds. 1 ft. 3 in. subtends at the centre of a circle an angle of 1.9 radians; find the radius of the circle in inches.

6. The flywheel of an engine makes 35 revolutions in a second; how long will it take to turn through 5 radians?  $\left[\pi = \frac{22}{7}\right]$ .

7. The large hand of a clock is 2 ft. 4 in. long; how many inches does its extremity move in 20 minutes?  $\left[\pi = \frac{22}{7}\right]$ .

8. A horse is tethered to a stake; how long must the rope be in order that, when the horse has moved through 52.36 yards at the extremity of the rope, the angle traced out by the rope may be 75 degrees?

9. Find the length of an arc which subtends 1 minute at the centre of the earth, supposed to be a sphere of diameter 7920 miles.

10. Find the number of seconds in the angle subtended at the centre of a circle of radius 1 mile by an arc  $5\frac{1}{2}$  inches long.

11. Two places on the same meridian are 145.2 miles apart; find their difference in latitude, taking  $\pi = \frac{22}{7}$ , and the earth's diameter as 7920 miles.

12. Find the radius of a globe such that the distance measured along its surface between two places on the same meridian whose latitudes differ by  $1\frac{3}{11}^\circ$  may be 1 foot, taking  $\pi = \frac{22}{7}$ .

**MISCELLANEOUS EXAMPLES. B.**

1. Express in degrees the angle whose circular measure is  $\cdot 15708$ .

2. If  $C=90^\circ$ ,  $A=30^\circ$ ,  $c=110$ , find  $b$  to two decimal places.

3. Find the number of degrees in the unit angle when the angle  $\frac{12\pi}{25}$  is represented by  $1\frac{3}{5}$ .

4. What is the radius of the circle in which an arc of 1 inch subtends an angle of  $1'$  at the centre?

5. Prove that

$$(1) \quad (\sin a + \cos a)(\tan a + \cot a) = \sec a + \operatorname{cosec} a;$$

$$(2) \quad (\sqrt{3}+1)(3-\cot 30^\circ) = \tan^3 60^\circ - 2 \sin 60^\circ.$$

6. Find the angle of elevation of the sun when a chimney 60 feet high throws a shadow  $20\sqrt{3}$  yards long.

7. Prove the identities :

$$(1) \quad (\tan \theta + 2)(2 \tan \theta + 1) = 5 \tan \theta + 2 \sec^2 \theta;$$

$$(2) \quad 1 + \frac{\cot^2 a}{1 + \operatorname{cosec} a} = \operatorname{cosec} a.$$

8. One angle of a triangle is  $45^\circ$  and another is  $\frac{5\pi}{8}$  radians; express the third angle both in sexagesimal and radian measure.

9. The number of degrees in an angle exceeds 14 times the number of radians in it by 51. Taking  $\pi = \frac{22}{7}$ , find the sexagesimal measure of the angle.

10. If  $B=30^\circ$ ,  $C=90^\circ$ ,  $b=6$ , find  $a$ ,  $c$ , and the perpendicular from  $C$  on the hypotenuse.

11. Shew that

$$(1) \quad \cot \theta + \cot \left( \frac{\pi}{2} - \theta \right) = \operatorname{cosec} \theta \operatorname{cosec} \left( \frac{\pi}{2} - \theta \right);$$

$$(2) \quad \operatorname{cosec}^2 \theta + \operatorname{cosec}^2 \left( \frac{\pi}{2} - \theta \right) = \operatorname{cosec}^2 \theta \operatorname{cosec}^2 \left( \frac{\pi}{2} - \theta \right).$$

12. The angle of elevation of the top of a pillar is  $30^\circ$ , and on approaching 20 feet nearer it is  $60^\circ$ : find the height of the pillar.

---

13. Shew that  $\tan^2 A - \sin^2 A = \sin^4 A \sec^2 A$ .

14. In a triangle the angle  $A$  is  $3x$  degrees, the angle  $B$  is  $x$  grades, and the angle  $C$  is  $\frac{\pi x}{300}$  radians: find the number of degrees in each of the angles.

15. Find the numerical value of

$$\sin^3 60^\circ \cot 30^\circ - 2 \sec^2 45^\circ + 3 \cos 60^\circ \tan 45^\circ - \tan^2 60^\circ.$$

16. Prove the identities :

$$(1) \quad (1 + \tan A)^2 + (1 + \cot A)^2 = (\sec A + \operatorname{cosec} A)^2;$$

$$(2) \quad (\sec a - 1)^2 - (\tan a - \sin a)^2 = (1 - \cos a)^2.$$

17. Which of the following statements is possible and which impossible?

$$(1) \quad \operatorname{cosec} \theta = \frac{a^2 + b^2}{2ab}; \quad (2) \quad 2 \sin \theta = a + \frac{1}{a}.$$

18. A balloon leaves the earth at the point  $A$  and rises at a uniform pace. At the end of 1.5 minutes an observer stationed at a distance of 660 feet from  $A$  finds the angular elevation of the balloon to be  $60^\circ$ ; at what rate in miles per hour is the balloon rising?

---

19. Find the number of radians in the angles of a triangle which are in arithmetical progression, the least angle being  $36^\circ$ .

20. Shew that

$$\sin^2 a \sec^2 \beta + \tan^2 \beta \cos^2 a = \sin^2 a + \tan^2 \beta.$$

21. In the triangle  $ABC$  if  $A = 42^\circ$ ,  $B = 116^\circ 33'$ , find the perpendicular from  $C$  upon  $AB$  produced; given

$$c = 55, \quad \tan 42^\circ = .9, \quad \tan 63^\circ 27' = 2.$$

22. Prove the identities :

$$(1) \cot a + \frac{\sin a}{1 + \cos a} = \operatorname{cosec} a;$$

$$(2) \operatorname{cosec} a (\sec a - 1) - \cot a (1 - \cos a) = \tan a - \sin a.$$

23. Shew that  $\left(\frac{1 + \cot 60^\circ}{1 - \cot 60^\circ}\right)^2 = \frac{1 + \cos 30^\circ}{1 - \cos 30^\circ}.$

24. A man walking N.W. sees a windmill which bears N.  $15^\circ$  W. In half-an-hour he reaches a place which he knows to be W.  $15^\circ$  S. of the windmill and a mile away from it. Find his rate of walking and his distance from the windmill at the first observation.

25. Find the number of radians in the complement of  $\frac{3\pi}{8}.$

26. Solve the equations :

$$(1) 3 \sin \theta + 4 \cos^2 \theta = 4\frac{1}{2}; \quad (2) \tan \theta + \sec 30^\circ = \cot \theta.$$

27. If  $5 \tan a = 4$ , find the value of

$$\frac{5 \sin a - 3 \cos a}{\sin a + 2 \cos a}.$$

28. Prove that

$$\frac{1 - \sin A \cos A}{\cos A (\sec A - \operatorname{cosec} A)} \times \frac{\sin^2 A - \cos^2 A}{\sin^3 A + \cos^3 A} = \sin A.$$

29. Find the distance of an observer from the top of a cliff which is 195.2 yards high, given that the angle of elevation is  $77^\circ 26'$ , and that  $\sin 77^\circ 26' = .976$ .


30. A horse is tethered to a stake by a rope 27 feet long. If the horse moves along the circumference of a circle always keeping the rope tight, find how far it will have gone when the rope has traced out an angle of  $70^\circ$ .  $\left[\pi = \frac{22}{7}\right].$

## CHAPTER VIII.

### TRIGONOMETRICAL RATIOS OF ANGLES OF ANY MAGNITUDE.

72. In the present chapter we shall find it necessary to take account not only of the magnitude of straight lines, but also of the direction in which they are measured.


Let  $O$  be a fixed point in a horizontal line  $XX'$ , then the position of any other point  $P$  in the line, whose distance from  $O$  is a given length  $a$ , will not be determined unless we know on which side of  $O$  the point  $P$  lies.



But there will be no ambiguity if it is agreed that distances measured in one direction are positive and distances measured in the opposite direction are negative.

Hence the following **Convention of Signs** is adopted :

*lines measured from  $O$  to the right are positive,*  
*lines measured from  $O$  to the left are negative.*



Thus in the above figure, if  $P$  and  $Q$  are two points on the line  $XX'$  at a distance  $a$  from  $O$ , their positions are indicated by the statements

$$OP = +a, \quad OQ = -a.$$


73. A similar *convention of signs* is used in the case of a plane surface.

Let  $O$  be any point in the plane; through  $O$  draw two straight lines  $XX'$  and  $YY'$  in the horizontal and vertical direction respectively, thus dividing the plane into four *quadrants*.



Then it is universally agreed to consider that


- (1) *horizontal lines to the right of  $YY'$  are positive, horizontal lines to the left of  $YY'$  are negative;*
- (2) *vertical lines above  $XX'$  are positive, vertical lines below  $XX'$  are negative.*



Thus  $OM_1$ ,  $OM_4$  are positive,  $OM_2$ ,  $OM_3$  are negative;  
 $M_1P_1$ ,  $M_2P_2$  are positive,  $M_3P_3$ ,  $M_4P_4$  are negative.

**74. Convention of Signs for Angles.** In Art. 2 an angle has been defined as the amount of revolution which the radius vector makes in passing from its initial to its final position.

In the adjoining figure the straight line  $OP$  may be supposed to have arrived at its present position from the position occupied by  $OA$  by revolution about the point  $O$  in *either* of the two directions indicated by the arrows. The angle  $AOP$  may thus be regarded in two senses according as we suppose the revolution to have been in the same direction as the hands of a clock or in the opposite direction. To distinguish between these cases we adopt the following convention:







*when the revolution of the radius vector is counter-clockwise the angle is positive,*

*when the revolution is clockwise the angle is negative.*



### Trigonometrical Ratios of any Angle.

75. Let  $XX'$  and  $YY'$  be two straight lines intersecting at right angles in  $O$ , and let a radius vector starting from  $O$  revolve in either direction till it has traced out an angle  $A$ , taking up the position  $OP$ .



From  $P$  draw  $PM$  perpendicular to  $XX'$ ; then in the right-angled triangle  $OPM$ , due regard being paid to the signs of the lines,

$$\sin A = \frac{MP}{OP}, \quad \operatorname{cosec} A = \frac{OP}{MP},$$


$$\cos A = \frac{OM}{OP}, \quad \sec A = \frac{OP}{OM},$$

$$\tan A = \frac{MP}{OM}, \quad \cot A = \frac{OM}{MP}.$$

*The radius vector  $OP$  which only fixes the boundary of the angle is considered to be always positive.*

From these definitions it will be seen that any trigonometrical function will be positive or negative according as the fraction which expresses its value has the numerator and denominator of the same sign or of opposite sign.

76. The four diagrams of the last article may be conveniently included in one.



With centre  $O$  and fixed radius let a circle be described; then the diameters  $XX'$  and  $YY'$  divide the circle into four quadrants  $XOY$ ,  $YOX'$ ,  $X'OY'$ ,  $Y'OX$ , named *first*, *second*, *third*, *fourth* respectively.

Let the positions of the radius vector in the four quadrants be denoted by  $OP_1$ ,  $OP_2$ ,  $OP_3$ ,  $OP_4$ , and let perpendiculars  $P_1M_1$ ,  $P_2M_2$ ,  $P_3M_3$ ,  $P_4M_4$  be drawn to  $XX'$ ; then it will be seen that in the first quadrant all the lines are positive and therefore all the functions of  $A$  are positive.

In the second quadrant,  $OP_2$  and  $M_2P_2$  are positive,  $OM_2$  is negative; hence  $\sin A$  is positive,  $\cos A$  and  $\tan A$  are negative.

In the third quadrant,  $OP_3$  is positive,  $OM_3$  and  $M_3P_3$  are negative; hence  $\tan A$  is positive,  $\sin A$  and  $\cos A$  are negative.

In the fourth quadrant,  $OP_4$  and  $OM_4$  are positive,  $M_4P_4$  is negative; hence  $\cos A$  is positive,  $\sin A$  and  $\tan A$  are negative.

77. The following diagrams shew the *signs* of the trigonometrical functions in the four quadrants. It will be sufficient to consider the three principal functions only.

<i>sine</i>	<i>cosine</i>	<i>tangent</i>
<div style="display: flex; align-items: center; justify-content: center;"> <div style="border-right: 1px solid black; padding: 5px; margin-right: 5px;">+</div> <div style="padding: 5px;">+</div> </div> <div style="display: flex; align-items: center; justify-content: center;"> <div style="border-right: 1px solid black; padding: 5px; margin-right: 5px;">-</div> <div style="padding: 5px;">-</div> </div>	<div style="display: flex; align-items: center; justify-content: center;"> <div style="border-right: 1px solid black; padding: 5px; margin-right: 5px;">-</div> <div style="padding: 5px;">+</div> </div> <div style="display: flex; align-items: center; justify-content: center;"> <div style="border-right: 1px solid black; padding: 5px; margin-right: 5px;">-</div> <div style="padding: 5px;">+</div> </div>	<div style="display: flex; align-items: center; justify-content: center;"> <div style="border-right: 1px solid black; padding: 5px; margin-right: 5px;">-</div> <div style="padding: 5px;">+</div> </div> <div style="display: flex; align-items: center; justify-content: center;"> <div style="border-right: 1px solid black; padding: 5px; margin-right: 5px;">+</div> <div style="padding: 5px;">-</div> </div>

The diagram below exhibits the same results in another useful form.

<b>sine positive</b> <i>cosine negative</i> <i>tangent negative</i>	<b>all the</b> <b>ratios positive</b>
<b>tangent positive</b> <i>sine negative</i> <i>cosine negative</i>	<b>cosine positive</b> <i>sine negative</i> <i>tangent negative</i>

78. When an angle is increased or diminished by any multiple of four right angles, the radius vector is brought back again into the same position after one or more revolutions. There are thus an infinite number of angles which have the same boundary line. Such angles are called **coterminal angles**.

If  $n$  is *any* integer, all the angles coterminal with  $A$  may be represented by  $n \cdot 360^\circ + A$ . Similarly, in radian measure all the angles coterminal with  $\theta$  may be represented by  $2n\pi + \theta$ .

From the definitions of Art. 75, we see that the position of the boundary line is alone sufficient to determine the trigonometrical ratios of the angle; hence *all coterminal angles have the same trigonometrical ratios*.

For instance,  $\sin(n \cdot 360^\circ + 45^\circ) = \sin 45^\circ = \frac{1}{\sqrt{2}}$ ;


and  $\cos\left(2n\pi + \frac{\pi}{6}\right) = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$ .

*Example.* Draw the boundary lines of the angles  $780^\circ$ ,  $-130^\circ$ ,  $-400^\circ$ , and in each case state which of the trigonometrical functions are negative.


(1) Since  $780 = (2 \times 360) + 60$ , the radius vector has to make two complete revolutions and then turn through  $60^\circ$ . Thus the boundary line is in the first quadrant, so that all the functions are positive.



(2) Here the radius vector has to revolve through  $130^\circ$  in the negative direction. The boundary line is thus in the third quadrant, and since  $OM$  and  $MP$  are negative, the sine, cosine, cosecant, and secant are negative.



(3) Since  $-400 = -(360 + 40)$ , the radius vector has to make one complete revolution in the negative direction and then turn through  $40^\circ$ . The boundary line is thus in the fourth quadrant, and since  $MP$  is negative, the sine, tangent, cosecant, and cotangent are negative.



### EXAMPLES. VIII. a.

State the quadrant in which the radius vector lies after describing the following angles:

- |                       |                       |                        |                         |
|-----------------------|-----------------------|------------------------|-------------------------|
| 1. $135^\circ$ .      | 2. $265^\circ$ .      | 3. $-315^\circ$ .      | 4. $-120^\circ$ .       |
| 5. $\frac{2\pi}{3}$ . | 6. $\frac{5\pi}{6}$ . | 7. $\frac{10\pi}{3}$ . | 8. $-\frac{11\pi}{4}$ . |

For each of the following angles state which of the three principal trigonometrical functions are positive.

- |                         |                         |                          |
|-------------------------|-------------------------|--------------------------|
| 9. $470^\circ$ .        | 10. $330^\circ$ .       | 11. $575^\circ$ .        |
| 12. $-230^\circ$ .      | 13. $-620^\circ$ .      | 14. $-1200^\circ$ .      |
| 15. $-\frac{4\pi}{3}$ . | 16. $\frac{13\pi}{6}$ . | 17. $-\frac{13\pi}{6}$ . |

In each of the following cases write down the smallest positive coterminal angle, and the value of the expression.

$$18. \sin 420^\circ. \quad 19. \cos 390^\circ. \quad 20. \tan (-315^\circ).$$

$$21. \sec 405^\circ. \quad 22. \operatorname{cosec} (-330^\circ). \quad 23. \operatorname{cosec} 4380^\circ.$$

$$24. \cot \frac{17\pi}{4}. \quad 25. \sec \frac{25\pi}{3}. \quad 26. \tan \left(-\frac{5\pi}{3}\right).$$

79. Since the definitions of the functions given in Art. 75 are applicable to angles of any magnitude, positive or negative, it follows that all relations derived from these definitions must be true universally. Thus we shall find that the fundamental formulæ given in Art. 29 hold in all cases; that is,

$$\sin A \times \operatorname{cosec} A = 1, \quad \cos A \times \sec A = 1, \quad \tan A \times \cot A = 1;$$

$$\tan A = \frac{\sin A}{\cos A}, \quad \cot A = \frac{\cos A}{\sin A};$$

$$\sin^2 A + \cos^2 A = 1,$$

$$1 + \tan^2 A = \sec^2 A,$$

$$1 + \cot^2 A = \operatorname{cosec}^2 A.$$

It will be useful practice for the student to test the truth of these formulæ for different positions of the boundary line of the angle  $A$ . We shall give one illustration.

80. Let the radius vector revolve from its initial position  $OX$  till it has traced out an angle  $A$  and come into the position  $OP$  indicated in the figure. Draw  $PM$  perpendicular to  $XX'$ .

In the right-angled triangle  $OMP$ ,


$$MP^2 + OM^2 = OP^2 \dots\dots (1).$$

Divide each term by  $OP^2$ ; thus

$$\left(\frac{MP}{OP}\right)^2 + \left(\frac{OM}{OP}\right)^2 = 1;$$

that is,

$$\sin^2 A + \cos^2 A = 1.$$



Divide each term of (1) by  $OM^2$ ; thus

$$\left(\frac{MP}{OM}\right)^2 + 1 = \left(\frac{OP}{OM}\right)^2;$$

that is,  $\tan^2 A + 1 = \sec^2 A$ .

Divide each term of (1) by  $MP^2$ ; thus

$$1 + \left(\frac{OM}{MP}\right)^2 = \left(\frac{OP}{MP}\right)^2;$$

that is,  $1 + \cot^2 A = \operatorname{cosec}^2 A$ .

It thus appears that the truth of these relations depends only on the statement  $OP^2 = MP^2 + OM^2$  in the right-angled triangle  $OMP$ , and this will be the case in whatever quadrant  $OP$  lies.

NOTE.  $OM^2$  is positive, although the line  $OM$  in the figure is negative.

81. In the statement  $\cos A = \sqrt{1 - \sin^2 A}$ , either the positive or the negative sign may be placed before the radical. The sign of the radical hitherto has always been taken positively, because we have restricted ourselves to the consideration of acute angles. It will sometimes be necessary to examine which sign must be taken before the radical in any particular case.

*Example 1.* Given  $\cos 126^\circ 53' = -\frac{3}{5}$ , find  $\sin 126^\circ 53'$  and  $\cot 126^\circ 53'$ .

Since  $\sin^2 A + \cos^2 A = 1$  for angles of any magnitude, we have

$$\sin A = \pm \sqrt{1 - \cos^2 A}.$$

Denote  $126^\circ 53'$  by  $A$ ; then the boundary line of  $A$  lies in the second quadrant, and therefore  $\sin A$  is positive. Hence the sign  $+$  must be placed before the radical;

$$\therefore \sin 126^\circ 53' = +\sqrt{1 - \frac{9}{25}} = +\sqrt{\frac{16}{25}} = \frac{4}{5};$$

$$\cot 126^\circ 53' = \frac{\cos 126^\circ 53'}{\sin 126^\circ 53'} = \left(-\frac{3}{5}\right) \div \left(\frac{4}{5}\right) = -\frac{3}{4}.$$

The same results may also be obtained by the method used in the following example. The appropriate signs of the lines are shewn in the figure.



*Example 2.* If  $\tan A = -\frac{15}{8}$ , find  $\sin A$  and  $\cos A$ .


The boundary line of  $A$  will lie either in the second or in the fourth quadrant, as  $OP$  or  $OP'$ . In either position,

$$\begin{aligned}\text{the radius vector} &= \sqrt{(15)^2 + (8)^2} \\ &= \sqrt{289} = 17.\end{aligned}$$

$$\text{Hence } \sin XOP = \frac{15}{17}, \quad \cos XOP = -\frac{8}{17};$$

$$\text{and } \sin XOP' = -\frac{15}{17}, \quad \cos XOP' = \frac{8}{17}.$$

Thus corresponding to  $\tan A$ , there are two values of  $\sin A$  and two values of  $\cos A$ . If however it is known in which quadrant the boundary line of  $A$  lies,  $\sin A$  and  $\cos A$  have each a single value.



### EXAMPLES. VIII. b.

1. Given  $\sin 120^\circ = \frac{\sqrt{3}}{2}$ , find  $\tan 120^\circ$ .
2. Given  $\tan 135^\circ = -1$ , find  $\sin 135^\circ$ .
3. Find  $\cos 240^\circ$ , given that  $\tan 240^\circ = \sqrt{3}$ .
4. If  $A = 202^\circ 37'$  and  $\sin A = -\frac{5}{13}$ , find  $\cos A$  and  $\cot A$ .
5. If  $A = 143^\circ 8'$  and  $\operatorname{cosec} A = 1\frac{2}{3}$ , find  $\sec A$  and  $\tan A$ .
6. If  $A = 216^\circ 52'$  and  $\cos A = -\frac{4}{5}$ , find  $\cot A$  and  $\sin A$ .
7. Given  $\sec \frac{2\pi}{3} = -2$ , find  $\sin \frac{2\pi}{3}$  and  $\cot \frac{2\pi}{3}$ .
8. Given  $\sin \frac{5\pi}{4} = -\frac{1}{\sqrt{2}}$ , find  $\tan \frac{5\pi}{4}$  and  $\sec \frac{5\pi}{4}$ .
9. If  $\cos A = \frac{12}{13}$ , find  $\sin A$  and  $\tan A$ .



## CHAPTER IX.

### VARIATIONS OF THE TRIGONOMETRICAL FUNCTIONS.

82. A CAREFUL perusal of the following remarks will render the explanations which follow more easily intelligible.

Consider the fraction  $\frac{a}{x}$  in which the numerator  $a$  has a *certain fixed value* and the denominator  $x$  is a *quantity subject to change*; then it is clear that the smaller  $x$  becomes the larger does the value of the fraction  $\frac{a}{x}$  become. For instance

$$\frac{\frac{a}{1}}{10} = 10a, \quad \frac{\frac{a}{1}}{1000} = 1000a, \quad \frac{\frac{a}{1}}{10000000} = 10000000a.$$

By making the denominator  $x$  sufficiently small the value of the fraction  $\frac{a}{x}$  can be made as large as we please; that is, as  $x$  approaches to the value 0, the fraction  $\frac{a}{x}$  becomes infinitely great.

The symbol  $\infty$  is used to express a quantity infinitely great, or more shortly *infinity*, and the above statement is concisely written

$$\text{when } x=0, \text{ the limit of } \frac{a}{x} = \infty.$$

Again, if  $x$  is a quantity which gradually increases and finally becomes infinitely large, the fraction  $\frac{a}{x}$  becomes infinitely small; that is,

$$\text{when } x=\infty, \text{ the limit of } \frac{a}{x} = 0.$$

**83. DEFINITION.** If  $y$  is a function of  $x$ , and if when  $x$  approaches nearer and nearer to the fixed quantity  $a$ , the value of  $y$  approaches nearer and nearer to the fixed quantity  $b$  and can be made to differ from it by as little as we please, then  $b$  is called the **limiting value** or the **limit** of  $y$  when  $x=a$ .

#### 84. Trigonometrical Functions of $0^\circ$ .

Let  $XOP$  be an angle traced out by a radius vector  $OP$  of fixed length.



Draw  $PM$  perpendicular to  $OX$ ; then

$$\sin POM = \frac{MP}{OP}.$$

If we suppose the angle  $POM$  to be gradually decreasing,  $MP$  will also gradually decrease, and if  $OP$  ultimately come into coincidence with  $OM$  the angle  $POM$  vanishes and  $MP=0$ .

Hence 
$$\sin 0^\circ = \frac{0}{OP} = 0.$$

Again,  $\cos POM = \frac{OM}{OP}$ ; but when the angle  $POM$  vanishes  $OP$  becomes coincident with  $OM$ .

Hence 
$$\cos 0^\circ = \frac{OM}{OM} = 1.$$

Also when the angle  $POM$  vanishes,

$$\tan 0^\circ = \frac{0}{OM} = 0.$$


And 
$$\operatorname{cosec} 0^\circ = \frac{1}{\sin 0^\circ} = \frac{1}{0} = \infty;$$

$$\sec 0^\circ = \frac{1}{\cos 0^\circ} = \frac{1}{1} = 1;$$

$$\cot 0^\circ = \frac{1}{\tan 0^\circ} = \frac{1}{0} = \infty.$$

85. Trigonometrical Functions of  $90^\circ$  or  $\frac{\pi}{2}$ .

Let  $XOP$  be an angle traced out by a radius vector of fixed length.



Draw  $PM$  perpendicular to  $OX$ , and  $OY$  perpendicular to  $OX$ .

By definition,

$$\sin POM = \frac{MP}{OP}, \quad \cos POM = \frac{OM}{OP}, \quad \tan POM = \frac{MP}{OM}.$$

If we suppose the angle  $POM$  to be gradually increasing,  $MP$  will gradually increase and  $OM$  decrease. When  $OP$  comes into coincidence with  $OY$  the angle  $POM$  becomes equal to  $90^\circ$ , and  $OM$  vanishes, while  $MP$  becomes equal to  $OP$ .

$$\begin{aligned} \text{Hence} \quad \sin 90^\circ &= \frac{OP}{OP} = 1; \\ \cos 90^\circ &= \frac{0}{OP} = 0; \\ \tan 90^\circ &= \frac{MP}{OM} = \frac{OP}{0} = \infty. \end{aligned}$$


$$\text{And} \quad \cot 90^\circ = \frac{1}{\tan 90^\circ} = \frac{1}{\infty} = 0;$$

$$\sec 90^\circ = \frac{1}{\cos 90^\circ} = \frac{1}{0} = \infty;$$

$$\operatorname{cosec} 90^\circ = \frac{1}{\sin 90^\circ} = 1.$$

86. To trace the changes in sign and magnitude of  $\sin A$  as  $A$  increases from  $0^\circ$  to  $360^\circ$ .

Let  $XX'$  and  $YY'$  be two straight lines intersecting at right angles in  $O$ .



With centre  $O$  and any radius  $OP$  describe a circle, and suppose the angle  $A$  to be traced out by the revolution of  $OP$  through the four quadrants starting from  $OX$ .

Draw  $PM$  perpendicular to  $OX$  and let  $OP=r$ ; then

$$\sin A = \frac{MP}{r},$$

and since  $r$  does not alter in sign or magnitude, we have only to consider the changes of  $MP$  as  $P$  moves round the circle.

When  $A=0^\circ$ ,  $MP=0$ , and  $\sin 0^\circ = \frac{0}{r} = 0$ .

*In the first quadrant*,  $MP$  is positive and increasing;

$\therefore \sin A$  is positive and increasing.

When  $A=90^\circ$ ,  $MP=r$ , and  $\sin 90^\circ = \frac{r}{r} = 1$ .

*In the second quadrant*,  $MP$  is positive and decreasing;

$\therefore \sin A$  is positive and decreasing.

When  $A=180^\circ$ ,  $MP=0$ , and  $\sin 180^\circ = \frac{0}{r} = 0$ .

*In the third quadrant,  $MP$  is negative and increasing;*

$\therefore \sin A$  is negative and increasing.

When  $A = 270^\circ$ ,  $MP$  is equal to  $r$ , but is negative; hence

$$\sin 270^\circ = -\frac{r}{r} = -1.$$

*In the fourth quadrant,  $MP$  is negative and decreasing;*

$\therefore \sin A$  is negative and decreasing.

When  $A = 360^\circ$ ,  $MP = 0$ , and  $\sin 360^\circ = \frac{0}{r} = 0$ .

87. The results of the previous article are concisely shewn in the following diagram:

		$\sin 90^\circ = 1$	
	$\sin A$ positive and decreasing	$\sin A$ positive and increasing	
$\sin 180^\circ = 0$			$\sin 0^\circ = 0$
	$\sin A$ negative and increasing	$\sin A$ negative and decreasing	
		$\sin 270^\circ = -1$	

88. We leave as an exercise to the student the investigation of the changes in sign and magnitude of  $\cos A$  as  $A$  increases from  $0^\circ$  to  $360^\circ$ . The following diagram exhibits these changes.

		$\cos 90^\circ = 0$	
	$\cos A$ negative and increasing	$\cos A$ positive and decreasing	
$\cos 180^\circ = -1$			$\cos 0^\circ = 1$
	$\cos A$ negative and decreasing	$\cos A$ positive and increasing	
		$\cos 270^\circ = 0$	

89. To trace the changes in sign and magnitude of  $\tan A$  as  $A$  increases from  $0^\circ$  to  $360^\circ$ .

With the figure of Art. 86,  $\tan A = \frac{MP}{OM}$ , and its changes will therefore depend on those of  $MP$  and  $OM$ .

When  $A = 0^\circ$ ,  $MP = 0$ ,  $OM = r$ ;  $\therefore \tan 0^\circ = \frac{0}{r} = 0$ .

*In the first quadrant,*

$MP$  is positive and increasing,

$OM$  is positive and decreasing;

$\therefore \tan A$  is positive and increasing.

When  $A = 90^\circ$ ,  $MP = r$ ,  $OM = 0$ ;  $\therefore \tan 90^\circ = \frac{r}{0} = \infty$ .

*In the second quadrant,*

$MP$  is positive and decreasing,

$OM$  is negative and increasing;

$\therefore \tan A$  is negative and decreasing.

When  $A = 180^\circ$ ,  $MP = 0$ ;  $\therefore \tan 180^\circ = 0$ .

*In the third quadrant,*

$MP$  is negative and increasing,

$OM$  is negative and decreasing;

$\therefore \tan A$  is positive and increasing.

When  $A = 270^\circ$ ,  $OM = 0$ ;  $\therefore \tan 270^\circ = \infty$ .

*In the fourth quadrant,*

$MP$  is negative and decreasing,

$OM$  is positive and increasing;

$\therefore \tan A$  is negative and decreasing.

When  $A = 360^\circ$ ,  $MP = 0$ ;  $\therefore \tan 360^\circ = 0$ .

NOTE. When the numerator of a fraction changes continually from a small positive to a small negative quantity the fraction changes sign by passing through the value 0. When the denominator changes continually from a small positive to a small negative quantity the fraction changes sign by passing through the value  $\infty$ . For instance, as  $A$  passes through the value  $90^\circ$ ,  $OM$  changes from a small positive to a small negative quantity, hence  $\frac{OM}{OP}$ , that is  $\cos A$ , changes sign by passing through the value 0, while  $\frac{PM}{OM}$ , that is  $\tan A$ , changes sign by passing through the value  $\infty$ .

90. The results of Art. 89 are shewn in the following diagram :

$\tan 90^\circ = \infty$	
<i>tan A negative</i> <i>and decreasing</i>	<i>tan A positive</i> <i>and increasing</i>
$\tan 180^\circ = 0$	$\tan 0^\circ = 0$
<i>tan A positive</i> <i>and increasing</i>	<i>tan A negative</i> <i>and decreasing</i>
$\tan 270^\circ = \infty$	

The student will now have no difficulty in tracing the variations in sign and magnitude of the other functions.

91. In Arts. 86 and 89 we have seen that the variations of the trigonometrical functions of the angle  $\angle O P$  depend on the position of  $P$  as  $P$  moves round the circumference of the circle. On this account the trigonometrical functions of an angle are called **circular functions**. This name is one that we shall use frequently.

### EXAMPLES. IX.

Trace the changes in sign and magnitude of

1.  $\cot A$ , between  $0^\circ$  and  $360^\circ$ .
2.  $\operatorname{cosec} \theta$ , between 0 and  $\pi$ .
3.  $\cos \theta$ , between  $\pi$  and  $2\pi$ .
4.  $\tan A$ , between  $-90^\circ$  and  $-270^\circ$ .
5.  $\sec \theta$ , between  $\frac{\pi}{2}$  and  $\frac{3\pi}{2}$ .


Find the value of

6.  $\cos 0^\circ \sin^2 270^\circ - 2 \cos 180^\circ \tan 45^\circ$ .
7.  $3 \sin 0^\circ \sec 180^\circ + 2 \operatorname{cosec} 90^\circ - \cos 360^\circ$ .
8.  $2 \sec^2 \pi \cos 0 + 3 \sin^3 \frac{3\pi}{2} - \operatorname{cosec} \frac{\pi}{2}$ .
9.  $\tan \pi \cos \frac{3\pi}{2} + \sec 2\pi - \operatorname{cosec} \frac{3\pi}{2}$ .



*Note on the old definitions of the Trigonometrical Functions.*

Formerly, Mathematicians considered the trigonometrical functions with reference to the *arc* of a given circle, and did not regard them as *ratios* but as the *lengths* of certain straight lines drawn in relation to this arc.



Let  $OA$  and  $OB$  be two radii of a circle at right angles, and let  $P$  be any point on the circumference. Draw  $PM$  and  $PN$  perpendicular to  $OA$  and  $OB$  respectively, and let the tangents at  $A$  and  $B$  meet  $OP$  produced in  $T$  and  $t$  respectively.

The lines  $PM$ ,  $AT$ ,  $OT$ ,  $AM$  were named respectively the sine, tangent, secant, versed-sine of the arc  $AP$ , and  $PN$ ,  $Bt$ ,  $Ot$ ,  $BN$ , which are the sine, tangent, secant, versed-sine of the complementary arc  $BP$ , were named respectively the cosine, cotangent, cosecant, coversed-sine of the arc  $AP$ .

As thus defined each trigonometrical function of the *arc* is equal to the corresponding function of the *angle*, which it subtends at the centre of the circle, multiplied by the radius. Thus

$$\frac{AT}{OA} = \tan POA; \text{ that is, } AT = OA \times \tan POA;$$

and 
$$\frac{Ot}{OB} = \sec BOP = \operatorname{cosec} POA; \text{ that is, } Ot = OB \times \operatorname{cosec} POA.$$

The values of the functions of the arc therefore depended on the length of the radius of the circle as well as on the angle subtended by the arc at the centre of the circle, so that in Tables of the functions it was necessary to state the magnitude of the radius.

The names of the trigonometrical functions and the abbreviations for them now in use were introduced by different Mathematicians chiefly towards the end of the sixteenth and during the seventeenth century, but were not generally employed until their re-introduction by Euler. The development of the science of Trigonometry may be considered to date from the publication in 1748 of Euler's *Introductio in analysin Infinitorum*.

The reader will find some interesting information regarding the progress of Trigonometry in Ball's *Short History of Mathematics*.

## MISCELLANEOUS EXAMPLES. C.

1. Draw the boundary lines of the angles whose tangent is equal to  $-\frac{3}{4}$ , and find the cosine of these angles.

2. Shew that

$$\cos A (2 \sec A + \tan A) (\sec A - 2 \tan A) = 2 \cos A - 3 \tan A.$$

3. Given  $C=90^\circ$ ,  $b=10\cdot5$ ,  $c=21$ , solve the triangle.

4. If  $\sec A = -\frac{25}{7}$ , and  $A$  lies between  $180^\circ$  and  $270^\circ$ , find  $\cot A$ .

5. The latitude of Bombay is  $19^\circ \text{N.}$ : find its distance from the equator, taking the diameter of the earth to be 7920 miles.

6. From the top of a cliff 200 ft. high, the angles of depression of two boats due east of the observer are  $34^\circ 30'$  and  $18^\circ 40'$ : find their distance apart, given

$$\cot 34^\circ 30' = 1\cdot455, \quad \cot 18^\circ 40' = 2\cdot96.$$

7. If  $A$  lies between  $180^\circ$  and  $270^\circ$ , and  $3 \tan A = 4$ , find the value of  $2 \cot A - 5 \cos A + \sin A$ .

8. Find, correct to three decimal places, the radius of a circle in which an arc 15 inches long subtends at the centre an angle of  $71^\circ 36' 3\cdot6''$ .

9. Shew that

$$\frac{\tan^3 \theta}{1 + \tan^2 \theta} + \frac{\cot^3 \theta}{1 + \cot^2 \theta} = \frac{1 - 2 \sin^2 \theta \cos^2 \theta}{\sin \theta \cos \theta}.$$

10. The angle of elevation of the top of a tower is  $68^\circ 11'$ , and a flagstaff 24 ft. high on the summit of the tower subtends an angle of  $2^\circ 10'$  at the observer's eye. Find the height of the tower, given


$$\tan 70^\circ 21' = 2\cdot8, \quad \cot 68^\circ 11' = \cdot4.$$

## CHAPTER X.

### CIRCULAR FUNCTIONS OF CERTAIN ALLIED ANGLES.

#### 92. Circular Functions of $180^\circ - A$ .

Take any straight line  $XOX'$ , and let a radius vector starting from  $O$  revolve until it has traced the angle  $A$ , taking up the position  $OP$ .



Again, let the radius vector starting from  $O$  revolve through  $180^\circ$  into the position  $OX'$  and then *back again* through an angle  $A$  taking up the final position  $OP'$ . Thus  $XOP'$  is the angle  $180^\circ - A$ .

From  $P$  and  $P'$  draw  $PM$  and  $P'M'$  perpendicular to  $XX'$ ; then by Euc. I. 26 the triangles  $OPM$  and  $OP'M'$  are geometrically equal.

By definition,

$$\sin(180^\circ - A) = \frac{M'P'}{OP'};$$

but  $M'P'$  is equal to  $MP$  in magnitude and is of the same sign;

$$\therefore \sin(180^\circ - A) = \frac{MP}{OP} = \sin A.$$

$$\text{Again,} \quad \cos(180^\circ - A) = \frac{OM'}{OP'};$$

and  $OM'$  is equal to  $OM$  in magnitude, but is of opposite sign;

$$\therefore \cos(180^\circ - A) = \frac{-OM}{OP} = -\frac{OM}{OP} = -\cos A.$$

$$\text{Also} \quad \tan(180^\circ - A) = \frac{M'P'}{OM'} = \frac{MP}{-OM} = -\frac{MP}{OM} = -\tan A.$$

93. In the last article, for the sake of simplicity we have supposed the angle  $A$  to be less than a right angle, but all the formulæ of this chapter may be shewn to be true for angles of any magnitude. A general proof of one case is given in Art. 102, and the same method may be applied to all the other cases.

94. If the angles are expressed in radian measure, the formulæ of Art. 92 become

$$\begin{aligned}\sin(\pi - \theta) &= \sin \theta, \\ \cos(\pi - \theta) &= -\cos \theta, \\ \tan(\pi - \theta) &= -\tan \theta.\end{aligned}$$

*Example 1.* Find the sine and cosine of  $120^\circ$ .

$$\begin{aligned}\sin 120^\circ &= \sin(180^\circ - 60^\circ) = \sin 60^\circ = \frac{\sqrt{3}}{2}. \\ \cos 120^\circ &= \cos(180^\circ - 60^\circ) = -\cos 60^\circ = -\frac{1}{2}.\end{aligned}$$

*Example 2.* Find the cosine and cotangent of  $\frac{5\pi}{6}$ .

$$\begin{aligned}\cos \frac{5\pi}{6} &= \cos\left(\pi - \frac{\pi}{6}\right) = -\cos \frac{\pi}{6} = -\frac{\sqrt{3}}{2}. \\ \cot \frac{5\pi}{6} &= \cot\left(\pi - \frac{\pi}{6}\right) = -\cot \frac{\pi}{6} = -\sqrt{3}.\end{aligned}$$

95. DEFINITION. When the sum of two angles is equal to two right angles each is said to be the **supplement** of the other and the angles are said to be **supplementary**. Thus if  $A$  is any angle its supplement is  $180^\circ - A$ .

96. The results of Art. 92 are so important in a later part of the subject that it is desirable to emphasize them. We therefore repeat them in a verbal form :


*the sines of supplementary angles are equal in magnitude and are of the same sign;*

*the cosines of supplementary angles are equal in magnitude but are of opposite sign;*

*the tangents of supplementary angles are equal in magnitude but are of opposite sign.*

### 97. Circular Functions of $180^\circ + A$ .

Take any straight line  $XOX'$  and let a radius vector starting from  $O$  revolve until it has traced the angle  $A$ , taking up the position  $OP$ .



Again, let the radius vector starting from  $O$  revolve through  $180^\circ$  into the position  $OX'$ , and then further through an angle  $A$ , taking up the final position  $OP'$ . Thus  $XOP'$  is the angle  $180^\circ + A$ .

From  $P$  and  $P'$  draw  $PM$  and  $P'M'$  perpendicular to  $XX'$ ; then  $OP$  and  $OP'$  are in the same straight line, and by Euc. I. 26 the triangles  $OPM$  and  $OP'M'$  are geometrically equal.

By definition,

$$\sin(180^\circ + A) = \frac{M'P'}{OP'};$$

and  $M'P'$  is equal to  $MP$  in magnitude but is of opposite sign;

$$\therefore \sin(180^\circ + A) = \frac{-MP}{OP} = -\frac{MP}{OP} = -\sin A.$$

$$\text{Again,} \quad \cos(180^\circ + A) = \frac{OM'}{OP'};$$

and  $OM'$  is equal to  $OM$  in magnitude but is of opposite sign;

$$\therefore \cos(180^\circ + A) = \frac{-OM}{OP} = -\frac{OM}{OP} = -\cos A.$$

$$\text{Also} \quad \tan(180^\circ + A) = \frac{M'P'}{OM'} = \frac{-MP}{-OM} = \frac{MP}{OM} = \tan A.$$

Expressed in radian measure, the above formulæ are written  
 $\sin(\pi + \theta) = -\sin \theta, \quad \cos(\pi + \theta) = -\cos \theta, \quad \tan(\pi + \theta) = \tan \theta.$

In these results we may draw especial attention to the fact that an angle may be increased or diminished by two right angles as often as we please without altering the value of the tangent.


*Example.* Find the value of  $\cot 210^\circ$ .

$$\cot 210^\circ = \cot(180^\circ + 30^\circ) = \cot 30^\circ = \sqrt{3}.$$

# 98. Circular Functions of $90^\circ + A$ .

Take any straight line  $XOX'$ , and let a radius vector starting from  $OX$  revolve until it has traced the angle  $A$ , taking up the position  $OP$ .

Again, let the radius vector starting from  $OX$  revolve through  $90^\circ$  into the position  $OY$ , and then further through  $X'$  an angle  $A$ , taking up the final position  $OP'$ . Thus  $X'OP'$  is the angle  $90^\circ + A$ .



From  $P$  and  $P'$  draw  $PM$  and  $P'M'$  perpendicular to  $XX'$ ; then  $\angle M'P'O = \angle P'OY = A = \angle POM$ .

By Euc. I. 26, the triangles  $OPM$  and  $OP'M'$  are geometrically equal; hence

$M'P'$  is equal to  $OM$  in magnitude and is of the same sign, and  $OM'$  is equal to  $MP$  in magnitude but is of opposite sign.

By definition,

$$\sin(90^\circ + A) = \frac{M'P'}{OP'} = \frac{OM}{OP} = \cos A;$$

$$\cos(90^\circ + A) = \frac{OM'}{OP'} = -\frac{MP}{OP} = -\sin A;$$

$$\tan(90^\circ + A) = \frac{M'P'}{OM'} = \frac{OM}{-MP} = -\cot A.$$

Expressed in radian measure the above formulæ become

$$\sin\left(\frac{\pi}{2} + \theta\right) = \cos \theta, \quad \cos\left(\frac{\pi}{2} + \theta\right) = -\sin \theta, \quad \tan\left(\frac{\pi}{2} + \theta\right) = -\cot \theta.$$

*Example 1.* Find the value of  $\sin 120^\circ$ .

$$\sin 120^\circ = \sin(90^\circ + 30^\circ) = \cos 30^\circ = \frac{\sqrt{3}}{2}.$$



*Example 2.* Find the values of  $\tan (270^\circ + A)$  and  $\cos \left( \frac{3\pi}{2} + \theta \right)$ .


$$\tan (270^\circ + A) = \tan (180^\circ + \overline{90^\circ + A}) = \tan (90^\circ + A) = -\cot A;$$

$$\cos \left( \frac{3\pi}{2} + \theta \right) = \cos \left( \pi + \overline{\frac{\pi}{2} + \theta} \right) = -\cos \left( \frac{\pi}{2} + \theta \right) = \sin \theta.$$

## 99. Circular Functions of $-A$ .

Take any straight line  $OX$  and let a radius vector starting from  $O$  revolve until it has traced the angle  $A$ , taking up the position  $OP$ .

Again, let the radius vector starting from  $O$  revolve in the *opposite* direction until it has traced the angle  $A$ , taking up the position  $OP'$ . Join  $PP'$ ; then  $MP'$  is equal to  $MP$  in magnitude, and the angles at  $M$  are right angles. [Euc. I. 4.]



By definition,

$$\sin(-A) = \frac{MP'}{OP'} = -\frac{MP}{OP} = -\sin A;$$

$$\cos(-A) = \frac{OM}{OP'} = \frac{OM}{OP} = \cos A;$$

$$\tan(-A) = \frac{MP'}{OM} = -\frac{MP}{OM} = -\tan A.$$

It is especially worthy of notice that *we may change the sign of an angle without altering the value of its cosine.*

*Example.* Find the values of

$$\operatorname{cosec}(-210^\circ) \text{ and } \cos(A - 270^\circ).$$

$$\operatorname{cosec}(-210^\circ) = -\operatorname{cosec} 210^\circ = -\operatorname{cosec}(180^\circ + 30^\circ) = \operatorname{cosec} 30^\circ = 2.$$

$$\begin{aligned} \cos(A - 270^\circ) &= \cos(270^\circ - A) = \cos(180^\circ + \overline{90^\circ - A}) \\ &= -\cos(90^\circ - A) = -\sin A. \end{aligned}$$



100. If  $f(A)$  denotes a function of  $A$  which is unaltered in magnitude and sign when  $-A$  is written for  $A$ , then  $f(A)$  is said to be an **even function** of  $A$ . In this case  $f(-A) = f(A)$ .

If when  $-A$  is written for  $A$ , the sign of  $f(A)$  is changed while the magnitude remains unaltered,  $f(A)$  is said to be an **odd function** of  $A$ , and in this case  $f(-A) = -f(A)$ .

From the last article it will be seen that

$\cos A$  and  $\sec A$  are even functions of  $A$ ,  
 $\sin A$ ,  $\operatorname{cosec} A$ ,  $\tan A$ ,  $\cot A$  are odd functions of  $A$ .

### EXAMPLES. X. a.

Find the numerical value of

- |  |   |   |
|--|---|---|
| 1. $\cos 135^\circ$ .                                    | 2. $\sin 150^\circ$ .                     | 3. $\tan 240^\circ$ .                     |
| 4. $\operatorname{cosec} 225^\circ$ .                    | 5. $\sin (-120^\circ)$ .                  | 6. $\cot (-135^\circ)$ .                  |
| 7. $\cot 315^\circ$ .                                    | 8. $\cos (-240^\circ)$ .                  | 9. $\sec (-300^\circ)$ .                  |
| 10. $\tan \frac{3\pi}{4}$ .                              | 11. $\sin \frac{4\pi}{3}$ .               | 12. $\sec \frac{2\pi}{3}$ .               |
| 13. $\operatorname{cosec} \left(-\frac{\pi}{6}\right)$ . | 14. $\cos \left(-\frac{3\pi}{4}\right)$ . | 15. $\cot \left(-\frac{5\pi}{6}\right)$ . |

Express as functions of  $A$ :

- |                              |                              |                             |
|------------------------------|------------------------------|-----------------------------|
| 16. $\cos (270^\circ + A)$ . | 17. $\cot (270^\circ - A)$ . | 18. $\sin (A - 90^\circ)$ . |
| 19. $\sec (A - 180^\circ)$ . | 20. $\sin (270^\circ - A)$ . | 21. $\cot (A - 90^\circ)$ . |

Express as functions of  $\theta$ :


- |  |                             |   |
|--|-----------------------------|---|
| 22. $\sin \left(\theta - \frac{\pi}{2}\right)$ . | 23. $\tan (\theta - \pi)$ . | 24. $\sec \left(\frac{3\pi}{2} - \theta\right)$ . |
|--|-----------------------------|---|

Express in the simplest form:

- |  |
|--|
| 25. $\tan (180^\circ + A) \sin (90^\circ + A) \sec (90^\circ - A)$ .                         |
| 26. $\cos (90^\circ + A) + \sin (180^\circ - A) - \sin (180^\circ + A) - \sin (-A)$ .        |
| 27. $\sec (180^\circ + A) \sec (180^\circ - A) + \cot (90^\circ + A) \tan (180^\circ + A)$ . |

101. In Art. 38 we have established the relations which subsist between the trigonometrical ratios of  $90^\circ - A$  and those of  $A$ , when  $A$  is an acute angle. We shall now give a general proof which is applicable whatever be the magnitude of  $A$ .

### 102. Circular Functions of $90^\circ - A$ for any value of $A$ .



Let a radius vector starting from  $OX$  revolve until it has traced the angle  $A$ , taking up the position  $OP$  in each of the two figures.

Again, let the radius vector starting from  $OX$  revolve through  $90^\circ$  into the position  $OY$  and then *back again* through an angle  $A$ , taking up the final position  $OP'$  in each of the two figures.

Draw  $PM$  and  $P'M'$  perpendicular to  $XX'$ ; then whatever be the value of  $A$ , it will be found that  $\angle OPM' = \angle POM$ , so that the triangles  $OMP$  and  $OM'P'$  are geometrically equal, having  $MP$  equal to  $OM'$ , and  $OM$  equal to  $M'P'$ , *in magnitude*.

When  $P$  is above  $XX'$ ,  $P'$  is to the right of  $YY'$ ,  
and when  $P$  is below  $XX'$ ,  $P'$  is to the left of  $YY'$ .

When  $P'$  is above  $XX'$ ,  $P$  is to the right of  $YY'$ ,  
and when  $P'$  is below  $XX'$ ,  $P$  is to the left of  $YY'$ .

Hence  $MP$  is equal to  $OM'$  in magnitude and is always of the same sign as  $OM'$ ;

and  $M'P'$  is equal to  $OM$  in magnitude and is always of the same sign as  $OM$ .

By definition,

$$\sin (90^{\circ} - A) = \frac{M'P'}{OP'} = \frac{OM}{OP} = \cos A ;$$

$$\cos (90^{\circ} - A) = \frac{OM'}{OP'} = \frac{MP}{OP} = \sin A ;$$

$$\tan (90^{\circ} - A) = \frac{M'P'}{OM'} = \frac{OM}{MP} = \cot A .$$

A general method similar to the above may be applied to all the other cases of this chapter.

### 103. Circular Functions of $n \cdot 360^{\circ} + A$ .

If  $n$  is any integer,  $n \cdot 360^{\circ}$  represents  $n$  complete revolutions of the radius vector, and therefore the boundary line of the angle  $n \cdot 360^{\circ} + A$  is coincident with that of  $A$ . The value of each function of the angle  $n \cdot 360^{\circ} + A$  is thus the same as the value of the corresponding function of  $A$  both in magnitude and in sign.

104. Since the functions of all coterminal angles are equal, there is a *recurrence* of the values of the functions each time the boundary line completes its revolution and comes round into its original position. This is otherwise expressed by saying that *the circular functions are periodic*, and  $360^{\circ}$  is said to be *the amplitude of the period*.

In radian measure, the amplitude of the period is  $2\pi$ .

NOTE. In the case of the tangent and cotangent the amplitude of the period is half that of the other circular functions, being  $180^{\circ}$  or  $\pi$  radians. [Art. 97.]

### 105. Circular Functions of $n \cdot 360^{\circ} - A$ .

If  $n$  is any integer, the boundary line of  $n \cdot 360^{\circ} - A$  is coincident with that of  $-A$ . The value of each function of  $n \cdot 360^{\circ} - A$  is thus the same as the value of the corresponding function of  $-A$  both in magnitude and in sign ; hence

$$\sin (n \cdot 360^{\circ} - A) = \sin (-A) = -\sin A ;$$

$$\cos (n \cdot 360^{\circ} - A) = \cos (-A) = \cos A ;$$

$$\tan (n \cdot 360^{\circ} - A) = \tan (-A) = -\tan A .$$

**106.** We can always express the functions of any angle in terms of the functions of some positive acute angle. In the arrangement of the work it is advisable to follow a uniform plan.

(1) If the angle is negative, use the relations connecting the functions of  $-A$  and  $A$ . [Art. 99.]

$$\text{Thus} \quad \sin(-30^\circ) = -\sin 30^\circ = -\frac{1}{2};$$

$$\cos(-845^\circ) = \cos 845^\circ.$$

(2) If the angle is greater than  $360^\circ$ , by taking off multiples of  $360^\circ$  the angle may be replaced by a coterminal angle less than  $360^\circ$ . [Art. 103.]

$$\text{Thus} \quad \tan 735^\circ = \tan(2 \times 360^\circ + 15^\circ) = \tan 15^\circ.$$

(3) If the angle is still greater than  $180^\circ$ , use the relations connecting the functions of  $180^\circ + A$  and  $A$ . [Art. 97.]

$$\begin{aligned} \text{Thus} \quad \cot 585^\circ &= \cot(360^\circ + 225^\circ) = \cot 225^\circ \\ &= \cot(180^\circ + 45^\circ) = \cot 45^\circ = 1. \end{aligned}$$

(4) If the angle is still greater than  $90^\circ$ , use the relations connecting the functions of  $180^\circ - A$  and  $A$ . [Art. 92.]

$$\begin{aligned} \text{Thus} \quad \cos 675^\circ &= \cos(360^\circ + 315^\circ) = \cos 315^\circ \\ &= \cos(180^\circ + 135^\circ) = -\cos 135^\circ \\ &= -\cos(180^\circ - 45^\circ) = \cos 45^\circ = \frac{1}{\sqrt{2}}. \end{aligned}$$

*Example.* Express  $\sin(-1190^\circ)$ ,  $\tan 1000^\circ$ ,  $\cos(-3860^\circ)$  as functions of positive acute angles.

$$\begin{aligned} \sin(-1190^\circ) &= -\sin 1190^\circ = -\sin(3 \times 360^\circ + 110^\circ) = -\sin 110^\circ \\ &= -\sin(180^\circ - 70^\circ) = -\sin 70^\circ. \end{aligned}$$

$$\begin{aligned} \tan 1000^\circ &= \tan(2 \times 360^\circ + 280^\circ) = \tan 280^\circ \\ &= \tan(180^\circ + 100^\circ) = \tan 100^\circ \\ &= \tan(180^\circ - 80^\circ) = -\tan 80^\circ. \end{aligned}$$

$$\begin{aligned} \cos(-3860^\circ) &= \cos 3860^\circ = \cos(10 \times 360^\circ + 260^\circ) = \cos 260^\circ \\ &= \cos(180^\circ + 80^\circ) = -\cos 80^\circ. \end{aligned}$$

107. From the investigations of this chapter we see that the number of angles which have the same circular function is unlimited. Thus if  $\tan \theta = 1$ ,  $\theta$  may be any one of the angles coterminial with  $45^\circ$  or  $225^\circ$ .

*Example.* Draw the boundary lines of  $A$  when  $\sin A = \frac{\sqrt{3}}{2}$ , and write down all the angles numerically less than  $360^\circ$  which satisfy the equation.


Since  $\sin 60^\circ = \frac{\sqrt{3}}{2}$ , if we draw  $OP$  making  $\angle XOP = 60^\circ$ , then  $OP$  is one position of the boundary line.

Again,  $\sin 60^\circ = \sin (180^\circ - 60^\circ) = \sin 120^\circ$ , so that another position of the boundary line will be found by making  $\angle XOP' = 120^\circ$ .

There will be no position of the boundary line in the third or fourth quadrant, since in these quadrants the sine is negative.

Thus in one complete revolution  $OP$  and  $OP'$  are the only two positions of the boundary line of the angle  $A$ .

Hence the positive angles are  $60^\circ$  and  $120^\circ$ ;  
and the negative angles are  $-(360^\circ - 120^\circ)$  and  $-(360^\circ - 60^\circ)$ ; that is,  $-240^\circ$  and  $-300^\circ$ .



### EXAMPLES. X. b.

Find the numerical value of

- |  |  |  |
|--|--|--|
| 1. $\cos 480^\circ$ .                    | 2. $\sin 960^\circ$ .                                      | 3. $\cos (-780^\circ)$ .                 |
| 4. $\sin (-870^\circ)$ .                 | 5. $\sec 900^\circ$ .                                      | 6. $\tan (-855^\circ)$ .                 |
| 7. $\operatorname{cosec} (-660^\circ)$ . | 8. $\cot 840^\circ$ .                                      | 9. $\operatorname{cosec} (-765^\circ)$ . |
| 10. $\cos 4005^\circ$ .                  | 11. $\cot 990^\circ$ .                                     | 12. $\sin 3015^\circ$ .                  |
| 13. $\sec 2745^\circ$ .                  | 14. $\cos 2400^\circ$ .                                    | 15. $\sec (-5895^\circ)$ .               |
| 16. $\sin \frac{15\pi}{4}$ .             | 17. $\cot \frac{23\pi}{4}$ .                               | 18. $\sec \frac{7\pi}{3}$ .              |
| 19. $\cot \frac{16\pi}{3}$ .             | 20. $\sec \left( \frac{3\pi}{2} + \frac{\pi}{3} \right)$ . |  |

Find all the angles numerically less than  $360^\circ$  which satisfy the equations :

$$21. \quad \cos \theta = \frac{\sqrt{3}}{2}.$$

$$22. \quad \sin \theta = -\frac{1}{2}.$$

$$23. \quad \tan \theta = -\sqrt{3}.$$

$$24. \quad \cot \theta = -1.$$

If  $A$  is less than  $90^\circ$ , prove geometrically

$$25. \quad \sec (A - 180^\circ) = -\sec A.$$

$$26. \quad \tan (270^\circ + A) = -\cot A.$$

$$27. \quad \cos (A - 90^\circ) = \sin A.$$

28. Prove that

$$\tan A + \tan (180^\circ - A) + \cot (90^\circ + A) = \tan (360^\circ - A).$$

29. Shew that

$$\frac{\sin (180^\circ - A)}{\tan (180^\circ + A)} \cdot \frac{\cot (90^\circ - A)}{\tan (90^\circ + A)} \cdot \frac{\cos (360^\circ - A)}{\sin (-A)} = \sin A.$$

Express in the simplest form

$$30. \quad \frac{\sin (-A)}{\sin (180^\circ + A)} - \frac{\tan (90^\circ + A)}{\cot A} + \frac{\cos A}{\sin (90^\circ + A)}.$$

$$31. \quad \frac{\operatorname{cosec} (180^\circ - A)}{\sec (180^\circ + A)} \cdot \frac{\cos (-A)}{\cos (90^\circ + A)}.$$

$$32. \quad \frac{\cos (90^\circ + A) \sec (-A) \tan (180^\circ - A)}{\sec (360^\circ + A) \sin (180^\circ + A) \cot (90^\circ - A)}.$$

$$33. \quad \text{Prove that } \sin \left( \frac{\pi}{2} + \theta \right) \cos (\pi - \theta) \cot \left( \frac{3\pi}{2} + \theta \right) \\ = \sin \left( \frac{\pi}{2} - \theta \right) \sin \left( \frac{3\pi}{2} - \theta \right) \cot \left( \frac{\pi}{2} + \theta \right).$$

$$34. \quad \text{When } a = \frac{11\pi}{4}, \text{ find the numerical value of}$$

$$\sin^2 a - \cos^2 a + 2 \tan a - \sec^2 a.$$



## CHAPTER XI.

### FUNCTIONS OF COMPOUND ANGLES.

108. WHEN an angle is made up by the algebraical sum of two or more angles it is called a **compound angle**; thus  $A + B$ ,  $A - B$ , and  $A + B - C$  are compound angles.

109. Hitherto we have only discussed the properties of the functions of single angles, such as  $A$ ,  $B$ ,  $\alpha$ ,  $\theta$ . In the present chapter we shall prove some fundamental properties relating to the functions of compound angles. We shall begin by finding expressions for the sine, cosine, and tangent of  $A + B$  and  $A - B$  in terms of the functions of  $A$  and  $B$ .

It may be useful to caution the student against the prevalent mistake of supposing that a function of  $A + B$  is equal to the sum of the corresponding functions of  $A$  and  $B$ , and a function of  $A - B$  to the difference of the corresponding functions.

Thus  $\sin(A + B)$  is not equal to  $\sin A + \sin B$ ,  
and  $\cos(A - B)$  is not equal to  $\cos A - \cos B$ .

A numerical instance will illustrate this.

Thus if  $A = 60^\circ$ ,  $B = 30^\circ$ , then  $A + B = 90^\circ$ ,  
so that  $\cos(A + B) = \cos 90^\circ = 0$ ;  
but  $\cos A + \cos B = \cos 60^\circ + \cos 30^\circ = \frac{1}{2} + \frac{\sqrt{3}}{2}$ .

Hence  $\cos(A + B)$  is not equal to  $\cos A + \cos B$ .

In like manner,  $\sin(A + A)$  is not equal to  $\sin A + \sin A$ ;  
that is,  $\sin 2A$  is not equal to  $2 \sin A$ .  
Similarly  $\tan 3A$  is not equal to  $3 \tan A$ .




110. To prove the formulæ

$$\sin(A+B) = \sin A \cos B + \cos A \sin B,$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B.$$

Let  $\angle LOM = A$ , and  $\angle MON = B$ ; then  $\angle LON = A + B$ .



In  $ON$ , the boundary line of the compound angle  $A + B$ , take any point  $P$ , and draw  $PQ$  and  $PR$  perpendicular to  $OL$  and  $OM$  respectively; also draw  $RS$  and  $RT$  perpendicular to  $OL$  and  $PQ$  respectively.

By definition,

$$\begin{aligned} \sin(A+B) &= \frac{PQ}{OP} = \frac{RS+PT}{OP} = \frac{RS}{OP} + \frac{PT}{OP} \\ &= \frac{RS}{OR} \cdot \frac{OR}{OP} + \frac{PT}{PR} \cdot \frac{PR}{OP} \\ &= \sin A \cdot \cos B + \cos A \sin B. \end{aligned}$$

But  $\angle TPR = 90^\circ - \angle TRP = \angle TRO = \angle ROS = A$ ;

$$\therefore \sin(A+B) = \sin A \cos B + \cos A \sin B.$$


$$\begin{aligned} \text{Also } \cos(A+B) &= \frac{OQ}{OP} = \frac{OS-TR}{OP} = \frac{OS}{OP} - \frac{TR}{OP} \\ &= \frac{OS}{OR} \cdot \frac{OR}{OP} - \frac{TR}{PR} \cdot \frac{PR}{OP} \\ &= \cos A \cdot \cos B - \sin A \sin B \\ &= \cos A \cos B - \sin A \sin B. \end{aligned}$$

111. To prove the formulæ

$$\sin (A-B)=\sin A \cos B-\cos A \sin B,$$

$$\cos (A-B)=\cos A \cos B+\sin A \sin B.$$

Let  $\angle LOM=A$ , and  $\angle MON=B$ ; then  $\angle LON=A-B$ .



In  $ON$ , the boundary line of the compound angle  $A-B$ , take any point  $P$ , and draw  $PQ$  and  $PR$  perpendicular to  $OL$  and  $OM$  respectively; also draw  $RS$  and  $RT$  perpendicular to  $OL$  and  $QP$  respectively.

By definition,

$$\begin{aligned}\sin (A-B) &= \frac{PQ}{OP} = \frac{RS-PT}{OP} = \frac{RS}{OP} - \frac{PT}{OP} \\ &= \frac{RS}{OR} \cdot \frac{OR}{OP} - \frac{PT}{PR} \cdot \frac{PR}{OP} \\ &= \sin A \cdot \cos B - \cos TPR \cdot \sin B.\end{aligned}$$

But  $\angle TPR = 90^\circ - \angle TRP = \angle MRT = \angle MOL = A$ ;

$$\therefore \sin (A-B) = \sin A \cos B - \cos A \sin B.$$

$$\begin{aligned}\text{Also } \cos (A-B) &= \frac{OQ}{OP} = \frac{OS+RT}{OP} = \frac{OS}{OP} + \frac{RT}{OP} \\ &= \frac{OS}{OR} \cdot \frac{OR}{OP} + \frac{RT}{RP} \cdot \frac{RP}{OP} \\ &= \cos A \cdot \cos B + \sin TPR \cdot \sin B \\ &= \cos A \cos B + \sin A \sin B.\end{aligned}$$

112. The *expansions* of  $\sin(A \pm B)$  and  $\cos(A \pm B)$  are frequently called the "Addition Formulæ." We shall sometimes refer to them as the " $A + B$ " and " $A - B$ " formulæ.

113. In the foregoing geometrical proofs we have supposed that the angles  $A$ ,  $B$ ,  $A + B$  are all less than a right angle, and that  $A - B$  is positive. If the angles are not so restricted some modification of the figures will be required. It is however unnecessary to consider these cases in detail, as in Chap. XXII. we shall shew by the Method of Projections that the Addition Formulæ hold universally. In the meantime the student may assume that they are always true.

*Example 1.* Find the value of  $\cos 75^\circ$ .

$$\begin{aligned}\cos 75^\circ &= \cos(45^\circ + 30^\circ) = \cos 45^\circ \cos 30^\circ - \sin 45^\circ \sin 30^\circ \\ &= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \cdot \frac{1}{2} = \frac{\sqrt{3}-1}{2\sqrt{2}}.\end{aligned}$$

*Example 2.* If  $\sin A = \frac{4}{5}$  and  $\sin B = \frac{5}{13}$ , find  $\sin(A - B)$ .

$$\sin(A - B) = \sin A \cos B - \cos A \sin B.$$

But  $\cos A = \sqrt{1 - \sin^2 A} = \sqrt{1 - \frac{16}{25}} = \frac{3}{5};$

and  $\cos B = \sqrt{1 - \sin^2 B} = \sqrt{1 - \frac{25}{169}} = \frac{12}{13};$

$$\therefore \sin(A - B) = \frac{4}{5} \cdot \frac{12}{13} - \frac{3}{5} \cdot \frac{5}{13} = \frac{33}{65}.$$

NOTE. Strictly speaking  $\cos A = \pm \frac{3}{5}$  and  $\cos B = \pm \frac{12}{13}$ , so that  $\sin(A - B)$  has *four* values. We shall however suppose that in similar cases only the positive value of the square root is taken.

114. To prove that  $\sin(A + B) \sin(A - B) = \sin^2 A - \sin^2 B$ .

The first side

$$\begin{aligned}&= (\sin A \cos B + \cos A \sin B)(\sin A \cos B - \cos A \sin B) \\ &= \sin^2 A \cos^2 B - \cos^2 A \sin^2 B \\ &= \sin^2 A (1 - \sin^2 B) - (1 - \sin^2 A) \sin^2 B \\ &= \sin^2 A - \sin^2 B.\end{aligned}$$

# EXAMPLES. XI. a.

[The examples printed in more prominent type are important, and should be regarded as standard formulæ.]

Prove that

1.  $\sin (A + 45^\circ) = \frac{1}{\sqrt{2}} (\sin A + \cos A).$
2.  $\cos (A + 45^\circ) = \frac{1}{\sqrt{2}} (\cos A - \sin A).$
3.  $2 \sin (30^\circ - A) = \cos A - \sqrt{3} \sin A.$
4. If  $\cos A = \frac{4}{5}$ ,  $\cos B = \frac{3}{5}$ , find  $\sin (A + B)$  and  $\cos (A - B).$
5. If  $\sin A = \frac{3}{5}$ ,  $\cos B = \frac{12}{13}$ , find  $\cos (A + B)$  and  $\sin (A - B).$
6. If  $\sec A = \frac{17}{8}$ ,  $\operatorname{cosec} B = \frac{5}{4}$ , find  $\sec (A + B).$

Prove that

7.  $\sin 75^\circ = \cos 15^\circ = \frac{\sqrt{3}+1}{2\sqrt{2}}.$
8.  $\sin 15^\circ = \cos 75^\circ = \frac{\sqrt{3}-1}{2\sqrt{2}}.$
9.  $\frac{\sin (a + \beta)}{\cos a \cos \beta} = \tan a + \tan \beta.$
10.  $\frac{\sin (a - \beta)}{\sin a \sin \beta} = \cot \beta - \cot a.$
11.  $\frac{\cos (a - \beta)}{\cos a \sin \beta} = \cot \beta + \tan a.$
12.  $\cos (A + B) \cos (A - B) = \cos^2 A - \sin^2 B.$
13.  $\sin (A + B) \sin (A - B) = \cos^2 B - \cos^2 A.$
14.  $\cos (45^\circ - A) - \sin (45^\circ + A) = 0.$
15.  $\cos (45^\circ + A) + \sin (A - 45^\circ) = 0.$
16.  $\cos (A - B) - \sin (A + B) = (\cos A - \sin A)(\cos B - \sin B).$
17.  $\cos (A + B) + \sin (A - B) = (\cos A + \sin A)(\cos B - \sin B).$

Prove the following identities :

$$18. \quad 2 \sin (A + 45^\circ) \sin (A - 45^\circ) = \sin^2 A - \cos^2 A.$$

$$19. \quad 2 \cos \left( \frac{\pi}{4} + a \right) \cos \left( \frac{\pi}{4} - a \right) = \cos^2 a - \sin^2 a.$$

$$20. \quad 2 \sin \left( \frac{\pi}{4} + a \right) \cos \left( \frac{\pi}{4} + \beta \right) = \cos (a + \beta) + \sin (a - \beta).$$

$$21. \quad \frac{\sin (\beta - \gamma)}{\cos \beta \cos \gamma} + \frac{\sin (\gamma - a)}{\cos \gamma \cos a} + \frac{\sin (a - \beta)}{\cos a \cos \beta} = 0.$$

115. To expand  $\tan (A + B)$  in terms of  $\tan A$  and  $\tan B$ .

$$\tan (A + B) = \frac{\sin (A + B)}{\cos (A + B)} = \frac{\sin A \cos B + \cos A \sin B}{\cos A \cos B - \sin A \sin B}.$$

To express this fraction in terms of *tangents*, divide each term of numerator and denominator by  $\cos A \cos B$ ;

$$\therefore \tan (A + B) = \frac{\frac{\sin A}{\cos A} + \frac{\sin B}{\cos B}}{1 - \frac{\sin A}{\cos A} \cdot \frac{\sin B}{\cos B}};$$

$$\text{that is,} \quad \tan (A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}.$$

A geometrical proof of this result is given in Chap. XXII.

Similarly, we may prove that

$$\tan (A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}.$$

*Example.* Find the value of  $\tan 75^\circ$ .

$$\tan 75^\circ = \tan (45^\circ + 30^\circ) = \frac{\tan 45^\circ + \tan 30^\circ}{1 - \tan 45^\circ \tan 30^\circ}$$

$$= \frac{1 + \frac{1}{\sqrt{3}}}{1 - \frac{1}{\sqrt{3}}} = \frac{\sqrt{3} + 1}{\sqrt{3} - 1}$$

$$= \frac{(\sqrt{3} + 1)(\sqrt{3} + 1)}{3 - 1} = \frac{4 + 2\sqrt{3}}{2}$$

$$= 2 + \sqrt{3}.$$

116. To expand  $\cot(A+B)$  in terms of  $\cot A$  and  $\cot B$ .

$$\cot(A+B) = \frac{\cos(A+B)}{\sin(A+B)} = \frac{\cos A \cos B - \sin A \sin B}{\sin A \cos B + \cos A \sin B}.$$

To express this fraction in terms of *cotangents*, divide each term of numerator and denominator by  $\sin A \sin B$ ;

$$\therefore \cot(A+B) = \frac{\frac{\cos A \cos B}{\sin A \sin B} - 1}{\frac{\cos B}{\sin B} + \frac{\cos A}{\sin A}} = \frac{\cot A \cot B - 1}{\cot B + \cot A}.$$

Similarly, we may prove that

$$\cot(A-B) = \frac{\cot A \cot B + 1}{\cot B - \cot A}.$$

117. To find the expansion of  $\sin(A+B+C)$ .

$$\begin{aligned} \sin(A+B+C) &= \sin\{(A+B)+C\} \\ &= \sin(A+B) \cos C + \cos(A+B) \sin C \\ &= (\sin A \cos B + \cos A \sin B) \cos C \\ &\quad + (\cos A \cos B - \sin A \sin B) \sin C \\ &= \sin A \cos B \cos C + \cos A \sin B \cos C \\ &\quad + \cos A \cos B \sin C - \sin A \sin B \sin C. \end{aligned}$$

118. To find the expansion of  $\tan(A+B+C)$ .

$$\begin{aligned} \tan(A+B+C) &= \tan\{(A+B)+C\} = \frac{\tan(A+B) + \tan C}{1 - \tan(A+B) \tan C} \\ &= \frac{\frac{\tan A + \tan B}{1 - \tan A \tan B} + \tan C}{1 - \frac{\tan A + \tan B}{1 - \tan A \tan B} \cdot \tan C} \\ &= \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A}. \end{aligned}$$

COR. If  $A+B+C=180^\circ$ , then  $\tan(A+B+C)=0$ ; hence the numerator of the above expression must be zero.

$$\therefore \tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

**EXAMPLES. XI. b.**

[The examples printed in more prominent type are important, and should be regarded as standard formulæ.]

1. Find  $\tan(A+B)$  when  $\tan A = \frac{1}{2}$ ,  $\tan B = \frac{1}{3}$ .
2. If  $\tan A = \frac{4}{3}$ , and  $B = 45^\circ$ , find  $\tan(A-B)$ .
3. If  $\cot A = \frac{5}{7}$ ,  $\cot B = \frac{7}{5}$ , find  $\cot(A+B)$  and  $\tan(A-B)$ .
4. If  $\cot A = \frac{11}{2}$ ,  $\tan B = \frac{7}{24}$ , find  $\cot(A-B)$  and  $\tan(A+B)$ .
5.  $\tan(45^\circ + A) = \frac{1 + \tan A}{1 - \tan A}$ .
6.  $\tan(45^\circ - A) = \frac{1 - \tan A}{1 + \tan A}$ .
7.  $\cot\left(\frac{\pi}{4} - \theta\right) = \frac{\cot \theta + 1}{\cot \theta - 1}$ .
8.  $\cot\left(\frac{\pi}{4} + \theta\right) = \frac{\cot \theta - 1}{\cot \theta + 1}$ .
9.  $\tan 15^\circ = 2 - \sqrt{3}$ .
10.  $\cot 15^\circ = 2 + \sqrt{3}$ .
11. Find the expansions of  
 $\cos(A+B+C)$  and  $\sin(A-B+C)$ .
12. Express  $\tan(A-B-C)$  in terms of  $\tan A$ ,  $\tan B$ ,  $\tan C$ .
13. Express  $\cot(A+B+C)$  in terms of  $\cot A$ ,  $\cot B$ ,  $\cot C$ .

119. Beginners not unfrequently find a difficulty in the converse use of the  $A+B$  and  $A-B$  formulæ; that is, they fail to recognise when an expression is merely an expansion belonging to one of the standard forms.

*Example 1.* Simplify  $\cos(\alpha - \beta) \cos(\alpha + \beta) - \sin(\alpha - \beta) \sin(\alpha + \beta)$ .

This expression is the expansion of the cosine of the compound angle  $(\alpha + \beta) + (\alpha - \beta)$ , and is therefore equal to  $\cos\{(\alpha + \beta) + (\alpha - \beta)\}$ ; that is, to  $\cos 2\alpha$ .

*Example 2.* Shew that  $\frac{\tan A + \tan 2A}{1 - \tan A \tan 2A} = \tan 3A$ .

By Art. 115, the first side is the expansion of  $\tan(A+2A)$ , and is therefore equal to  $\tan 3A$ .



*Example 3.* Prove that  $\cot 2A + \tan A = \operatorname{cosec} 2A$ .

$$\begin{aligned} \text{The first side} &= \frac{\cos 2A}{\sin 2A} + \frac{\sin A}{\cos A} = \frac{\cos 2A \cos A + \sin 2A \sin A}{\sin 2A \cos A} \\ &= \frac{\cos (2A - A)}{\sin 2A \cos A} = \frac{\cos A}{\sin 2A \cos A} \\ &= \frac{1}{\sin 2A} = \operatorname{cosec} 2A. \end{aligned}$$

*Example 4.* Prove that

$$\cos 4\theta \cos \theta + \sin 4\theta \sin \theta = \cos 2\theta \cos \theta - \sin 2\theta \sin \theta.$$

$$\begin{aligned} \text{The first side} &= \cos (4\theta - \theta) = \cos 3\theta = \cos (2\theta + \theta) \\ &= \cos 2\theta \cos \theta - \sin 2\theta \sin \theta. \end{aligned}$$

### EXAMPLES. XI. c.

Prove the following identities :

1.  $\cos (A + B) \cos B + \sin (A + B) \sin B = \cos A.$
2.  $\sin 3A \cos A - \cos 3A \sin A = \sin 2A.$
3.  $\cos 2a \cos a + \sin 2a \sin a = \cos a.$
4.  $\cos (30^\circ + A) \cos (30^\circ - A) - \sin (30^\circ + A) \sin (30^\circ - A) = \frac{1}{2}.$
5.  $\sin (60^\circ - A) \cos (30^\circ + A) + \cos (60^\circ - A) \sin (30^\circ + A) = 1.$
6.  $\frac{\cos 2a}{\sec a} - \frac{\sin 2a}{\operatorname{cosec} a} = \cos 3a.$
7.  $\frac{\tan (a - \beta) + \tan \beta}{1 - \tan (a - \beta) \tan \beta} = \tan a.$
8.  $\frac{\cot (a + \beta) \cot a + 1}{\cot a - \cot (a + \beta)} = \cot \beta.$
9.  $\frac{\tan 4A - \tan 3A}{1 + \tan 4A \tan 3A} = \tan A.$
10.  $\cot \theta - \cot 2\theta = \operatorname{cosec} 2\theta.$
11.  $1 + \tan 2\theta \tan \theta = \sec 2\theta.$
12.  $1 + \cot 2\theta \cot \theta = \operatorname{cosec} 2\theta \cot \theta.$
13.  $\sin 2\theta \cos \theta + \cos 2\theta \sin \theta = \sin 4\theta \cos \theta - \cos 4\theta \sin \theta.$
14.  $\cos 4a \cos a - \sin 4a \sin a = \cos 3a \cos 2a - \sin 3a \sin 2a.$

### Functions of Multiple Angles.

120. *To express  $\sin 2A$  in terms of  $\sin A$  and  $\cos A$ .*

$$\sin 2A = \sin (A + A) = \sin A \cos A + \cos A \sin A;$$

that is,  $\sin 2A = 2 \sin A \cos A.$

Since  $A$  may have any value, this is a perfectly general formula for the sine of an angle in terms of the sine and cosine of the half angle. Thus if  $2A$  be replaced by  $\theta$ , we have

$$\sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}.$$

Similarly,  $\sin 4A = 2 \sin 2A \cos 2A$   
 $= 4 \sin A \cos A \cos 2A.$

121. *To express  $\cos 2A$  in terms of  $\cos A$  and  $\sin A$ .*

$$\cos 2A = \cos (A + A) = \cos A \cos A - \sin A \sin A;$$

that is,  $\cos 2A = \cos^2 A - \sin^2 A \dots\dots\dots(1).$

There are two other useful forms in which  $\cos 2A$  may be expressed, one involving  $\cos A$  only, the other  $\sin A$  only.

Thus from (1),

$$\cos 2A = \cos^2 A - (1 - \cos^2 A);$$

that is,  $\cos 2A = 2 \cos^2 A - 1 \dots\dots\dots(2).$

Again, from (1),

$$\cos 2A = (1 - \sin^2 A) - \sin^2 A;$$

that is,  $\cos 2A = 1 - 2 \sin^2 A \dots\dots\dots(3).$

From formulæ (2) and (3), we obtain by transposition

$$1 + \cos 2A = 2 \cos^2 A \dots\dots\dots(4),$$

and  $1 - \cos 2A = 2 \sin^2 A \dots\dots\dots(5).$

By division,  $\frac{1 + \cos 2A}{1 - \cos 2A} = \tan^2 A \dots\dots\dots(6).$

*Example.* Express  $\cos 4a$  in terms of  $\sin a$ .

From (3),  $\cos 4a = 1 - 2 \sin^2 2a = 1 - 2 (4 \sin^2 a \cos^2 a)$   
 $= 1 - 8 \sin^2 a (1 - \sin^2 a)$   
 $= 1 - 8 \sin^2 a + 8 \sin^4 a.$

122. The six formulæ of the last article deserve special attention. They are universally true so long as one of the angles involved is double of the other. For instance,

$$\cos a = \cos^2 \frac{a}{2} - \sin^2 \frac{a}{2},$$

$$\cos a = 2 \cos^2 \frac{a}{2} - 1, \quad \cos a = 1 - 2 \sin^2 \frac{a}{2},$$

$$1 + \cos \theta = 2 \cos^2 \frac{\theta}{2}, \quad 1 - \cos \theta = 2 \sin^2 \frac{\theta}{2}.$$

*Example.* If  $\cos \theta = .28$ , find the value of  $\tan \frac{\theta}{2}$ .

$$\tan^2 \frac{\theta}{2} = \frac{1 - \cos \theta}{1 + \cos \theta} = \frac{1 - .28}{1 + .28} = \frac{.72}{1.28} = \frac{72}{128} = \frac{9}{16};$$

$$\therefore \tan \frac{\theta}{2} = \pm \frac{3}{4}.$$

123. To express  $\tan 2A$  in terms of  $\tan A$ .

$$\tan 2A = \tan (A + A) = \frac{\tan A + \tan A}{1 - \tan A \tan A};$$

that is, 
$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}.$$

124. To express  $\sin 2A$  and  $\cos 2A$  in terms of  $\tan A$ .

$$\sin 2A = 2 \sin A \cos A = 2 \frac{\sin A}{\cos A} \cos^2 A = 2 \tan A \cos^2 A;$$

$$\therefore \sin 2A = \frac{2 \tan A}{\sec^2 A} = \frac{2 \tan A}{1 + \tan^2 A}.$$

Again,

$$\cos 2A = \cos^2 A - \sin^2 A = \cos^2 A (1 - \tan^2 A);$$

$$\therefore \cos 2A = \frac{1 - \tan^2 A}{\sec^2 A} = \frac{1 - \tan^2 A}{1 + \tan^2 A}.$$

*Example.* Shew that  $\frac{1 - \tan^2(45^\circ - A)}{1 + \tan^2(45^\circ - A)} = \sin 2A$ .

The first side  $= \cos 2(45^\circ - A) = \cos(90^\circ - 2A) = \sin 2A$ .

**EXAMPLES. XI. d.**

[The examples printed in more prominent type are important, and should be regarded as standard formulæ.]

1. If  $\cos A = \frac{1}{3}$ , find  $\cos 2A$ .
2. Find  $\cos 2A$  when  $\sin A = \frac{2}{5}$ .
3. If  $\sin A = \frac{3}{5}$ , find  $\sin 2A$ .
4. If  $\tan \theta = \frac{1}{3}$ , find  $\tan 2\theta$ .
5. If  $\tan \theta = \frac{1}{7}$ , find  $\sin 2\theta$  and  $\cos 2\theta$ .
6. If  $\cos a = \frac{4}{5}$ , find  $\tan \frac{a}{2}$ .
7. Find  $\tan A$  when  $\cos 2A = .96$ .

Prove the following identities :

- |  |   |
|--|---|
| 8. $\frac{\sin 2A}{1 + \cos 2A} = \tan A.$                             | 9. $\frac{\sin 2A}{1 - \cos 2A} = \cot A.$  |
| 10. $\frac{1 - \cos A}{\sin A} = \tan \frac{A}{2}.$                    | 11. $\frac{1 + \cos A}{\sin A} = \cot \frac{A}{2}.$   |
| 12. $2 \operatorname{cosec} 2a = \sec a \operatorname{cosec} a.$       |   |
| 13. $\tan a + \cot a = 2 \operatorname{cosec} 2a.$                     |   |
| 14. $\cos^4 a - \sin^4 a = \cos 2a.$                                   | 15. $\cot a - \tan a = 2 \cot 2a.$  |
| 16. $\cot 2A = \frac{\cot^2 A - 1}{2 \cot A}.$                         | 17. $\frac{\cot A - \tan A}{\cot A + \tan A} = \cos 2A.$                                      |
| 18. $\frac{1 + \cot^2 A}{2 \cot A} = \operatorname{cosec} 2A.$         | 19. $\frac{\cot^2 A + 1}{\cot^2 A - 1} = \sec 2A.$  |
| 20. $\frac{1 + \sec \theta}{\sec \theta} = 2 \cos^2 \frac{\theta}{2}.$ | 21. $\frac{\sec \theta - 1}{\sec \theta} = 2 \sin^2 \frac{\theta}{2}.$                        |
| 22. $\frac{2 - \sec^2 \theta}{\sec^2 \theta} = \cos 2\theta.$          | 23. $\frac{\operatorname{cosec}^2 \theta - 2}{\operatorname{cosec}^2 \theta} = \cos 2\theta.$ |

Prove the following identities :

$$24. \left( \sin \frac{A}{2} + \cos \frac{A}{2} \right)^2 = 1 + \sin A.$$

$$25. \left( \sin \frac{A}{2} - \cos \frac{A}{2} \right)^2 = 1 - \sin A.$$

$$26. \frac{\cos 2a}{1 + \sin 2a} = \tan (45^\circ - a).$$

$$27. \frac{\cos 2a}{1 - \sin 2a} = \cot (45^\circ - a).$$

$$28. \sin 8A = 8 \sin A \cos A \cos 2A \cos 4A.$$

$$29. \cos 4A = 8 \cos^4 A - 8 \cos^2 A + 1.$$

$$30. \sin A = 1 - 2 \sin^2 \left( 45^\circ - \frac{A}{2} \right).$$

$$31. \cos^2 \left( \frac{\pi}{4} - a \right) - \sin^2 \left( \frac{\pi}{4} - a \right) = \sin 2a.$$

$$32. \tan (45^\circ + A) - \tan (45^\circ - A) = 2 \tan 2A.$$

$$33. \tan (45^\circ + A) + \tan (45^\circ - A) = 2 \sec 2A.$$

### 125. Functions of $3A$ .

$$\begin{aligned} \sin 3A &= \sin (2A + A) = \sin 2A \cos A + \cos 2A \sin A \\ &= 2 \sin A \cos^2 A + (1 - 2 \sin^2 A) \sin A \\ &= 2 \sin A (1 - \sin^2 A) + (1 - 2 \sin^2 A) \sin A ; \\ &= 3 \sin A - 4 \sin^3 A. \end{aligned}$$

Similarly it may be proved that

$$\cos 3A = 4 \cos^3 A - 3 \cos A.$$

$$\text{Again, } \tan 3A = \tan (2A + A) = \frac{\tan 2A + \tan A}{1 - \tan 2A \tan A} ;$$

$$\text{by putting } \tan 2A = \frac{2 \tan A}{1 - \tan^2 A},$$

we obtain on reduction

$$\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}.$$

These formulæ are perfectly general and may be applied to cases of any two angles, one of which is three times the other; thus

$$\begin{aligned}\cos 6a &= 4 \cos^3 2a - 3 \cos 2a; \\ \sin 9A &= 3 \sin 3A - 4 \sin^3 3A.\end{aligned}$$

126. To find the value of  $\sin 18^\circ$ .

Let  $A = 18^\circ$ , then  $5A = 90^\circ$ , so that  $2A = 90^\circ - 3A$ .

$$\therefore \sin 2A = \sin (90^\circ - 3A) = \cos 3A;$$

$$\therefore 2 \sin A \cos A = 4 \cos^3 A - 3 \cos A.$$

Divide by  $\cos A$  (which is not equal to zero);

$$\therefore 2 \sin A = 4 \cos^2 A - 3 = 4(1 - \sin^2 A) - 3;$$

$$\therefore 4 \sin^2 A + 2 \sin A - 1 = 0;$$

$$\therefore \sin A = \frac{-2 \pm \sqrt{4+16}}{8} = \frac{-1 \pm \sqrt{5}}{4}.$$

Since  $18^\circ$  is an acute angle, we take the positive sign;

$$\therefore \sin 18^\circ = \frac{\sqrt{5}-1}{4}.$$

*Example.* Find  $\cos 18^\circ$  and  $\sin 54^\circ$ .

$$\cos 18^\circ = \sqrt{1 - \sin^2 18^\circ} = \sqrt{1 - \frac{6-2\sqrt{5}}{16}} = \frac{\sqrt{10+2\sqrt{5}}}{4}.$$

Since  $54^\circ$  and  $36^\circ$  are complementary,  $\sin 54^\circ = \cos 36^\circ$ .

$$\text{Now } \cos 36^\circ = 1 - 2 \sin^2 18^\circ = 1 - \frac{2(6-2\sqrt{5})}{16} = \frac{\sqrt{5}+1}{4};$$

$$\therefore \sin 54^\circ = \frac{\sqrt{5}+1}{4}.$$

### EXAMPLES. XI. e.

1. If  $\cos A = \frac{1}{3}$ , find  $\cos 3A$ .
2. Find  $\sin 3A$  when  $\sin A = \frac{3}{5}$ .
3. Given  $\tan A = 3$ , find  $\tan 3A$ .



Prove the following identities:

$$4. \frac{\sin 3A}{\sin A} - \frac{\cos 3A}{\cos A} = 2. \quad 5. \cot 3A = \frac{\cot^3 A - 3 \cot A}{3 \cot^2 A - 1}.$$

$$6. \frac{3 \cos a + \cos 3a}{3 \sin a - \sin 3a} = \cot^3 a. \quad 7. \frac{\sin 3a + \sin^3 a}{\cos^3 a - \cos 3a} = \cot a.$$

$$8. \frac{\cos^3 a - \cos 3a}{\cos a} + \frac{\sin^3 a + \sin 3a}{\sin a} = 3.$$

$$9. \sin 18^\circ + \sin 30^\circ = \sin 54^\circ. \quad 10. \cos 36^\circ - \sin 18^\circ = \frac{1}{2}.$$

$$11. \cos^2 36^\circ + \sin^2 18^\circ = \frac{3}{4}. \quad 12. 4 \sin 18^\circ \cos 36^\circ = 1.$$

127. The following examples further illustrate the formulæ proved in this chapter.

*Example 1.* Shew that  $\cos^6 a + \sin^6 a = 1 - \frac{3}{4} \sin^2 2a$ .

$$\begin{aligned} \text{The first side} &= (\cos^2 a + \sin^2 a)(\cos^4 a + \sin^4 a - \cos^2 a \sin^2 a) \\ &= (\cos^2 a + \sin^2 a)^2 - 3 \cos^2 a \sin^2 a \\ &= 1 - \frac{3}{4} (4 \cos^2 a \sin^2 a) \\ &= 1 - \frac{3}{4} \sin^2 2a. \end{aligned}$$

*Example 2.* Prove that  $\frac{\cos A - \sin A}{\cos A + \sin A} = \sec 2A - \tan 2A$ .

$$\text{The right side} = \frac{1}{\cos 2A} - \frac{\sin 2A}{\cos 2A} = \frac{1 - \sin 2A}{\cos 2A},$$

and since  $\cos 2A = \cos^2 A - \sin^2 A = (\cos A + \sin A)(\cos A - \sin A)$ , this suggests that we should multiply the numerator and denominator of the left side by  $\cos A - \sin A$ ; thus

$$\begin{aligned} \text{the first side} &= \frac{(\cos A - \sin A)(\cos A - \sin A)}{(\cos A + \sin A)(\cos A - \sin A)} \\ &= \frac{\cos^2 A + \sin^2 A - 2 \cos A \sin A}{\cos^2 A - \sin^2 A} \\ &= \frac{1 - \sin 2A}{\cos 2A} = \sec 2A - \tan 2A. \end{aligned}$$

*Example 3.* Shew that  $\frac{1}{\tan 3A - \tan A} - \frac{1}{\cot 3A - \cot A} = \cot 2A$ .

$$\begin{aligned} \text{The first side} &= \frac{1}{\frac{\sin 3A}{\cos 3A} - \frac{\sin A}{\cos A}} - \frac{1}{\frac{\cos 3A}{\sin 3A} - \frac{\cos A}{\sin A}} \\ &= \frac{\cos 3A \cos A}{\sin 3A \cos A - \cos 3A \sin A} - \frac{\sin 3A \sin A}{\cos 3A \sin A - \sin 3A \cos A} \\ &= \frac{\cos 3A \cos A + \sin 3A \sin A}{\sin 3A \cos A - \cos 3A \sin A} \\ &= \frac{\cos (3A - A)}{\sin (3A - A)} = \frac{\cos 2A}{\sin 2A} = \cot 2A. \end{aligned}$$

NOTE. This example has been given to emphasize the fact that in identities involving the functions of  $2A$  and  $3A$  it is sometimes best not to substitute their equivalents in terms of functions of  $A$ .

### EXAMPLES. XI. f.

Prove the following identities :

1.  $\tan 2A - \sec A \sin A = \tan A \sec 2A$ .
2.  $\tan 2A + \cos A \operatorname{cosec} A = \cot A \sec 2A$ .
3.  $\frac{1 - \cos 2\theta + \sin 2\theta}{1 + \cos 2\theta + \sin 2\theta} = \tan \theta$ .
4.  $\frac{1 + \cos \theta + \cos \frac{\theta}{2}}{\sin \theta + \sin \frac{\theta}{2}} = \cot \frac{\theta}{2}$ .
5.  $\cos^6 a - \sin^6 a = \cos 2a \left( 1 - \frac{1}{4} \sin^2 2a \right)$ .
6.  $4(\cos^6 \theta + \sin^6 \theta) = 1 + 3 \cos^2 2\theta$ .
7.  $\frac{\cos 3a + \sin 3a}{\cos a - \sin a} = 1 + 2 \sin 2a$ .
8.  $\frac{\cos 3a - \sin 3a}{\cos a + \sin a} = 1 - 2 \sin 2a$ .
9.  $\frac{\cos a + \sin a}{\cos a - \sin a} = \tan 2a + \sec 2a$ .

Prove the following identities :

$$10. \quad \frac{\cot a - 1}{\cot a + 1} = \frac{1 - \sin 2a}{\cos 2a}.$$

$$11. \quad \frac{1 + \sin \theta}{\cos \theta} = \frac{1 + \tan \frac{\theta}{2}}{1 - \tan \frac{\theta}{2}}.$$

$$12. \quad \frac{\cos \theta}{1 - \sin \theta} = \frac{\cot \frac{\theta}{2} + 1}{\cot \frac{\theta}{2} - 1}.$$

$$13. \quad \sec A - \tan A = \tan \left( 45^\circ - \frac{A}{2} \right).$$

$$14. \quad \tan A + \sec A = \cot \left( 45^\circ - \frac{A}{2} \right).$$

$$15. \quad \frac{1 + \sin \theta}{1 - \sin \theta} = \tan^2 \left( \frac{\pi}{4} + \frac{\theta}{2} \right).$$

$$16. \quad (2 \cos A + 1)(2 \cos A - 1) = 2 \cos 2A + 1.$$

$$17. \quad \frac{\sin 2A}{1 + \cos 2A} \cdot \frac{\cos A}{1 + \cos A} = \tan \frac{A}{2}.$$

$$18. \quad \frac{\sin 2A}{1 - \cos 2A} \cdot \frac{1 - \cos A}{\cos A} = \tan \frac{A}{2}.$$

$$19. \quad 4 \sin^3 a \cos 3a + 4 \cos^3 a \sin 3a = 3 \sin 4a.$$

[Put  $4 \sin^3 a = 3 \sin a - \sin 3a$  and  $4 \cos^3 a = 3 \cos a + \cos 3a$ .]

$$20. \quad \cos^3 a \cos 3a + \sin^3 a \sin 3a = \cos^3 2a.$$

$$21. \quad 4 (\cos^3 20^\circ + \cos^3 40^\circ) = 3 (\cos 20^\circ + \cos 40^\circ).$$

$$22. \quad 4 (\cos^3 10^\circ + \sin^3 20^\circ) = 3 (\cos 10^\circ + \sin 20^\circ).$$

$$23. \quad \tan 3A - \tan 2A - \tan A = \tan 3A \tan 2A \tan A.$$

[Use  $\tan 3A = \tan (2A + A)$ .]

$$24. \quad \frac{\cot \theta}{\cot \theta - \cot 3\theta} + \frac{\tan \theta}{\tan \theta - \tan 3\theta} = 1.$$

$$25. \quad \frac{1}{\tan 3\theta + \tan \theta} - \frac{1}{\cot 3\theta + \cot \theta} = \cot 4\theta.$$

## CHAPTER XII.

### TRANSFORMATION OF PRODUCTS AND SUMS.

#### Transformation of products into sums or differences.

128. In the last chapter we have proved that

$$\sin A \cos B + \cos A \sin B = \sin (A + B),$$

and  $\sin A \cos B - \cos A \sin B = \sin (A - B).$

By addition,

$$2 \sin A \cos B = \sin (A + B) + \sin (A - B) \dots\dots\dots(1);$$

by subtraction

$$2 \cos A \sin B = \sin (A + B) - \sin (A - B) \dots\dots\dots(2).$$

These formulæ enable us to express the product of a sine and cosine as the sum or difference of two sines.

Again,  $\cos A \cos B - \sin A \sin B = \cos (A + B),$

and  $\cos A \cos B + \sin A \sin B = \cos (A - B).$

By addition,

$$2 \cos A \cos B = \cos (A + B) + \cos (A - B) \dots\dots\dots(3);$$

by subtraction,

$$2 \sin A \sin B = \cos (A - B) - \cos (A + B) \dots\dots\dots(4).$$

These formulæ enable us to express

- (i) the product of two cosines as the sum of two cosines;
- (ii) the product of two sines as the difference of two cosines.

129. In each of the four formulæ of the previous article it should be noticed that on the left side we have any two angles  $A$  and  $B$ , and on the right side the sum and difference of these angles.

For practical purposes the following verbal statements of the results are more useful.

$$2 \sin A \cos B = \sin (\text{sum}) + \sin (\text{difference});$$

$$2 \cos A \sin B = \sin (\text{sum}) - \sin (\text{difference});$$

$$2 \cos A \cos B = \cos (\text{sum}) + \cos (\text{difference});$$

$$2 \sin A \sin B = \cos (\text{difference}) - \cos (\text{sum}).$$

N.B. In the last of these formulæ, *the difference precedes the sum.*

$$\begin{aligned} \text{Example 1. } 2 \sin 7A \cos 4A &= \sin (\text{sum}) + \sin (\text{difference}) \\ &= \sin 11A + \sin 3A. \end{aligned}$$

$$\begin{aligned} \text{Example 2. } 2 \cos 3\theta \sin 6\theta &= \sin (3\theta + 6\theta) - \sin (3\theta - 6\theta) \\ &= \sin 9\theta - \sin (-3\theta) \\ &= \sin 9\theta + \sin 3\theta. \end{aligned}$$

$$\begin{aligned} \text{Example 3. } \cos \frac{3A}{2} \cos \frac{5A}{2} &= \frac{1}{2} \left\{ \cos \left( \frac{3A}{2} + \frac{5A}{2} \right) + \cos \left( \frac{3A}{2} - \frac{5A}{2} \right) \right\} \\ &= \frac{1}{2} \{ \cos 4A + \cos (-A) \} \\ &= \frac{1}{2} (\cos 4A + \cos A). \end{aligned}$$

$$\begin{aligned} \text{Example 4. } 2 \sin 75^\circ \sin 15^\circ &= \cos (75^\circ - 15^\circ) - \cos (75^\circ + 15^\circ) \\ &= \cos 60^\circ - \cos 90^\circ \\ &= \frac{1}{2} - 0 \\ &= \frac{1}{2}. \end{aligned}$$

**130.** After a little practice the student will be able to omit some of the steps and find the equivalent very rapidly.

$$\text{Example 1. } 2 \cos \left( \frac{\pi}{4} + \theta \right) \cos \left( \frac{\pi}{4} - \theta \right) = \cos \frac{\pi}{2} + \cos 2\theta = \cos 2\theta.$$

$$\begin{aligned} \text{Example 2. } \sin (\alpha - 2\beta) \cos (\alpha + 2\beta) &= \frac{1}{2} \{ \sin 2\alpha + \sin (-4\beta) \} \\ &= \frac{1}{2} (\sin 2\alpha - \sin 4\beta). \end{aligned}$$

**EXAMPLES. XII. a.**

Express in the form of a sum or difference

- |   |  |
|---|--|
| 1. $2 \sin 3\theta \cos \theta$ .                       | 2. $2 \cos 6\theta \sin 3\theta$ .                   |
| 3. $2 \cos 7A \cos 5A$ .                                | 4. $2 \sin 3A \sin 2A$ .                             |
| 5. $2 \cos 5\theta \sin 4\theta$ .                      | 6. $2 \sin 4\theta \cos 8\theta$ .                   |
| 7. $2 \sin 9\theta \sin 3\theta$ .                      | 8. $2 \cos 9\theta \sin 7\theta$ .                   |
| 9. $2 \cos 2a \cos 11a$ .                               | 10. $2 \sin 5a \sin 10a$ .                           |
| 11. $\sin 4a \cos 7a$ .                                 | 12. $\sin 3a \sin a$ .                               |
| 13. $\cos \frac{A}{2} \sin \frac{3A}{2}$ .              | 14. $\sin \frac{5A}{2} \cos \frac{7A}{2}$ .          |
| 15. $2 \cos \frac{2\theta}{3} \cos \frac{5\theta}{3}$ . | 16. $\sin \frac{\theta}{4} \sin \frac{3\theta}{4}$ . |
| 17. $2 \cos 2\beta \cos (a - \beta)$ .                  | 18. $2 \sin 3a \sin (a + \beta)$ .                   |
| 19. $2 \sin (2\theta + \phi) \cos (\theta - 2\phi)$ .   |  |
| 20. $2 \cos (3\theta + \phi) \sin (\theta - 2\phi)$ .   |  |
| 21. $\cos (60^\circ + a) \sin (60^\circ - a)$ .         |  |

**Transformation of sums or differences into products.**

131. Since  $\sin (A + B) = \sin A \cos B + \cos A \sin B$ ,  
 and  $\sin (A - B) = \sin A \cos B - \cos A \sin B$ ;  
 by addition,  

$$\sin (A + B) + \sin (A - B) = 2 \sin A \cos B \dots\dots\dots(1);$$
  
 by subtraction,  

$$\sin (A + B) - \sin (A - B) = 2 \cos A \sin B \dots\dots\dots(2).$$
  
 Again,  $\cos (A + B) = \cos A \cos B - \sin A \sin B$ ,  
 and  $\cos (A - B) = \cos A \cos B + \sin A \sin B$ .  
 By addition,  

$$\cos (A + B) + \cos (A - B) = 2 \cos A \cos B \dots\dots\dots(3);$$
  
 by subtraction,  

$$\begin{aligned} \cos (A + B) - \cos (A - B) &= -2 \sin A \sin B \\ &= 2 \sin A \sin (-B) \dots\dots\dots(4). \end{aligned}$$



Let  $A+B=C$ , and  $A-B=D$ ;  
 then  $A=\frac{C+D}{2}$ , and  $B=\frac{C-D}{2}$ .

By substituting for  $A$  and  $B$  in the formulæ (1), (2), (3), (4), we obtain

$$\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2},$$

$$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2},$$

$$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2},$$

$$\cos C - \cos D = 2 \sin \frac{C+D}{2} \sin \frac{D-C}{2}.$$

132. In practice, it is more convenient to quote the formulæ we have just obtained verbally as follows :

sum of two sines  $= 2 \sin (\text{half-sum}) \cos (\text{half-difference})$ ;

difference of two sines  $= 2 \cos (\text{half-sum}) \sin (\text{half-difference})$ ;

sum of two cosines  $= 2 \cos (\text{half-sum}) \cos (\text{half-difference})$ ;

difference of two cosines

$$= 2 \sin (\text{half-sum}) \sin (\text{half-difference reversed})$$

*Example 1.*  $\sin 14\theta + \sin 6\theta = 2 \sin \frac{14\theta + 6\theta}{2} \cos \frac{14\theta - 6\theta}{2}$   
 $= 2 \sin 10\theta \cos 4\theta.$

*Example 2.*  $\sin 9A - \sin 7A = 2 \cos \frac{9A + 7A}{2} \sin \frac{9A - 7A}{2}$   
 $= 2 \cos 8A \sin A.$

*Example 3.*  $\cos A + \cos 8A = 2 \cos \frac{9A}{2} \cos \left( -\frac{7A}{2} \right)$   
 $= 2 \cos \frac{9A}{2} \cos \frac{7A}{2}.$

*Example 4.*  $\cos 70^\circ - \cos 10^\circ = 2 \sin 40^\circ \sin (-30^\circ)$   
 $= -2 \sin 40^\circ \sin 30^\circ = -\sin 40^\circ.$

## EXAMPLES. XII. b.

Express in the form of a product

- |                                      |                                      |
|--------------------------------------|--------------------------------------|
| 1. $\sin 8\theta + \sin 4\theta.$    | 2. $\sin 5\theta - \sin \theta.$     |
| 3. $\cos 7\theta + \cos 3\theta.$    | 4. $\cos 9\theta - \cos 11\theta.$   |
| 5. $\sin 7a - \sin 5a.$              | 6. $\cos 3a + \cos 8a.$              |
| 7. $\sin 3a + \sin 13a.$             | 8. $\cos 5a - \cos a.$               |
| 9. $\cos 2A + \cos 9A.$              | 10. $\sin 3A - \sin 11A.$            |
| 11. $\cos 10^\circ - \cos 50^\circ.$ | 12. $\sin 70^\circ + \sin 50^\circ.$ |

Prove that

- |  |   |
|--|---|
| 13. $\frac{\cos a - \cos 3a}{\sin 3a - \sin a} = \tan 2a.$   | 14. $\frac{\sin 2a + \sin 3a}{\cos 2a - \cos 3a} = \cot \frac{a}{2}.$                   |
| 15. $\frac{\cos 4\theta - \cos \theta}{\sin \theta - \sin 4\theta} = \tan \frac{5\theta}{2}.$                                      | 16. $\frac{\cos 2\theta - \cos 12\theta}{\sin 12\theta + \sin 2\theta} = \tan 5\theta.$ |
| 17. $\sin (60^\circ + A) - \sin (60^\circ - A) = \sin A.$  |   |
| 18. $\cos (30^\circ - A) + \cos (30^\circ + A) = \sqrt{3} \cos A.$   |   |
| 19. $\cos \left( \frac{\pi}{4} + a \right) - \cos \left( \frac{\pi}{4} - a \right) = -\sqrt{2} \sin a.$                            |   |
| 20. $\frac{\cos (2a - 3\beta) + \cos 3\beta}{\sin (2a - 3\beta) + \sin 3\beta} = \cot a.$  |   |
| 21. $\frac{\cos (\theta - 3\phi) - \cos (3\theta + \phi)}{\sin (3\theta + \phi) + \sin (\theta - 3\phi)} = \tan (\theta + 2\phi).$ |   |
| 22. $\frac{\sin (a + \beta) - \sin 4\beta}{\cos (a + \beta) + \cos 4\beta} = \tan \frac{a - 3\beta}{2}.$                           |   |

133. The eight formulæ proved in this chapter are of the utmost importance and very little further progress can be made until they have been thoroughly learnt. In the first group, the transformation is from products to sums and differences; in the second group, there is the converse transformation from sums and differences to products.

Many examples admit of solution by applying either of these transformations, but it is absolutely necessary that the student should master all the formulæ and apply them with equal readiness.

134. The following examples should be studied with great care.

*Example 1.* Prove that

$$\sin 5A + \sin 2A - \sin A = \sin 2A (2 \cos 3A + 1).$$

$$\begin{aligned} \text{The first side} &= (\sin 5A - \sin A) + \sin 2A \\ &= 2 \cos 3A \sin 2A + \sin 2A \\ &= \sin 2A (2 \cos 3A + 1). \end{aligned}$$

*Example 2.* Prove that

$$\cos 2\theta \cos \theta - \sin 4\theta \sin \theta = \cos 3\theta \cos 2\theta.$$

$$\begin{aligned} \text{The first side} &= \frac{1}{2} (\cos 3\theta + \cos \theta) - \frac{1}{2} (\cos 3\theta - \cos 5\theta) \\ &= \frac{1}{2} (\cos \theta + \cos 5\theta) \\ &= \cos 3\theta \cos 2\theta. \end{aligned}$$

*Example 3.* Find the value of

$$\cos 20^\circ + \cos 100^\circ + \cos 140^\circ.$$

$$\begin{aligned} \text{The expression} &= \cos 20^\circ + (\cos 100^\circ + \cos 140^\circ) \\ &= \cos 20^\circ + 2 \cos 120^\circ \cos 20^\circ \\ &= \cos 20^\circ + 2 \left( -\frac{1}{2} \right) \cos 20^\circ \\ &= \cos 20^\circ - \cos 20^\circ = 0. \end{aligned}$$

*Example 4.* Express as the product of four sines

$$\sin (\beta + \gamma - \alpha) + \sin (\gamma + \alpha - \beta) + \sin (\alpha + \beta - \gamma) - \sin (\alpha + \beta + \gamma).$$

$$\begin{aligned} \text{The expression} &= 2 \sin \gamma \cos (\beta - \alpha) + 2 \cos (\alpha + \beta) \sin (-\gamma) \\ &= 2 \sin \gamma \{ \cos (\beta - \alpha) - \cos (\alpha + \beta) \} \\ &= 2 \sin \gamma (2 \sin \beta \sin \alpha) \\ &= 4 \sin \alpha \sin \beta \sin \gamma. \end{aligned}$$

*Example 5.* Express  $4 \cos \alpha \cos \beta \cos \gamma$  as the sum of four cosines.

$$\begin{aligned} \text{The expression} &= 2 \cos \alpha \{ \cos (\beta + \gamma) + \cos (\beta - \gamma) \} \\ &= 2 \cos \alpha \cos (\beta + \gamma) + 2 \cos \alpha \cos (\beta - \gamma) \\ &= \cos (\alpha + \beta + \gamma) + \cos (\alpha - \beta - \gamma) + \cos (\alpha + \beta - \gamma) + \cos (\alpha - \beta + \gamma) \\ &= \cos (\alpha + \beta + \gamma) + \cos (\beta + \gamma - \alpha) + \cos (\gamma + \alpha - \beta) + \cos (\alpha + \beta - \gamma). \end{aligned}$$

*Example 6.* Prove that  $\sin^2 5x - \sin^2 3x = \sin 8x \sin 2x$ .

*First solution.*

$$\begin{aligned}\sin^2 5x - \sin^2 3x &= (\sin 5x + \sin 3x)(\sin 5x - \sin 3x) \\ &= (2 \sin 4x \cos x)(2 \cos 4x \sin x) \\ &= (2 \sin 4x \cos 4x)(2 \sin x \cos x) \\ &= \sin 8x \sin 2x.\end{aligned}$$

*Second solution.*

$$\begin{aligned}\sin 8x \sin 2x &= \frac{1}{2} (\cos 6x - \cos 10x) \\ &= \frac{1}{2} \{1 - 2 \sin^2 3x - (1 - 2 \sin^2 5x)\} \\ &= \sin^2 5x - \sin^2 3x.\end{aligned}$$

*Third solution.*

By using the formula of Art. 114 we have at once

$$\sin^2 5x - \sin^2 3x = \sin (5x + 3x) \sin (5x - 3x) = \sin 8x \sin 2x.$$

## EXAMPLES. XII. c.

Prove the following identities :

1.  $\cos 3A + \sin 2A - \sin 4A = \cos 3A (1 - 2 \sin A).$
2.  $\sin 3\theta - \sin \theta - \sin 5\theta = \sin 3\theta (1 - 2 \cos 2\theta).$
3.  $\cos \theta + \cos 2\theta + \cos 5\theta = \cos 2\theta (1 + 2 \cos 3\theta).$
4.  $\sin a - \sin 2a + \sin 3a = 4 \sin \frac{a}{2} \cos a \cos \frac{3a}{2}.$
5.  $\sin 3a + \sin 7a + \sin 10a = 4 \sin 5a \cos \frac{7a}{2} \cos \frac{3a}{2}.$
6.  $\sin A + 2 \sin 3A + \sin 5A = 4 \sin 3A \cos^2 A.$
7.  $\frac{\sin 2a + \sin 5a - \sin a}{\cos 2a + \cos 5a + \cos a} = \tan 2a.$
8.  $\frac{\sin a + \sin 2a + \sin 4a + \sin 5a}{\cos a + \cos 2a + \cos 4a + \cos 5a} = \tan 3a.$
9.  $\frac{\cos 7\theta + \cos 3\theta - \cos 5\theta - \cos \theta}{\sin 7\theta - \sin 3\theta - \sin 5\theta + \sin \theta} = \cot 2\theta.$
10.  $\cos 3A \sin 2A - \cos 4A \sin A = \cos 2A \sin A.$

Prove the following identities :

$$11. \cos 5A \cos 2A - \cos 4A \cos 3A = -\sin 2A \sin A.$$

$$12. \sin 4\theta \cos \theta - \sin 3\theta \cos 2\theta = \sin \theta \cos 2\theta.$$

$$13. \cos 5^\circ - \sin 25^\circ = \sin 35^\circ.$$

[Use  $\sin 25^\circ = \cos 65^\circ$ .]

$$14. \sin 65^\circ + \cos 65^\circ = \sqrt{2} \cos 20^\circ.$$

$$15. \cos 80^\circ + \cos 40^\circ - \cos 20^\circ = 0.$$

$$16. \sin 78^\circ - \sin 18^\circ + \cos 132^\circ = 0.$$

$$17. \sin^2 5A - \sin^2 2A = \sin 7A \sin 3A.$$

$$18. \cos 2A \cos 5A = \cos^2 \frac{7A}{2} - \sin^2 \frac{3A}{2}.$$

$$19. \sin (a + \beta + \gamma) + \sin (a - \beta - \gamma) + \sin (a + \beta - \gamma) \\ + \sin (a - \beta + \gamma) = 4 \sin a \cos \beta \cos \gamma.$$

$$20. \cos (\beta + \gamma - a) - \cos (\gamma + a - \beta) + \cos (a + \beta - \gamma) \\ - \cos (a + \beta + \gamma) = 4 \sin a \cos \beta \sin \gamma.$$

$$21. \sin 2a + \sin 2\beta + \sin 2\gamma - \sin 2(a + \beta + \gamma) \\ = 4 \sin (\beta + \gamma) \sin (\gamma + a) \sin (a + \beta).$$

$$22. \cos a + \cos \beta + \cos \gamma + \cos (a + \beta + \gamma) \\ = 4 \cos \frac{\beta + \gamma}{2} \cos \frac{\gamma + a}{2} \cos \frac{a + \beta}{2}.$$

$$23. 4 \sin A \sin (60^\circ + A) \sin (60^\circ - A) = \sin 3A.$$

$$24. 4 \cos \theta \cos \left( \frac{2\pi}{3} + \theta \right) \cos \left( \frac{2\pi}{3} - \theta \right) = \cos 3\theta.$$

$$25. \cos \theta + \cos \left( \frac{2\pi}{3} - \theta \right) + \cos \left( \frac{2\pi}{3} + \theta \right) = 0.$$

$$26. \cos^2 A + \cos^2 (60^\circ + A) + \cos^2 (60^\circ - A) = \frac{3}{2}.$$

[Put  $2 \cos^2 A = 1 + \cos 2A$ .]

$$27. \sin^2 A + \sin^2 (120^\circ + A) + \sin^2 (120^\circ - A) = \frac{3}{2}.$$

$$28. \cos 20^\circ \cos 40^\circ \cos 80^\circ = \frac{1}{8}.$$

$$29. \sin 20^\circ \sin 40^\circ \sin 80^\circ = \frac{1}{8} \sqrt{3}.$$

**135.** Many interesting identities can be established connecting the functions of the three angles  $A, B, C$ , which satisfy the relation  $A + B + C = 180^\circ$ . In proving these it will be necessary to keep clearly in view the properties of complementary and supplementary angles. [Arts. 39 and 96.]

From the given relation, the sum of any two of the angles is the supplement of the third; thus

$$\begin{aligned}\sin(B+C) &= \sin A, & \cos(A+B) &= -\cos C, \\ \tan(C+A) &= -\tan B, & \cos B &= -\cos(C+A), \\ \sin C &= \sin(A+B), & \cot A &= -\cot(B+C).\end{aligned}$$

Again,  $\frac{A}{2} + \frac{B}{2} + \frac{C}{2} = 90^\circ$ , so that each half angle is the complement of the sum of the other two; thus

$$\begin{aligned}\cos \frac{A+B}{2} &= \sin \frac{C}{2}, & \sin \frac{C+A}{2} &= \cos \frac{B}{2}, & \tan \frac{B+C}{2} &= \cot \frac{A}{2}, \\ \cos \frac{C}{2} &= \sin \frac{A+B}{2}, & \sin \frac{A}{2} &= \cos \frac{B+C}{2}, & \tan \frac{B}{2} &= \cot \frac{C+A}{2}.\end{aligned}$$

*Example 1.* If  $A + B + C = 180^\circ$ , prove that

$$\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C.$$

The first side  $= 2 \sin(A+B) \cos(A-B) + 2 \sin C \cos C$

$$= 2 \sin C \cos(A-B) + 2 \sin C \cos C$$

$$= 2 \sin C \{\cos(A-B) + \cos C\}$$

$$= 2 \sin C \{\cos(A-B) - \cos(A+B)\}$$

$$= 2 \sin C \times 2 \sin A \sin B$$

$$= 4 \sin A \sin B \sin C.$$

*Example 2.* If  $A + B + C = 180^\circ$ , prove that

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

Since  $A + B$  is the supplement of  $C$ , we have

$$\tan(A+B) = -\tan C;$$

$$\therefore \frac{\tan A + \tan B}{1 - \tan A \tan B} = -\tan C;$$

whence by multiplying up and rearranging,

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$



*Example 3.* If  $A + B + C = 180^\circ$ , prove that

$$\cos A + \cos B + \cos C = 1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$$

$$\begin{aligned} \text{The first side} &= 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2} + \cos C \\ &= 2 \sin \frac{C}{2} \cos \frac{A-B}{2} + 1 - 2 \sin^2 \frac{C}{2} \\ &= 1 + 2 \sin \frac{C}{2} \left( \cos \frac{A-B}{2} - \sin \frac{C}{2} \right) \\ &= 1 + 2 \sin \frac{C}{2} \left( \cos \frac{A-B}{2} - \cos \frac{A+B}{2} \right) \\ &= 1 + 2 \sin \frac{C}{2} \left( 2 \sin \frac{A}{2} \sin \frac{B}{2} \right) \\ &= 1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}. \end{aligned}$$

### EXAMPLES. XII. d.

If  $A + B + C = 180^\circ$ , prove that

1.  $\sin 2A - \sin 2B + \sin 2C = 4 \cos A \sin B \cos C.$
  2.  $\sin 2A - \sin 2B - \sin 2C = -4 \sin A \cos B \cos C.$
  3.  $\sin A + \sin B + \sin C = 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}.$
  4.  $\sin A + \sin B - \sin C = 4 \sin \frac{A}{2} \sin \frac{B}{2} \cos \frac{C}{2}.$
  5.  $\cos A - \cos B + \cos C = 4 \cos \frac{A}{2} \sin \frac{B}{2} \cos \frac{C}{2} - 1.$
  6.  $\frac{\sin B + \sin C - \sin A}{\sin A + \sin B + \sin C} = \tan \frac{B}{2} \tan \frac{C}{2}.$
  7.  $\tan \frac{B}{2} \tan \frac{C}{2} + \tan \frac{C}{2} \tan \frac{A}{2} + \tan \frac{A}{2} \tan \frac{B}{2} = 1.$
- [Use  $\tan \frac{A+B}{2} = \cot \frac{C}{2}$ , and therefore  $\tan \frac{A+B}{2} \tan \frac{C}{2} = 1.$ ]



If  $A + B + C = 180^\circ$ , prove that

8.  $\frac{1 + \cos A - \cos B + \cos C}{1 + \cos A + \cos B - \cos C} = \tan \frac{B}{2} \cot \frac{C}{2}.$
9.  $\cos 2A + \cos 2B + \cos 2C + 4 \cos A \cos B \cos C + 1 = 0.$
10.  $\cot B \cot C + \cot C \cot A + \cot A \cot B = 1.$
11.  $(\cot B + \cot C)(\cot C + \cot A)(\cot A + \cot B)$   
 $\quad = \operatorname{cosec} A \operatorname{cosec} B \operatorname{cosec} C.$
12.  $\cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C = 1.$   
 $\quad \quad \quad [Use \ 2 \cos^2 A = 1 + \cos 2A.]$
13.  $\sin^2 \frac{A}{2} + \sin^2 \frac{B}{2} + \sin^2 \frac{C}{2} = 1 - 2 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$
14.  $\cos^2 2A + \cos^2 2B + \cos^2 2C = 1 + 2 \cos 2A \cos 2B \cos 2C.$
15.  $\frac{\cot B + \cot C}{\tan B + \tan C} + \frac{\cot C + \cot A}{\tan C + \tan A} + \frac{\cot A + \cot B}{\tan A + \tan B} = 1.$
16.  $\frac{\tan A + \tan B + \tan C}{(\sin A + \sin B + \sin C)^2} = \frac{\tan \frac{A}{2} \tan \frac{B}{2} \tan \frac{C}{2}}{2 \cos A \cos B \cos C}.$

136. The following examples further illustrate the formulæ proved in this and the preceding chapter.

*Example 1.* Prove that  $\cot (A + 15^\circ) - \tan (A - 15^\circ) = \frac{4 \cos 2A}{2 \sin 2A + 1}.$

$$\begin{aligned}
 \text{The first side} &= \frac{\cos (A + 15^\circ)}{\sin (A + 15^\circ)} - \frac{\sin (A - 15^\circ)}{\cos (A - 15^\circ)} \\
 &= \frac{\cos (A + 15^\circ) \cos (A - 15^\circ) - \sin (A + 15^\circ) \sin (A - 15^\circ)}{\sin (A + 15^\circ) \cos (A - 15^\circ)} \\
 &= \frac{\cos \{(A + 15^\circ) + (A - 15^\circ)\}}{\sin (A + 15^\circ) \cos (A - 15^\circ)} \\
 &= \frac{2 \cos 2A}{2 \sin (A + 15^\circ) \cos (A - 15^\circ)} = \frac{2 \cos 2A}{\sin 2A + \sin 30^\circ} \\
 &= \frac{4 \cos 2A}{2 \sin 2A + 1}.
 \end{aligned}$$

NOTE. In dealing with expressions which involve numerical angles it is usually advisable to effect some simplification before substituting the known values of the functions of the angles, especially if these contain surds.

*Example 2.* Prove that

$$\cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} = 4 \cos \frac{\pi - A}{4} \cos \frac{\pi - B}{4} \cos \frac{\pi - C}{4}.$$

$$\begin{aligned} \text{The second side} &= 2 \cos \frac{\pi - A}{4} \left[ \cos \frac{2\pi - (B + C)}{4} + \cos \frac{B - C}{4} \right] \\ &= 2 \cos \frac{\pi - A}{4} \cos \frac{\pi + A}{4} + 2 \cos \frac{\pi - A}{4} \cos \frac{B - C}{4} \\ &= \left( \cos \frac{\pi}{2} + \cos \frac{A}{2} \right) + 2 \cos \frac{B + C}{4} \cos \frac{B - C}{4} \\ &= \cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2}. \end{aligned}$$

### EXAMPLES. XII. e.

Prove the following identities :

$$1. \quad \cos(a + \beta) \sin(a - \beta) + \cos(\beta + \gamma) \sin(\beta - \gamma) + \cos(\gamma + \delta) \sin(\gamma - \delta) + \cos(\delta + a) \sin(\delta - a) = 0.$$

$$2. \quad \frac{\sin(\beta - \gamma)}{\sin \beta \sin \gamma} + \frac{\sin(\gamma - a)}{\sin \gamma \sin a} + \frac{\sin(a - \beta)}{\sin a \sin \beta} = 0.$$

$$3. \quad \frac{\sin a + \sin \beta + \sin(a + \beta)}{\sin a + \sin \beta - \sin(a + \beta)} = \cot \frac{a}{2} \cot \frac{\beta}{2}.$$

$$4. \quad \sin a \cos(\beta + \gamma) - \sin \beta \cos(a + \gamma) = \cos \gamma \sin(a - \beta).$$

$$5. \quad \cos a \cos(\beta + \gamma) - \cos \beta \cos(a + \gamma) = \sin \gamma \sin(a - \beta).$$

$$6. \quad (\cos A - \sin A)(\cos 2A - \sin 2A) = \cos A - \sin 3A.$$

$$7. \quad \text{If } \tan \theta = \frac{b}{a}, \text{ prove that } a \cos 2\theta + b \sin 2\theta = a.$$

[See Art. 124.]

$$8. \quad \text{Prove that } \sin 2A + \cos 2A = \frac{(1 + \tan A)^2 - 2 \tan^2 A}{1 + \tan^2 A}.$$

$$9. \quad \text{Prove that } \sin 4A = \frac{4 \tan A (1 - \tan^2 A)}{(1 + \tan^2 A)^2}.$$

$$10. \quad \text{If } A + B = 45^\circ, \text{ prove that}$$

$$(1 + \tan A)(1 + \tan B) = 2.$$

Prove the following identities :

$$11. \cot(15^\circ - A) + \tan(15^\circ + A) = \frac{4 \cos 2A}{1 - 2 \sin 2A}.$$

$$12. \cot(15^\circ + A) + \tan(15^\circ + A) = \frac{4}{\cos 2A + \sqrt{3} \sin 2A}.$$

$$13. \tan(A + 30^\circ) \tan(A - 30^\circ) = \frac{1 - 2 \cos 2A}{1 + 2 \cos 2A}.$$

$$14. (2 \cos A + 1)(2 \cos A - 1)(2 \cos 2A - 1) = 2 \cos 4A + 1.$$

$$15. \tan(\beta - \gamma) + \tan(\gamma - \alpha) + \tan(\alpha - \beta) \\ = \tan(\beta - \gamma) \tan(\gamma - \alpha) \tan(\alpha - \beta).$$

$$16. \sin(\beta - \gamma) + \sin(\gamma - \alpha) + \sin(\alpha - \beta) \\ + 4 \sin \frac{\beta - \gamma}{2} \sin \frac{\gamma - \alpha}{2} \sin \frac{\alpha - \beta}{2} = 0.$$

$$17. \cos^2(\beta - \gamma) + \cos^2(\gamma - \alpha) + \cos^2(\alpha - \beta) \\ = 1 + 2 \cos(\beta - \gamma) \cos(\gamma - \alpha) \cos(\alpha - \beta).$$

$$18. \cos^2 \alpha + \cos^2 \beta - 2 \cos \alpha \cos \beta \cos(\alpha + \beta) = \sin^2(\alpha + \beta).$$

$$19. \sin^2 \alpha + \sin^2 \beta + 2 \sin \alpha \sin \beta \cos(\alpha + \beta) = \sin^2(\alpha + \beta).$$

$$20. \cos 12^\circ + \cos 60^\circ + \cos 84^\circ = \cos 24^\circ + \cos 48^\circ.$$

If  $A + B + C = 180^\circ$ , shew that

$$21. \cos \frac{A}{2} + \cos \frac{B}{2} + \cos \frac{C}{2} = 4 \cos \frac{B+C}{4} \cos \frac{C+A}{4} \cos \frac{A+B}{4}.$$

$$22. \cos \frac{A}{2} - \cos \frac{B}{2} + \cos \frac{C}{2} = 4 \cos \frac{\pi+A}{4} \cos \frac{\pi-B}{4} \cos \frac{\pi+C}{4}.$$

$$23. \sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2} = 1 + 4 \sin \frac{\pi-A}{4} \sin \frac{\pi-B}{4} \sin \frac{\pi-C}{4}.$$

If  $\alpha + \beta + \gamma = \frac{\pi}{2}$ , shew that

$$24. \frac{\sin 2\alpha + \sin 2\beta + \sin 2\gamma}{\sin 2\alpha + \sin 2\beta - \sin 2\gamma} = \cot \alpha \cot \beta.$$

$$25. \tan \beta \tan \gamma + \tan \gamma \tan \alpha + \tan \alpha \tan \beta = 1.$$

## CHAPTER XIII.

### RELATIONS BETWEEN THE SIDES AND ANGLES OF A TRIANGLE.

137. *In any triangle the sides are proportional to the sines of the opposite angles; that is,*

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$

(1) Let the triangle  $ABC$  be acute-angled.


From  $A$  draw  $AD$  perpendicular to the opposite side; then

$$AD = AB \sin ABD = c \sin B,$$

and  $AD = AC \sin ACD = b \sin C;$

$$\therefore b \sin C = c \sin B,$$

that is, 
$$\frac{b}{\sin B} = \frac{c}{\sin C}.$$



(2) Let the triangle  $ABC$  have an obtuse angle  $B$ .

Draw  $AD$  perpendicular to  $CB$  produced; then


$$AD = AC \sin ACD = b \sin C,$$

and  $AD = AB \sin ABD$

$$= c \sin (180^\circ - B) = c \sin B;$$

$$\therefore b \sin C = c \sin B;$$

that is, 
$$\frac{b}{\sin B} = \frac{c}{\sin C}.$$



In like manner it may be proved that either of these ratios is equal to  $\frac{a}{\sin A}.$

Thus 
$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$

**138.** *To find an expression for one side of a triangle in terms of the other two sides and the included angle.*


(1) Let  $ABC$  be an acute-angled triangle.

Draw  $BD$  perpendicular to  $AC$ ;  
then by Euc. II. 13,

$$AB^2 = BC^2 + CA^2 - 2AC \cdot CD;$$

$$\therefore c^2 = a^2 + b^2 - 2b \cdot a \cos C$$

$$= a^2 + b^2 - 2ab \cos C.$$



(2) Let the triangle  $ABC$  have an obtuse angle  $C$ .


Draw  $BD$  perpendicular to  $AC$   
produced; then by Euc. II. 12,

$$AB^2 = BC^2 + CA^2 + 2AC \cdot CD;$$

$$\therefore c^2 = a^2 + b^2 + 2b \cdot a \cos BCD$$

$$= a^2 + b^2 + 2ab \cos (180^\circ - C)$$

$$= a^2 + b^2 - 2ab \cos C.$$



Hence in each case,  $c^2 = a^2 + b^2 - 2ab \cos C$ .

Similarly it may be shewn that

$$a^2 = b^2 + c^2 - 2bc \cos A,$$

and

$$b^2 = c^2 + a^2 - 2ca \cos B.$$

**139.** From the formulæ of the last article, we obtain

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}; \quad \cos B = \frac{c^2 + a^2 - b^2}{2ca}; \quad \cos C = \frac{a^2 + b^2 - c^2}{2ab}.$$

These results enable us to find the cosines of the angles when the numerical values of the sides are given.

**140.** *To express one side of a triangle in terms of the adjacent angles and the other two sides.*


(1) Let  $ABC$  be an acute-angled triangle.

Draw  $AD$  perpendicular to  $BC$ ; then

$$BC = BD + CD$$

$$= AB \cos ABD + AC \cos ACD;$$

that is,  $a = c \cos B + b \cos C$ .



(2) Let the triangle  $ABC$  have an obtuse angle  $C$ .


Draw  $AD$  perpendicular to  $BC$  produced; then

$$BC = BD - CD$$

$$= AB \cos ABD - AC \cos ACD;$$

$$\therefore a = c \cos B - b \cos (180^\circ - C)$$

$$= c \cos B + b \cos C.$$



Thus in each case  $a = b \cos C + c \cos B$ .

Similarly it may be shewn that

$$b = c \cos A + a \cos C, \text{ and } c = a \cos B + b \cos A.$$

NOTE. The formulæ we have proved in this chapter are quite general and may be regarded as the fundamental relations subsisting between the sides and angles of a triangle. The modified forms which they assume in the case of right-angled triangles have already been considered in Chap. V.; it will therefore be unnecessary in the present chapter to make any direct reference to right-angled triangles.

141. The sets of formulæ in Arts. 137, 138, and 140 have been established independently of one another; they are however not independent, for from any one set the other two may be derived by the help of the relation  $A + B + C = 180^\circ$ .

For instance, suppose we have proved as in Art. 137 that

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C};$$

then since  $\sin A = \sin (B + C) = \sin B \cos C + \sin C \cos B$ ;

$$\therefore 1 = \frac{\sin B}{\sin A} \cos C + \frac{\sin C}{\sin A} \cos B;$$

$$\therefore 1 = \frac{b}{a} \cos C + \frac{c}{a} \cos B;$$

$$\therefore a = b \cos C + c \cos B.$$

Similarly, we may prove that

$$b = c \cos A + a \cos C, \text{ and } c = a \cos B + b \cos A.$$

Multiplying these last three equations by  $a$ ,  $b$ ,  $-c$  respectively and adding, we have

$$a^2 + b^2 - c^2 = 2ab \cos C;$$

$$\therefore c^2 = a^2 + b^2 - 2ab \cos C.$$

Similarly the other relations of Art. 138 may be deduced.



### Solution of Triangles.

142. When any three parts of a triangle are given, provided that one at least of these is a side, the relations we have proved enable us to find the numerical values of the unknown parts. For from any equation which connects four quantities three of which are known the fourth may be found. Thus if  $c, a, B$  are given, we can find  $b$  from the formula

$$b^2 = c^2 + a^2 - 2ca \cos B;$$

and if  $B, C, b$  are given, we find  $c$  from the formula

$$\frac{c}{\sin C} = \frac{b}{\sin B}.$$

We may remark that if the three angles alone are given, the formula

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

enables us to find the *ratios* of the sides but not their actual *lengths*, and thus the triangle cannot be completely solved. In such a case there may be an infinite number of equiangular triangles all satisfying the data of the question. [See Euc. VI. 4.]

143. CASE I. *To solve a triangle having given the three sides.*

The angles  $A$  and  $B$  may be found from the formulæ

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}, \text{ and } \cos B = \frac{c^2 + a^2 - b^2}{2ca};$$

then the angle  $C$  is known from the equation  $C = 180^\circ - A - B$ .

*Example 1.* If  $a=7, b=5, c=8$ , find the angles  $A$  and  $B$ , having given that  $\cos 38^\circ 11' = \frac{11}{14}$ .

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{5^2 + 8^2 - 7^2}{2 \times 5 \times 8} = \frac{40}{2 \times 5 \times 8} = \frac{1}{2};$$

$$\therefore A = 60^\circ.$$

$$\cos B = \frac{c^2 + a^2 - b^2}{2ca} = \frac{8^2 + 7^2 - 5^2}{2 \times 8 \times 7} = \frac{88}{2 \times 8 \times 7} = \frac{11}{14};$$

$$\therefore B = 38^\circ 11'.$$



*Example 2.* Find the greatest angle of the triangle whose side are 6, 13, 11, having given that  $\cos 84^\circ 47' = \frac{1}{11}$ .

Let  $a=6$ ,  $b=13$ ,  $c=11$ . Since the greatest angle is opposite to the greatest side, the required angle is  $B$ .

$$\text{And } \cos B = \frac{c^2 + a^2 - b^2}{2ca} = \frac{11^2 + 6^2 - 13^2}{2 \times 11 \times 6} = \frac{-12}{2 \times 11 \times 6};$$

$$\therefore \cos B = -\frac{1}{11} = -\cos 84^\circ 47';$$

$$\therefore B = 180^\circ - 84^\circ 47' = 95^\circ 13'.$$

Thus the required angle is  $95^\circ 13'$ .

**144.** CASE II. *To solve a triangle having given two sides and the included angle.*

Let  $b, c, A$  be given; then  $a$  can be found from the formula

$$a^2 = b^2 + c^2 - 2bc \cos A.$$

We may now obtain  $B$  from either of the formulæ

$$\cos B = \frac{c^2 + a^2 - b^2}{2ca}, \text{ or } \sin B = \frac{b \sin A}{a};$$

then  $C$  is known from the equation  $C = 180^\circ - A - B$ .

*Example.* If  $a=3$ ,  $b=7$ ,  $C=98^\circ 13'$ , solve the triangle, having given  $\cos 81^\circ 47' = \frac{1}{7}$ .

$$\begin{aligned} c^2 &= a^2 + b^2 - 2ab \cos C \\ &= 9 + 49 - 2 \times 3 \times 7 \cos 98^\circ 13'. \end{aligned}$$

But  $98^\circ 13'$  is the supplement of  $81^\circ 47'$ ;

$$\begin{aligned} \therefore c^2 &= 58 + (2 \times 3 \times 7 \cos 81^\circ 47') \\ &= 58 + \left( 2 \times 3 \times 7 \times \frac{1}{7} \right) = 58 + 6 = 64; \end{aligned}$$

$$\therefore c = 8.$$

$$\cos B = \frac{c^2 + a^2 - b^2}{2ca} = \frac{64 + 9 - 49}{2 \times 8 \times 3} = \frac{24}{2 \times 8 \times 3} = \frac{1}{2};$$

$$\therefore B = 60^\circ.$$

$$C = 180^\circ - 60^\circ - 98^\circ 13' = 21^\circ 47'.$$

**145. CASE III.** *To solve a triangle having given two angles and a side.*

Let  $B, C, a$  be given.

The angle  $A$  is found from  $A = 180^\circ - B - C$ ; and the sides  $b$  and  $c$  from

$$b = \frac{a \sin B}{\sin A} \quad \text{and} \quad c = \frac{a \sin C}{\sin A}.$$

*Example.* If  $A = 105^\circ, C = 60^\circ, b = 4$ , solve the triangle.

$$B = 180^\circ - 105^\circ - 60^\circ = 15^\circ.$$

$$\therefore c = \frac{b \sin C}{\sin B} = \frac{4 \sin 60^\circ}{\sin 15^\circ} = \frac{4\sqrt{3}}{2} \cdot \frac{2\sqrt{2}}{\sqrt{3}-1} = \frac{4\sqrt{6}}{\sqrt{3}-1}$$

$$= \frac{4\sqrt{6}(\sqrt{3}+1)}{3-1} = 2\sqrt{6}(\sqrt{3}+1);$$

$$\therefore c = 6\sqrt{2} + 2\sqrt{6}.$$

$$a = \frac{b \sin A}{\sin B} = \frac{4 \sin 105^\circ}{\sin 15^\circ} = \frac{4 \sin 75^\circ}{\sin 15^\circ}$$

$$= 4 \times \frac{\sqrt{3}+1}{2\sqrt{2}} \times \frac{2\sqrt{2}}{\sqrt{3}-1} = \frac{4(\sqrt{3}+1)}{\sqrt{3}-1};$$

$$\therefore a = 4(2 + \sqrt{3}).$$

### EXAMPLES. XIII. a.

1. If  $a=15, b=7, c=13$ , find  $C$ .
2. If  $a=7, b=3, c=5$ , find  $A$ .
3. If  $a=5, b=5\sqrt{3}, c=5$ , find the angles.
4. If  $a=25, b=31, c=7\sqrt{2}$ , find  $A$ .
5. The sides of a triangle are 2,  $2\frac{2}{3}$ ,  $3\frac{1}{3}$ , find the greatest angle.
6. Solve the triangle when  $a=\sqrt{3}+1, b=2, c=\sqrt{6}$ .
7. Solve the triangle when  $a=\sqrt{2}, b=2, c=\sqrt{3}-1$ .
8. If  $a=8, b=5, c=\sqrt{19}$ , find  $C$ ; given  $\cos 28^\circ 56' = \frac{7}{8}$ .
9. If the sides are as 4 : 7 : 5, find the greatest angle;

$$\text{given } \cos 78^\circ 27' = \frac{1}{5}.$$

10. If  $a=2$ ,  $b=\sqrt{3}+1$ ,  $C=60^\circ$ , find  $c$ .
11. Given  $a=3$ ,  $c=5$ ,  $B=120^\circ$ , find  $b$ .
12. Given  $b=7$ ,  $c=6$ ,  $A=75^\circ 31'$ , find  $a$ ; given  $\cos 75^\circ 31' = .25$ .
13. If  $b=8$ ,  $c=11$ ,  $A=93^\circ 35'$ , find  $a$ ; given  $\cos 86^\circ 25' = .0625$ .
14. If  $a=7$ ,  $c=3$ ,  $B=123^\circ 12'$ , find  $b$ ; given  $\cos 56^\circ 48' = \frac{23}{42}$ .
15. Solve the triangle when  $a=2\sqrt{6}$ ,  $c=6-2\sqrt{3}$ ,  $B=75^\circ$ .
16. Solve the triangle when  $A=72^\circ$ ,  $b=2$ ,  $c=\sqrt{5}+1$ .
17. Given  $A=75^\circ$ ,  $B=30^\circ$ ,  $b=\sqrt{8}$ , solve the triangle.
18. If  $B=60^\circ$ ,  $C=15^\circ$ ,  $b=\sqrt{6}$ , solve the triangle.
19. If  $A=45^\circ$ ,  $B=105^\circ$ ,  $c=\sqrt{2}$ , solve the triangle.
20. Given  $A=45^\circ$ ,  $B=60^\circ$ , shew that  $c : a = \sqrt{3}+1 : 2$ .
21. If  $C=120^\circ$ ,  $c=2\sqrt{3}$ ,  $a=2$ , find  $b$ .
22. If  $B=60^\circ$ ,  $a=3$ ,  $b=3\sqrt{3}$ , find  $c$ .
23. Given  $(a+b+c)(b+c-a)=3bc$ , find  $A$ .
24. Find the angles of the triangle whose sides are  
 $3+\sqrt{3}$ ,  $2\sqrt{3}$ ,  $\sqrt{6}$ .
25. Find the angles of the triangle whose sides are  
 $\frac{\sqrt{3}+1}{2\sqrt{2}}$ ,  $\frac{\sqrt{3}-1}{2\sqrt{2}}$ ,  $\frac{\sqrt{3}}{2}$ .
26. Two sides of a triangle are  $\frac{1}{\sqrt{6}-\sqrt{2}}$  and  $\frac{1}{\sqrt{6}+\sqrt{2}}$ , and the included angle is  $60^\circ$ : solve the triangle.

146. When an angle of a triangle is obtained through the medium of the sine there may be ambiguity, for the sines of supplementary angles are equal in magnitude and are of the same sign, so that there are two angles less than  $180^\circ$  which have the same sine. When an angle is obtained through the medium of the cosine there is no ambiguity, for there is only one angle less than  $180^\circ$  whose cosine is equal to a given quantity.

Thus if  $\sin A = \frac{1}{2}$ , then  $A = 30^\circ$  or  $150^\circ$ ;  
 if  $\cos A = \frac{1}{2}$ , then  $A = 60^\circ$ .

*Example.* If  $C=60^\circ$ ,  $b=2\sqrt{3}$ ,  $c=3\sqrt{2}$ , find  $A$ .

From the equation  $\sin B = \frac{b \sin C}{c}$ ,

we have 
$$\sin B = \frac{2\sqrt{3}}{3\sqrt{2}} \cdot \frac{\sqrt{3}}{2} = \frac{1}{\sqrt{2}};$$

$$\therefore B = 45^\circ \text{ or } 135^\circ.$$

The value  $B=135^\circ$  is inadmissible, for in this case the sum of  $B$  and  $C$  would be greater than  $180^\circ$ .

Thus  $A = 180^\circ - 60^\circ - 45^\circ = 75^\circ.$

**147. CASE IV.** *To solve a triangle having given two sides and an angle opposite to one of them.*

Let  $a, b, A$  be given; then  $B$  is to be found from the equation

$$\sin B = \frac{b}{a} \sin A.$$

(i) If  $a < b \sin A$ , then  $\frac{b \sin A}{a} > 1$ , so that  $\sin B > 1$ , which is impossible. Thus there is no solution.

(ii) If  $a = b \sin A$ , then  $\frac{b \sin A}{a} = 1$ , so that  $\sin B = 1$ , and  $B$  has only the value  $90^\circ$ .

(iii) If  $a > b \sin A$ , then  $\frac{b \sin A}{a} < 1$ , and two values for  $B$  may be found from  $\sin B = \frac{b \sin A}{a}$ . These values are supplementary, so that one angle is acute, the other obtuse.

(1) If  $a < b$ , then  $A < B$ , and therefore  $B$  may either be acute or obtuse, so that both values are admissible. This is known as **the ambiguous case**.

(2) If  $a = b$ , then  $A = B$ ; and if  $a > b$ , then  $A > B$ ; in either case  $B$  cannot be obtuse, and therefore only the smaller value of  $B$  is admissible.

When  $B$  is found,  $C$  is determined from  $C = 180^\circ - A - B$ . Finally,  $c$  may be found from the equation  $c = \frac{a \sin C}{\sin A}$ .


From the foregoing investigation it appears that the only case in which an ambiguous solution can arise is when the smaller of the two given sides is opposite to the given angle.

148. To discuss the Ambiguous Case geometrically.


Let  $a, b, A$  be the given parts. Take a line  $AX$  unlimited towards  $X$ ; make  $\angle XAC$  equal to  $A$ , and  $AC$  equal to  $b$ . Draw  $CD$  perpendicular to  $AX$ , then  $CD = b \sin A$ .

With centre  $C$  and radius equal to  $a$  describe a circle.

(i) If  $a < b \sin A$ , the circle will not meet  $AX$ ; thus no triangle can be constructed with the given parts.




(ii) If  $a = b \sin A$ , the circle will touch  $AX$  at  $D$ ; thus there is a right-angled triangle with the given parts.



(iii) If  $a > b \sin A$ , the circle will cut  $AX$  in two points  $B_1, B_2$ .

(1) These points will be both on the same side of  $A$ , when  $a < b$ , in which case there are two solutions, namely the triangles

$$AB_1C, AB_2C.$$




This is the Ambiguous Case.

(2) The points  $B_1, B_2$  will be on opposite sides of  $A$  when  $a > b$ .

In this case there is only one solution, for the angle  $CAB_2$  is the supplement of the given angle, and thus the triangle  $AB_2C$  does not satisfy the data.

(3) If  $a = b$ , the point  $B_2$  coincides with  $A$ , so that there is only one solution.




*Example.* Given  $B=45^\circ$ ,  $c=\sqrt{12}$ ,  $b=\sqrt{8}$ , solve the triangle.

We have 
$$\sin C = \frac{c \sin B}{b} = \frac{2\sqrt{3}}{2\sqrt{2}} \cdot \frac{1}{\sqrt{2}} = \frac{\sqrt{3}}{2}.$$

$$\therefore C = 60^\circ \text{ or } 120^\circ,$$

and since  $b < c$ , both these values are admissible. The two triangles which satisfy the data are shewn in the figure.

Denote the sides  $BC_1$ ,  $BC_2$  by  $a_1$ ,  $a_2$ , and the angles  $BAC_1$ ,  $BAC_2$  by  $A_1$ ,  $A_2$  respectively.



(i) In the  $\triangle ABC_1$ ,  $\angle A_1 = 75^\circ$ ;

hence 
$$a_1 = \frac{b \sin A_1}{\sin B} = \frac{2\sqrt{2}}{\frac{1}{\sqrt{2}}} \cdot \frac{\sqrt{3}+1}{2\sqrt{2}} = \sqrt{2}(\sqrt{3}+1).$$

(ii) In the  $\triangle ABC_2$ ,  $\angle A_2 = 15^\circ$ ;

hence 
$$a_2 = \frac{b \sin A_2}{\sin B} = \frac{2\sqrt{2}}{\frac{1}{\sqrt{2}}} \cdot \frac{\sqrt{3}-1}{2\sqrt{2}} = \sqrt{2}(\sqrt{3}-1).$$

Thus the complete solution is 
$$\begin{cases} C = 60^\circ, \text{ or } 120^\circ; \\ A = 75^\circ, \text{ or } 15^\circ; \\ a = \sqrt{6} + \sqrt{2}, \text{ or } \sqrt{6} - \sqrt{2}. \end{cases}$$

### EXAMPLES. XIII. b.

1. Given  $a=1$ ,  $b=\sqrt{3}$ ,  $A=30^\circ$ , solve the triangle.
2. Given  $b=3\sqrt{2}$ ,  $c=2\sqrt{3}$ ,  $C=45^\circ$ , solve the triangle.
3. If  $C=60^\circ$ ,  $a=2$ ,  $c=\sqrt{6}$ , solve the triangle.



4. If  $A=30^\circ$ ,  $a=2$ ,  $c=5$ , solve the triangle.
5. If  $B=30^\circ$ ,  $b=\sqrt{6}$ ,  $c=2\sqrt{3}$ , solve the triangle.
6. If  $B=60^\circ$ ,  $b=3\sqrt{2}$ ,  $c=3+\sqrt{3}$ , solve the triangle.
7. If  $a=3+\sqrt{3}$ ,  $c=3-\sqrt{3}$ ,  $C=15^\circ$ , solve the triangle.
8. If  $A=18^\circ$ ,  $a=4$ ,  $b=4+\sqrt{80}$ , solve the triangle.
9. If  $B=135^\circ$ ,  $a=3\sqrt{2}$ ,  $b=2\sqrt{3}$ , solve the triangle.

149. Many relations connecting the sides and angles of a triangle may be proved by means of the formulæ we have established.

*Example 1.* Prove that  $(b-c) \cos \frac{A}{2} = a \sin \frac{B-C}{2}$ .

Let 
$$k = \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C};$$

then 
$$a = k \sin A, \quad b = k \sin B, \quad c = k \sin C;$$

$$\begin{aligned} \therefore (b-c) \cos \frac{A}{2} &= k (\sin B - \sin C) \cos \frac{A}{2} \\ &= 2k \cos \frac{B+C}{2} \sin \frac{B-C}{2} \cos \frac{A}{2} \\ &= 2k \sin \frac{A}{2} \cos \frac{A}{2} \sin \frac{B-C}{2} \\ &= k \sin A \sin \frac{B-C}{2} \\ &= a \sin \frac{B-C}{2}. \end{aligned}$$

*Example 2.* If  $a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2} = \frac{3b}{2}$ , shew that the sides of the triangle are in A.P.

Since 
$$2a \cos^2 \frac{C}{2} + 2c \cos^2 \frac{A}{2} = 3b,$$

$$\therefore a(1 + \cos C) + c(1 + \cos A) = 3b,$$

$$\therefore a + c + (a \cos C + c \cos A) = 3b,$$

$$\therefore a + c + b = 3b,$$

$$\therefore a + c = 2b.$$

Thus the sides  $a$ ,  $b$ ,  $c$  are in A.P.

*Example 3.* Prove that

$$(b^2 - c^2) \cot A + (c^2 - a^2) \cot B + (a^2 - b^2) \cot C = 0.$$

Let  $k = \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ ; then

the first side

$$\begin{aligned} &= k^2 \left\{ (\sin^2 B - \sin^2 C) \frac{\cos A}{\sin A} + \dots + \dots \right\} \\ &= k^2 \left\{ \sin(B+C) \sin(B-C) \frac{\cos A}{\sin A} + \dots + \dots \right\}. \quad [\text{Art. 114}]. \end{aligned}$$

But  $\sin(B+C) = \sin A$ , and  $\cos A = -\cos(B+C)$ ;

$\therefore$  the first side

$$\begin{aligned} &= -k^2 \{ \sin(B-C) \cos(B+C) + \dots + \dots \} \\ &= -\frac{k^2}{2} \{ (\sin 2B - \sin 2C) + (\sin 2C - \sin 2A) + (\sin 2A - \sin 2B) \} \\ &= 0. \end{aligned}$$

### EXAMPLES. XIII. c.

Prove the following identities :

1.  $a(\sin B - \sin C) + b(\sin C - \sin A) + c(\sin A - \sin B) = 0.$
2.  $2(bc \cos A + ca \cos B + ab \cos C) = a^2 + b^2 + c^2.$
3.  $a(b \cos C - c \cos B) = b^2 - c^2.$
4.  $(b+c) \cos A + (c+a) \cos B + (a+b) \cos C = a+b+c.$
5.  $2 \left( a \sin^2 \frac{C}{2} + c \sin^2 \frac{A}{2} \right) = c + a - b.$
6.  $\frac{\cos B}{\cos C} = \frac{c-b \cos A}{b-c \cos A}.$
7.  $\tan A = \frac{a \sin C}{b-a \cos C}.$
8.  $(b+c) \sin \frac{A}{2} = a \cos \frac{B-C}{2}.$
9.  $\frac{a+b}{c} \sin^2 \frac{C}{2} = \frac{\cos A + \cos B}{2}.$
10.  $a \sin(B-C) + b \sin(C-A) + c \sin(A-B) = 0.$
11.  $\frac{\sin(A-B)}{\sin(A+B)} = \frac{a^2 - b^2}{c^2}.$
12.  $\frac{c \sin(A-B)}{b \sin(C-A)} = \frac{a^2 - b^2}{c^2 - a^2}.$

[All articles and examples marked with an asterisk may be omitted on the first reading of the subject.]

\*150. The *ambiguous case* may also be discussed by first finding the third side.

As before, let  $a, b, A$  be given, then

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc};$$

$$\therefore c^2 - 2b \cos A \cdot c + b^2 - a^2 = 0.$$

By solving this quadratic equation in  $c$ , we obtain

$$\begin{aligned} c &= b \cos A \pm \sqrt{b^2 \cos^2 A + a^2 - b^2} \\ &= b \cos A \pm \sqrt{a^2 - b^2 \sin^2 A}. \end{aligned}$$

(i) When  $a < b \sin A$ , the quantity under the radical is negative, and the values of  $c$  are impossible; so that there is no solution.

(ii) When  $a = b \sin A$ , the quantity under the radical is zero, and  $c = b \cos A$ . Since  $\sin A < 1$ , it follows that  $a < b$ , and therefore  $A < B$ . Hence the triangle is impossible unless the given angle  $A$  is acute, in which case  $c$  is positive and there is one solution.

(iii) When  $a > b \sin A$ , there are three cases to consider.

(1) Suppose  $a < b$ , then  $A < B$ , and as before the triangle is impossible unless  $A$  is acute. In this case  $b \cos A$  is positive.

Also  $\sqrt{a^2 - b^2 \sin^2 A}$  is real and  $< \sqrt{b^2 - b^2 \sin^2 A}$ ;

that is  $\sqrt{a^2 - b^2 \sin^2 A} < b \cos A$ ;

hence both values of  $c$  are real and positive, so that there are two solutions.

(2) Suppose  $a > b$ , then  $\sqrt{a^2 - b^2 \sin^2 A} > \sqrt{b^2 - b^2 \sin^2 A}$ ; that is  $\sqrt{a^2 - b^2 \sin^2 A} > b \cos A$ ;

hence one value of  $c$  is positive and one value is negative, whether  $A$  is acute or obtuse, and in each case there is only one solution.

(3) Suppose  $a = b$ , then  $\sqrt{a^2 - b^2 \sin^2 A} = b \cos A$ ;

$$\therefore c = 2b \cos A \text{ or } 0;$$

hence there is only one solution when  $A$  is acute, and when  $A$  is obtuse the triangle is impossible.

*Example.* If  $b, c, B$  are given, and if  $b < c$ , shew that

$$(a_1 - a_2)^2 + (a_1 + a_2)^2 \tan^2 B = 4b^2,$$

where  $a_1, a_2$  are the two values of the third side.

From the formula  $\cos B = \frac{c^2 + a^2 - b^2}{2ca},$

we have  $a^2 - 2c \cos B \cdot a + c^2 - b^2 = 0.$

But the roots of this equation are  $a_1$  and  $a_2$ ; hence by the theory of quadratic equations

$$a_1 + a_2 = 2c \cos B \text{ and } a_1 a_2 = c^2 - b^2.$$

$$\begin{aligned} \therefore (a_1 - a_2)^2 &= (a_1 + a_2)^2 - 4a_1 a_2 \\ &= 4c^2 \cos^2 B - 4(c^2 - b^2). \end{aligned}$$

$$\begin{aligned} \therefore (a_1 - a_2)^2 + (a_1 + a_2)^2 \tan^2 B &= 4c^2 \cos^2 B - 4(c^2 - b^2) + 4c^2 \cos^2 B \tan^2 B \\ &= 4c^2 (\cos^2 B + \sin^2 B) - 4c^2 + 4b^2 \\ &= 4c^2 - 4c^2 + 4b^2 \\ &= 4b^2. \end{aligned}$$

### \*EXAMPLES. XIII. d.

1. In a triangle in which each base angle is double of the third angle the base is 2 : solve the triangle.

2. If  $B=45^\circ, C=75^\circ$ , and the perpendicular from  $A$  on  $BC$  is 3, solve the triangle.

3. If  $a=2, b=4-2\sqrt{3}, c=3\sqrt{2}-\sqrt{6}$ , solve the triangle.

4. If  $A=18^\circ, b-a=2, ab=4$ , find the other angles.

5. Given  $B=30^\circ, c=150, b=50\sqrt{3}$ , shew that of the two triangles which satisfy the data one will be isosceles and the other right-angled.

Find the third side in the greater of these triangles. Would the solution be ambiguous if the data had been  $B=30^\circ, c=150, b=75$ ?

6. If  $A=36^\circ, a=4$ , and the perpendicular from  $C$  upon  $AB$  is  $\sqrt{5}-1$ , find the other angles.

7. If the angles adjacent to the base of a triangle are  $22\frac{1}{2}^\circ$  and  $112\frac{1}{2}^\circ$ , shew that the altitude is half the base.

8. If  $a=2b$  and  $A=3B$ , find the angles and express  $c$  in terms of  $a$ .

9. The sides of a triangle are  $2x+3$ ,  $x^2+3x+3$ ,  $x^2+2x$ : shew that the greatest angle is  $120^\circ$ .

Shew that in any triangle

$$10. (b-a) \cos C + c (\cos B - \cos A) = c \sin \frac{A-B}{2} \operatorname{cosec} \frac{A+B}{2}.$$

$$11. a \sin \left( \frac{A}{2} + B \right) = (b+c) \sin \frac{A}{2}.$$

$$12. \sin \left( B + \frac{C}{2} \right) \cos \frac{C}{2} = \frac{a+b}{b+c} \cos \frac{A}{2} \cos \frac{B-C}{2}.$$

$$13. \frac{1 + \cos(A-B) \cos C}{1 + \cos(A-C) \cos B} = \frac{a^2 + b^2}{a^2 + c^2}.$$

14. If  $c^4 - 2(a^2 + b^2)c^2 + a^4 + a^2b^2 + b^4 = 0$ , prove that  $C$  is  $60^\circ$  or  $120^\circ$ .

15. If  $a, b, A$  are given, and if  $c_1, c_2$  are the values of the third side in the ambiguous case, prove that if  $c_1 > c_2$ ,

$$(1) \quad c_1 - c_2 = 2a \cos B_1.$$

$$(2) \quad \cos \frac{C_1 - C_2}{2} = \frac{b \sin A}{a}.$$

$$(3) \quad c_1^2 + c_2^2 - 2c_1c_2 \cos 2A = 4a^2 \cos^2 A.$$

$$(4) \quad \sin \frac{C_1 + C_2}{2} \sin \frac{C_1 - C_2}{2} = \cos A \cos B_1.$$

16. If  $A = 45^\circ$ , and  $c_1, c_2$  be the two values of the ambiguous side, shew that

$$\cos B_1 \cos B_2 = \frac{2c_1c_2}{c_1^2 + c_2^2}.$$

17. If  $\cos A + 2 \cos C : \cos A + 2 \cos B = \sin B : \sin C$ , prove that the triangle is either isosceles or right-angled.

18. If  $a, b, c$  are in A.P., shew that

$$\cot \frac{A}{2}, \cot \frac{B}{2}, \cot \frac{C}{2} \text{ are also in A.P.}$$

19. Shew that

$$\frac{a^2 \sin(B-C)}{\sin B + \sin C} + \frac{b^2 \sin(C-A)}{\sin C + \sin A} + \frac{c^2 \sin(A-B)}{\sin A + \sin B} = 0.$$

### MISCELLANEOUS EXAMPLES. D.

1. Prove that (1)  $\tan 2\theta \cot \theta - 1 = \sec 2\theta$ ;  
(2)  $\sin a - \cot \theta \cos a = -\operatorname{cosec} \theta \cos (a + \theta)$ .
2. If  $a = 48$ ,  $b = 35$ ,  $C = 60^\circ$ , find  $c$ .
3. If  $\cos a = \frac{8}{17}$  and  $\cos \beta = \frac{15}{17}$ , find  
 $\tan (a + \beta)$  and  $\operatorname{cosec} (a + \beta)$ .
4. If  $a = \frac{\pi}{21}$ , find the value of  $\frac{\sin 23a - \sin 7a}{\sin 2a + \sin 14a}$ .
5. Prove that  $\sin \theta (\cos 2\theta + \cos 4\theta + \cos 6\theta) = \sin 3\theta \cos 4\theta$ .
6. If  $b = \sqrt{2}$ ,  $c = \sqrt{3} + 1$ ,  $A = 45^\circ$ , solve the triangle.
7. Prove that  
(1)  $2 \sin^2 36^\circ = \sqrt{5} \sin 18^\circ$ ; (2)  $4 \sin 36^\circ \cos 18^\circ = \sqrt{5}$ .
8. Prove that  $\frac{\sin 3a}{\sin a} + \frac{\cos 3a}{\cos a} = 4 \cos 2a$ .
9. If  $b = c = 2$ ,  $a = \sqrt{6} - \sqrt{2}$ , solve the triangle.
10. Shew that  
(1)  $\cos 2a - \cot 3a \sin 2a = \tan a (\sin 2a + \cot 3a \cos 2a)$ .  
(2)  $\cos a + \cos 2a + \cos 3a = 4 \cos a \cos \frac{a}{2} \cos \frac{3a}{2} - 1$ .
11. In any triangle, prove that  
(1)  $b^2 \sin 2C + c^2 \sin 2B = 2bc \sin A$ ;  
(2)  $\frac{a^2 \sin (B - C)}{\sin A} + \frac{b^2 \sin (C - A)}{\sin B} + \frac{c^2 \sin (A - B)}{\sin C} = 0$ .
12. If  $A, B, C, D$  are the angles of a quadrilateral, prove that  
$$\frac{\tan A + \tan B + \tan C + \tan D}{\cot A + \cot B + \cot C + \cot D} = \tan A \tan B \tan C \tan D.$$
  
[Use  $\tan (A + B) = \tan (360^\circ - C - D)$ .]



## CHAPTER XIV.

### LOGARITHMS.

**151. DEFINITION.** The **logarithm** of any number to a given **base** is the index of the power to which the base must be raised in order to equal the given number. Thus if  $a^x=N$ ,  $x$  is called the logarithm of  $N$  to the base  $a$ .

*Example 1.* Since  $3^4=81$ , the logarithm of 81 to base 3 is 4.

*Example 2.* Since  $10^1=10$ ,  $10^2=100$ ,  $10^3=1000$ ,..... the natural numbers 1, 2, 3,... are respectively the logarithms of 10, 100, 1000,..... to base 10.

*Example 3.* Find the logarithm of .008 to base 25.

Let  $x$  be the required logarithm; then by definition,

$$25^x = .008 = \frac{8}{1000} = \frac{1}{125} = \frac{1}{5^3};$$

that is,  $(5^2)^x = 5^{-3}$ , or  $5^{2x} = 5^{-3}$ ;

whence, by equating indices,  $2x = -3$ , and  $x = -1.5$ .

**152.** The logarithm of  $N$  to base  $a$  is usually written  $\log_a N$ , so that the same meaning is expressed by the two equations

$$a^x = N, \quad x = \log_a N.$$

From these equations it is evident that  $a^{\log_a N} = N$ .

*Example.* Find the value of  $\log_{.01} .00001$ .

Let  $\log_{.01} .00001 = x$ ; then  $(.01)^x = .00001$ ;

$$\therefore \left(\frac{1}{10^2}\right)^x = \frac{1}{100000}, \text{ or } \frac{1}{10^{2x}} = \frac{1}{10^5}.$$

$$\therefore 2x = 5, \text{ and } x = 2.5.$$

**153.** When it is understood that a particular system of logarithms is in use, the suffix denoting the base is omitted.

Thus in arithmetical calculations in which 10 is the base, we usually write  $\log 2, \log 3, \dots$  instead of  $\log_{10} 2, \log_{10} 3, \dots$ .

Logarithms to the base 10 are known as **Common Logarithms**; this system was first introduced in 1615 by Briggs, a contemporary of Napier the inventor of Logarithms.

Before discussing the properties of common logarithms we shall prove some general propositions which are true for all logarithms independently of any particular base.

154. *The logarithm of 1 is 0.*

For  $a^0 = 1$  for all values of  $a$ ; therefore  $\log 1 = 0$ , whatever the base may be.

155. *The logarithm of the base itself is 1.*

For  $a^1 = a$ ; therefore  $\log_a a = 1$ .

156. *To find the logarithm of a product.*

Let  $MN$  be the product; let  $a$  be the base of the system, and suppose

$$x = \log_a M, \quad y = \log_a N;$$

so that

$$a^x = M, \quad a^y = N.$$

Thus the product  $MN = a^x \times a^y = a^{x+y}$ ;

whence, by definition,  $\log_a MN = x + y$

$$= \log_a M + \log_a N.$$

Similarly,  $\log_a MNP = \log_a M + \log_a N + \log_a P$ ;  
and so on for any number of factors.

*Example.*  $\log 42 = \log (2 \times 3 \times 7) = \log 2 + \log 3 + \log 7$ .

157. *To find the logarithm of a fraction.*

Let  $\frac{M}{N}$  be the fraction, and suppose

$$x = \log_a M, \quad y = \log_a N;$$

so that

$$a^x = M, \quad a^y = N.$$

Thus the fraction  $\frac{M}{N} = \frac{a^x}{a^y} = a^{x-y}$ ;

whence, by definition,  $\log_a \frac{M}{N} = x - y$

$$= \log_a M - \log_a N.$$

*Example.*  $\log(2\frac{1}{7}) = \log \frac{15}{7} = \log 15 - \log 7$   
 $= \log(3 \times 5) - \log 7 = \log 3 + \log 5 - \log 7.$

**158.** *To find the logarithm of a number raised to any power, integral or fractional.*

Let  $\log_a(M^p)$  be required, and suppose

$$x = \log_a M, \text{ so that } a^x = M;$$

then  $M^p = (a^x)^p = a^{px};$

whence, by definition,  $\log_a(M^p) = px;$

that is,  $\log_a(M^p) = p \log_a M.$

Similarly,  $\log_a(M^{\frac{1}{r}}) = \frac{1}{r} \log_a M.$

**159.** It follows from the results we have proved that

(1) the logarithm of a product is equal to the sum of the logarithms of its factors;

(2) the logarithm of a fraction is equal to the logarithm of the numerator diminished by the logarithm of the denominator;

(3) the logarithm of the  $p$ th power of a number is  $p$  times the logarithm of the number;

(4) the logarithm of the  $r$ th root of a number is  $\frac{1}{r}$  of the logarithm of the number.

Thus by the use of logarithms the operations of multiplication and division may be replaced by those of addition and subtraction; the operations of involution and evolution by those of multiplication and division.

*Example.* Express  $\log \frac{a^5 \sqrt{b}}{\sqrt[3]{c^2}}$  in terms of  $\log a$ ,  $\log b$ ,  $\log c$ .

$$\begin{aligned} \text{The expression} &= \log(a^5 \sqrt{b}) - \log \sqrt[3]{c^2} \\ &= \log a^5 + \log \sqrt{b} - \frac{2}{3} \log c \\ &= 5 \log a + \frac{1}{2} \log b - \frac{2}{3} \log c. \end{aligned}$$

160. From the equation  $10^x = N$ , it is evident that common logarithms will not in general be integral, and that they will not always be positive.

For instance  $3154 > 10^3$  and  $< 10^4$ ;  
 $\therefore \log 3154 = 3 + \text{a fraction.}$

Again,  $\cdot 06 > 10^{-2}$  and  $< 10^{-1}$ ;  
 $\therefore \log \cdot 06 = -2 + \text{a fraction.}$

161. DEFINITION. The integral part of a logarithm is called the **characteristic**, and the decimal part is called the **mantissa**.

The characteristic of the logarithm of any number to the base 10 can be found by inspection, as we shall now shew.

162. *To determine the characteristic of the logarithm of any number greater than unity.*

It is clear that a number with two digits in its integral part lies between  $10^1$  and  $10^2$ ; a number with three digits in its integral part lies between  $10^2$  and  $10^3$ ; and so on. Hence a number with  $n$  digits in its integral part lies between  $10^{n-1}$  and  $10^n$ .

Let  $N$  be a number whose integral part contains  $n$  digits; then

$$N = 10^{(n-1) + \text{a fraction}};$$

$$\therefore \log N = (n-1) + \text{a fraction.}$$

Hence the characteristic is  $n-1$ ; that is, *the characteristic of the logarithm of a number greater than unity is less by one than the number of digits in its integral part, and is positive.*

163. *To determine the characteristic of the logarithm of a decimal fraction.*

A decimal with one cipher immediately after the decimal point, such as  $\cdot 0324$ , being greater than  $\cdot 01$  and less than  $\cdot 1$ , lies between  $10^{-2}$  and  $10^{-1}$ ; a number with two ciphers after the decimal point lies between  $10^{-3}$  and  $10^{-2}$ ; and so on. Hence a decimal fraction with  $n$  ciphers immediately after the decimal point lies between  $10^{-(n+1)}$  and  $10^{-n}$ .

Let  $D$  be a decimal beginning with  $n$  ciphers; then

$$D = 10^{-(n+1) + \text{a fraction}};$$

$$\therefore \log D = -(n+1) + \text{a fraction.}$$

Hence the characteristic is  $-(n+1)$ ; that is, *the characteristic of the logarithm of a decimal fraction is greater by unity than the number of ciphers immediately after the decimal point and is negative.*

164. The logarithms to base 10 of all integers from 1 to 200000 have been found and tabulated; in most Tables they are given to seven places of decimals.

The base 10 is chosen on account of two great advantages.

(1) From the results already proved it is evident that the characteristics can be written down by inspection, so that only the mantissæ have to be registered in the Tables.

(2) The mantissæ are the same for the logarithms of all numbers which have the same significant digits; so that it is sufficient to tabulate the mantissæ of the logarithms of *integers*.

This proposition we proceed to prove.

165. Let  $N$  be any number, then since multiplying or dividing by a power of 10 merely alters the position of the decimal point without changing the sequence of figures, it follows that  $N \times 10^p$ , and  $N \div 10^q$ , where  $p$  and  $q$  are any integers, are numbers whose significant digits are the same as those of  $N$ .

$$\begin{aligned} \text{Now} \quad \log(N \times 10^p) &= \log N + p \log 10 \\ &= \log N + p \dots \dots \dots (1). \end{aligned}$$

$$\begin{aligned} \text{Again,} \quad \log(N \div 10^q) &= \log N - q \log 10 \\ &= \log N - q \dots \dots \dots (2). \end{aligned}$$

In (1) an integer is added to  $\log N$ , and in (2) an integer is subtracted from  $\log N$ ; that is, the mantissa or decimal portion of the logarithm remains unaltered.

In this and the three preceding articles the mantissæ have been supposed positive. In order to secure the advantages of Briggs' system, we arrange our work so as *always to keep the mantissa positive*, so that when the mantissa of any logarithm has been taken from the Tables the characteristic is prefixed with its appropriate sign, according to the rules already given.

166. In the case of a negative logarithm the minus sign is written *over the characteristic*, and not before it, to indicate that the characteristic alone is negative, and not the whole expression.



Thus  $\bar{4} \cdot 30103$ , the logarithm of  $\cdot 0002$ , is equivalent to  $-4 + \cdot 30103$ , and must be distinguished from  $-4 \cdot 30103$ , an expression in which both the integer and the decimal are negative. In working with negative logarithms an arithmetical artifice will sometimes be necessary in order to make the mantissa positive. For instance, a result such as  $-3 \cdot 69897$ , in which the whole expression is negative, may be transformed by subtracting 1 from the integral part and adding 1 to the decimal part. Thus

$$-3 \cdot 69897 = -4 + (1 - \cdot 69897) = \bar{4} \cdot 30103.$$

*Example 1.* Required the logarithms of  $\cdot 0002432$ .

In the Tables we find that 3859636 is the mantissa of  $\log 2432$  (the decimal point as well as the characteristic being omitted); and, by Art. 163, the characteristic of the logarithm of the given number is  $-4$ ;

$$\therefore \log \cdot 0002432 = \bar{4} \cdot 3859636.$$

*Example 2.* Find the cube root of  $\cdot 0007$ , having given

$$\log 7 = \cdot 8450980, \quad \log 887904 = 5 \cdot 9483660.$$

Let  $x$  be the required cube root; then

$$\log x = \frac{1}{3} \log (\cdot 0007) = \frac{1}{3} (\bar{4} \cdot 8450980) = \frac{1}{3} (\bar{6} + 2 \cdot 8450980);$$

that is,

$$\log x = \bar{2} \cdot 9483660;$$

but

$$\log 887904 = 5 \cdot 9483660;$$

$$\therefore x = \cdot 0887904.$$

**167.** The logarithm of 5 and its powers can easily be obtained from  $\log 2$ ; for

$$\log 5 = \log \frac{10}{2} = \log 10 - \log 2 = 1 - \log 2.$$

*Example.* Find the value of the logarithm of the reciprocal of  $324 \sqrt[5]{125}$ , having given  $\log 2 = \cdot 3010300$ ,  $\log 3 = \cdot 4771213$ .

Since  $\log \frac{1}{a} = -\log a$ , the required value

$$= -\log (324 \sqrt[5]{125}) = -\log (2^2 \times 3^4 \times 5^{\frac{3}{5}})$$

$$= -\left(2 \log 2 + 4 \log 3 + \frac{3}{5} \log 5\right)$$

$$= -2 \cdot 9299272$$

$$= \bar{3} \cdot 0700728.$$

$$2 \log 2 = \cdot 6020600$$

$$4 \log 3 = 1 \cdot 9084852$$

$$\frac{6}{10} \log 5 = \cdot 4193820$$

$$\underline{2 \cdot 9299272}$$



**EXAMPLES. XIV. a.**

1. Find the logarithms respectively  
of the numbers 1024, 81,  $\cdot 125$ ,  $\cdot 01$ ,  $\cdot 3$ , 100,  
to the bases 2,  $\sqrt{3}$ , 4,  $\cdot 001$ ,  $\cdot 1$ ,  $\cdot 01$ .

2. Find the values of

$$\log_8 16, \log_{81} 243, \log_{\cdot 01} 10, \log_{49} 343\sqrt{7}.$$

3. Find the numbers whose logarithms respectively  
to the bases 49,  $\cdot 25$ ,  $\cdot 03$ , 1,  $\cdot 64$ , 100,  $\cdot 1$ ,  
are 2,  $\frac{1}{2}$ , -2, -1,  $-\frac{1}{2}$ , 1.5, -4.

4. Find the respective characteristics  
of the logarithms of 325, 1603, 2400, 10000, 19,  
to the bases 3, 11, 7, 9, 21.

5. Write down the characteristics of the common logarithms  
of 3.26, 523.1,  $\cdot 03$ , 1.5,  $\cdot 0002$ , 3000.1,  $\cdot 1$ .

6. The mantissa of  $\log 64439$  is  $\cdot 8091488$ , write down the  
logarithms of  $\cdot 64439$ , 6443900,  $\cdot 00064439$ .

7. The logarithm of 32.5 is 1.5118834, write down the  
numbers whose logarithms are

$$\cdot 5118834, 2.5118834, \bar{4}.5118834.$$

[When required the following logarithms may be used

$$\log 2 = \cdot 3010300, \log 3 = \cdot 4771213, \log 7 = \cdot 8450980.]$$

Find the value of

- |                            |                                    |                                      |
|----------------------------|------------------------------------|--------------------------------------|
| 8. $\log 768$ .            | 9. $\log 2352$ .                   | 10. $\log 35.28$ .                   |
| 11. $\log \sqrt{6804}$ .   | 12. $\log \sqrt[5]{\cdot 00162}$ . | 13. $\log \cdot 0217$ .              |
| 14. $\log \cos 60^\circ$ . | 15. $\log \sin^3 60^\circ$ .       | 16. $\log \sqrt[3]{\sec 45^\circ}$ . |

Find the numerical value of

$$17. 2 \log \frac{15}{8} - \log \frac{25}{162} + 3 \log \frac{4}{9}.$$

18. Evaluate  $16 \log \frac{10}{9} - 4 \log \frac{25}{24} - 7 \log \frac{80}{81}$ .
19. Find the seventh root of 7,  
given  $\log 1.320469 = .1207283$ .
20. Find the cube root of .00001764,  
given  $\log 260315 = 5.4154995$ .
21. Given  $\log 3571 = 3.5527899$ , find the logarithm of  
 $3.571 \times .03571 \times \sqrt[3]{3571}$ .
22. Given  $\log 11 = 1.0413927$ , find the logarithm of  
 $(.00011)^{\frac{1}{3}} \times (1.21)^2 \times (13.31)^{\frac{4}{3}} \div 12100000$ .
23. Find the number of digits in the integral parts of  
 $\left(\frac{21}{20}\right)^{300}$  and  $\left(\frac{126}{125}\right)^{1000}$ .

24. How many positive integers have characteristic 3 when the base is 7?

168. Suppose that we have a table of logarithms of numbers to base  $a$  and require to find the logarithms to base  $b$ .

Let  $N$  be one of the numbers, then  $\log_b N$  is required.

Let  $b^y = N$ , so that  $y = \log_b N$ .

$$\therefore \log_a (b^y) = \log_a N;$$

that is,

$$y \log_a b = \log_a N;$$

$$\therefore y = \frac{1}{\log_a b} \times \log_a N,$$

or

$$\log_b N = \frac{1}{\log_a b} \times \log_a N \dots\dots\dots(1).$$

Now since  $N$  and  $b$  are given,  $\log_a N$  and  $\log_a b$  are known from the Tables, and thus  $\log_b N$  may be found.

Hence it appears that to transform logarithms from base  $a$  to base  $b$  we have only to multiply them all by  $\frac{1}{\log_a b}$ ; this is a constant quantity and is given by the Tables; it is known as the *modulus*.

If in equation (1) we put  $a$  for  $N$ , we obtain

$$\log_b a = \frac{1}{\log_a b} \times \log_a a = \frac{1}{\log_a b};$$

$$\therefore \log_b a \times \log_a b = 1.$$

**169.** The following examples further illustrate the great use of logarithms in arithmetical work.

*Example 1.* Given  $\log 2 = .3010300$  and  $\log 4844544 = 6.6852530$ , find the value of  $(6.4)^{\frac{1}{10}} \times (\sqrt[4]{.256})^3 \div \sqrt{80}$ .

Let  $x$  be the value of the expression; then

$$\begin{aligned}\log x &= \frac{1}{10} \log \frac{64}{10} + \frac{3}{4} \log \frac{256}{1000} - \frac{1}{2} \log 80 \\ &= \frac{1}{10} (\log 2^6 - 1) + \frac{3}{4} (\log 2^8 - 3) - \frac{1}{2} (\log 2^3 + 1) \\ &= \left( \frac{6}{10} + 6 - \frac{3}{2} \right) \log 2 - \left( \frac{1}{10} + \frac{9}{4} + \frac{1}{2} \right) \\ &= \left( 5 + \frac{1}{10} \right) \log 2 - 2\frac{17}{20} \\ &= 1.5051500 + .0301030 - 2.85.\end{aligned}$$

Thus  $\log x = \overline{2}.6852530$ .

But  $\log 4844544 = 6.6852530$ ,  
 $\therefore x = .04844544$ .

*Example 2.* Find how many ciphers there are between the decimal point and the first significant digit in  $(.0504)^{10}$ ; having given

$$\log 2 = .301, \log 3 = .477, \log 7 = .845.$$

Denote the expression by  $E$ ; then

$$\begin{aligned}\log E &= 10 \log \frac{504}{10000} \\ &= 10 (\log 504 - 4) \\ &= 10 \{ \log (2^3 \times 3^2 \times 7) - 4 \} \\ &= 10 \{ 3 \log 2 + 2 \log 3 + \log 7 - 4 \} \\ &= 10 (2.702 - 4) = 10 (\overline{2}.702) \\ &= \overline{20} + 7.02 = \overline{13}.02.\end{aligned}$$

$$\begin{array}{r} 3 \log 2 = .903 \\ 2 \log 3 = .954 \\ \log 7 = .845 \\ \hline 2.702 \end{array}$$

Thus the number of ciphers is 12. [Art. 163.]

### Exponential equations.

170. If in an equation the unknown quantity appears as an exponent, the solution may be effected by the help of logarithms.

*Example 1.* Solve the equation  $8^{5-3x} = 12^{4-2x}$ , having given

$$\log 2 = \cdot 30103, \text{ and } \log 3 = \cdot 47712.$$

From the given equation, by taking logarithms, we have

$$(5 - 3x) \log 8 = (4 - 2x) \log 12;$$

$$\therefore 3(5 - 3x) \log 2$$

$$= (4 - 2x)(2 \log 2 + \log 3);$$

$$\therefore 15 \log 2 - 8 \log 2 - 4 \log 3$$

$$= x(9 \log 2 - 4 \log 2 - 2 \log 3);$$

$$\therefore x = \frac{7 \log 2 - 4 \log 3}{5 \log 2 - 2 \log 3} = \frac{\cdot 19873}{\cdot 55091}.$$

$$7 \log 2 = 2 \cdot 10721$$

$$4 \log 3 = 1 \cdot 90848$$

$$\hline \cdot 19873$$

$$5 \log 2 = 1 \cdot 50515$$

$$2 \log 3 = \cdot 95424$$

$$\hline \cdot 55091$$

$$55091 \mid 198730 \quad (36$$

$$165273$$

$$\hline 334570$$

Thus  $x = \cdot 36$  nearly.

*Example 2.* Given  $\log 2 = \cdot 30103$ , solve the simultaneous equations

$$2^x \cdot 5^y = 1, \quad 5^{x+1} \cdot 2^y = 2.$$

Take logarithms of the given equations;

$$\therefore x \log 2 + y \log 5 = 0, \quad (x + 1) \log 5 + y \log 2 = \log 2.$$

For shortness, put  $\log 2 = a$ ,  $\log 5 = b$ .

$$\text{Thus} \quad ax + by = 0,$$

$$\text{and} \quad b(x + 1) + ay = a, \text{ or } bx + ay = a - b.$$

By eliminating  $y$ ,  $x(a^2 - b^2) = -b(a - b)$ ,

$$\therefore x = -\frac{b}{a+b} = -\frac{\log 5}{\log 2 + \log 5} = -\frac{\log 5}{\log 10} = -\log 5 = -\cdot 69897.$$

$$\text{And} \quad y = -\frac{ax}{b} = \frac{a}{b} \log 5 = a = \log 2 = \cdot 30103.$$

**EXAMPLES. XIV. b.**

[When required the values of  $\log 2$ ,  $\log 3$ ,  $\log 7$  given on p. 145 may be used.]

Find the value of

1.  $\left(\frac{147 \times 375}{126 \times 16}\right)^{\frac{2}{3}}$ , given  $\log 9.076226 = .9579053$ .

2.  $\sqrt[3]{378} \times \sqrt{108} \div (\sqrt[6]{1008} \times \sqrt[3]{486})$ ,  
given  $\log 301824 = 5.4797536$ .

3.  $(1080)^{\frac{1}{2}} \times (.24)^{\frac{5}{3}} \times 810$ ,  
given  $\log 2467266 = 6.3922160$ .

Calculate to two decimal places the values of

4.  $\log_{20} 800$ .                      5.  $\log_3 49$ .                      6.  $\log_{125} 4000$ .

7. Find how many ciphers there are before the first significant digits in

$$(.00378)^{\frac{40}{3}} \text{ and } (.0259)^{50}.$$

8. To what base is 3 the logarithm of 11000?  
given  $\log 11 = 1.0413927$  and  $\log 222398 = 5.3471309$ .

Solve to two decimal places the equations:

9.  $2^{x-1} = 5$ .                      10.  $3^{x-4} = 7$ .                      11.  $5^{1-x} = 6^{x-3}$ .

12.  $5^x = 2^{-y}$  and  $5^{2+y} = 2^{2-x}$ .

13.  $2^x = 3^y$  and  $2^{y+1} = 3^{x-1}$ .

14. Given  $\log 28 = a$ ,  $\log 21 = b$ ,  $\log 25 = c$ , find  $\log 27$  and  $\log 224$  in terms of  $a$ ,  $b$ ,  $c$ .

15. Given  $\log 242 = a$ ,  $\log 80 = b$ ,  $\log 45 = c$ , find  $\log 36$  and  $\log 66$  in terms of  $a$ ,  $b$ ,  $c$ .

### MISCELLANEOUS EXAMPLES. E.

1. Prove that

$$\cos(30^\circ + A) \cos(30^\circ - A) - \cos(60^\circ + A) \cos(60^\circ - A) = \frac{1}{2}.$$

2. If  $A + B + C = 180^\circ$ , shew that

$$\frac{\sin 2A + \sin 2B + \sin 2C}{\sin A + \sin B + \sin C} = 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$$

3. If  $a=2$ ,  $c=\sqrt{2}$ ,  $B=15^\circ$ , solve the triangle.

4. Shew that  $\cos a + \tan \frac{a}{2} \sin a = \cot \frac{a}{2} \sin a - \cos a$ .

5. If  $b \cos A = a \cos B$ , shew that the triangle is isosceles.

6. Prove that

$$(1) \quad \sin \theta (\sin 3\theta + \sin 5\theta + \sin 7\theta + \sin 9\theta) = \sin 6\theta \sin 4\theta;$$

$$(2) \quad \frac{\sin a + \sin 3a + \sin 5a + \sin 7a}{\cos a + \cos 3a + \cos 5a + \cos 7a} = \tan 4a.$$

7. Shew that  $\frac{\cos 3a}{\sin a} + \frac{\sin 3a}{\cos a} = 2 \cot 2a$ .

8. If  $b=a(\sqrt{3}-1)$ ,  $C=30^\circ$ , find  $A$  and  $B$ .

9. Shew that  $\tan 4a = \frac{4 \tan a - 4 \tan^3 a}{1 - 6 \tan^2 a + \tan^4 a}$ .

10. In a triangle, shew that

$$(1) \quad a^2 \cos 2B + b^2 \cos 2A = a^2 + b^2 - 4ab \sin A \sin B;$$

$$(2) \quad 4 \left( bc \cos^2 \frac{A}{2} + ca \cos^2 \frac{B}{2} + ab \cos^2 \frac{C}{2} \right) = (a+b+c)^2.$$

11. If  $a^4 + b^4 + c^4 = 2c^2(a^2 + b^2)$ , prove that  $C=45^\circ$  or  $135^\circ$ .

[Solve as a quadratic in  $c^2$ .]

12. If in a triangle  $\cos 3A + \cos 3B + \cos 3C = 1$ , shew that one angle must be  $120^\circ$ .



## CHAPTER XV.

### THE USE OF LOGARITHMIC TABLES.

171. We shall now explain the use of logarithmic Tables to which reference has been made in the previous chapter.

In a book of Tables there will usually be found the *mantissæ* of the logarithms of all *integers* from 1 to 100000; the *characteristics* can be written down by inspection and are therefore omitted. [Art. 162.]

The logarithm of any number consisting of not more than 5 significant digits can be obtained directly from these Tables. For instance, suppose the logarithm of 336·34 is required. Opposite to 33634 we find the figures 5267785; this, with the decimal point prefixed, is the mantissa for the logarithms of all numbers whose significant digits are the same as 33634. We have therefore only to prefix the characteristic 2, and we obtain

$$\log 336\cdot34 = 2\cdot5267785.$$

Similarly,  $\log 33634 = 4\cdot5267785,$   
and  $\log \cdot0033634 = \bar{3}\cdot5267785.$

172. Suppose now that we required  $\log 33634\cdot392.$

Since this number contains more than 5 significant digits it cannot be obtained directly from the tables; but it lies between the two consecutive numbers 33634 and 33635, and therefore its logarithm lies between the logarithms of these two numbers. If we pass from 33634 to 33635, making an increase of 1 in the number, the corresponding increase in the logarithm as obtained from the tables is ·0000129. If now we pass from 33634 to 33634·392, making an increase of ·392 in the number, the increase in the logarithm will be  $\cdot392 \times \cdot0000129$ , provided that the increase in the logarithm is proportional to the increase in the number.

Now it can be proved that *when the increase made is small in comparison with the number, the increase in the logarithm is very nearly proportional to the increase in the number.*

This principle is known as the **Rule of Proportional Parts.**

The application of this rule will be illustrated in the examples which follow.

**173.** In order to make the explanations more intelligible we give here an Extract from Chambers' *Mathematical Tables.*

No.	0	1	2	3	4	5	6	7	8	9	Diff.
3361	526 4685	4814	4944	5073	5202	5331	5460	5590	5719	5848	
62	5977	6106	6235	6365	6494	6623	6752	6881	7010	7140	129
63	7269	7398	7527	7656	7785	7914	8043	8173	8302	8431	1 13
64	8560	8689	8818	8947	9076	9205	9334	9463	9593	9722	2 26
65	9851	9980	0109	0238	0367	0496	0625	0754	0883	1012	3 39
											4 52
66	527 1141	1270	1399	1528	1657	1786	1915	2044	2173	2302	5 65
67	2431	2560	2689	2818	2947	3076	3205	3334	3463	3592	6 77
68	3721	3850	3979	4108	4237	4366	4494	4623	4752	4881	7 96
69	5010	5139	5268	5397	5526	5655	5783	5912	6041	6170	8 103
70	6299	6428	6557	6686	6814	6943	7072	7201	7330	7459	9 116

**174.** Suppose that  $\log 33635$  is required.

In the third horizontal line we have the logarithms of numbers beginning with 3363. As the next digit is 5 we choose from this line the mantissa which stands under the column 5. We have now only to prefix the characteristic and we obtain  $\log 33635 = 4.5267914$ .

Similarly,  $\log 33651 = 4.5269980$ ,  
and  $\log 33652 = 4.5270109$ ,

the transition in the mantissæ from 526... to 527... being shewn by the bar drawn over 0109. This bar is repeated over each of the subsequent logarithms as far as the end of the line, and in the next line the mantissæ begin with 527.

*Example.* Find  $\log 33634.392$ .

From the Tables,  $\log 33635 = 4.5267914$   
 $\log 33634 = 4.5267785$   
difference for 1 =  $\overline{.0000129}$

Now by the Rule of Proportional Parts,  
 $\log 33634.392$  will be greater than  $\log 33634$   
by  $.392$  times the difference for 1; hence to  
7 places of decimals, we have

$\overline{.0000129}$	
$\overline{.392}$	
	$\overline{258}$
	11 61
	38 7
$\overline{.0000050}$	$\overline{568}$

$\log 33634 = 4.5267785$   
proportional difference for  $.392 = .0000051$   
 $\therefore \log 33634.392 = 4.5267836$

In practice, the difference for 1 is usually quoted without the ciphers; if therefore we *treat the difference 129 as a whole number*, on multiplying by  $.392$  we obtain the product 50.568, and we take the digits given by its integral part (51 approximately) as the proportional increase for  $.392$ .

**175.** The method of calculating the proportional difference for  $.392$  which we have explained is that which must be adopted when we have nothing given but the logarithms of two consecutive numbers between which lies the number whose logarithm we are seeking.

But when the Tables are used the calculation is facilitated by means of the proportional differences standing in the column to the right. This gives the differences for *tenths* of unity.

The difference for  $.392$  is obtained as follows.

$$.392 \times 129 = \left( \frac{3}{10} + \frac{9}{100} + \frac{2}{1000} \right) \times 129 = 39 + 11.6 + .26 = 50.86.$$

The difference for 9 quoted in the margin (really 9 *tenths*) is 116, and therefore the difference for 9 hundredths is 11.6; and similarly the difference for 2 thousandths is .26.

In practical work, the following arrangement is adopted.

$$\begin{array}{rcl}
 & \log 33634 & = 4.5267785 \\
 \text{add for} & \begin{array}{r} 3 \\ 9 \\ 2 \end{array} & \begin{array}{r} 39 \\ 116 \\ 26 \end{array} \\
 \hline
 \therefore \log 33634.392 & = & 4.5267836
 \end{array}$$

176. The following example is solved more concisely as a model for the student. In the column on the left we work from the data of the question; in the column on the right we obtain the logarithm by the use of the Tables independently of the two given logarithms.

*Example.* Find  $\log 33.656208$ , having given

$$\log 33656 = 4.5270625 \quad \text{and} \quad \log 33657 = 4.5270754.$$

$  \begin{array}{rcl}  \log 33.657 & = & 1.5270754 \\  \log 33.656 & = & 1.5270625 \\  \text{diff. for } .001 & = & \begin{array}{r} 129 \\ 208 \end{array} \\  & & \begin{array}{r} 1032 \\ 2580 \end{array} \\  \text{diff. for } .000208 & = & \begin{array}{r} 26832 \end{array} \\  \log 33.656 & = & 1.5270625 \\  \log 33.656208 & = & 1.5270652  \end{array}  $	<p>From the Tables, we have</p> $  \begin{array}{rcl}  \log 33.656 & = & 1.5270625 \\  \text{add for} & \begin{array}{r} 2 \\ 0 \\ 8 \end{array} & \begin{array}{r} 26 \\ 0 \\ 103 \end{array} \\  \hline  \log 33.656208 & = & 1.5270652  \end{array}  $
---	---

177. The Rule of Proportional Parts also enables us to find the number corresponding to a given logarithm.

*Example 1.* Find the number whose logarithm is  $\bar{2}.5274023$ , having given  $\log 3.3683 = .5274108$  and  $\log 3.3682 = .5273979$ .

Let  $x$  be the required number; then

$  \begin{array}{rcl}  \log x & = & \bar{2}.5274023 \\  \log .033682 & = & \bar{2}.5273979 \\  \text{diff.} & = & 44  \end{array}  $	$  \begin{array}{rcl}  \log .033683 & = & \bar{2}.5274108 \\  \log .033682 & = & \bar{2}.5273979 \\  \text{diff. for } .000001 & = & 129  \end{array}  $
--	--

hence  $x$  lies between  $.033682$  and  $.033683$ ,  
 and is greater than  $.033682$  by  $\frac{44}{129} \times .000001$ ,  
 that is by  $.00000034$ .

$$\therefore x = .03368234.$$

$$\begin{array}{r}
 129 \ ) \ 440 \ ( \ 34 \\
 \underline{387} \\
 530 \\
 \underline{516}
 \end{array}$$

In working from the Tables, we proceed as follows.

$$\begin{array}{rcl} \log x & = & \bar{2}.5274023 \\ \log .033682 & = & \bar{2}.5273979 \\ & & \hline & & 44 \\ & 3 & 39 \\ & & \hline & & 50 \\ & 4 & 52 \\ & & \hline \end{array}$$

$$\therefore x = .03368234.$$

We are saved the trouble of the division, as the multiples of 129 which occur during the work are given in the approximate forms 39 and 52 in the difference column opposite to the numbers 3 and 4.

*Example 2.* Find the fifth root of .0025612, having given  
 $\log 2.5612 = .4084435$ ,  $\log 3.0317 = .4816862$ ,  $\log 3.0318 = .4817005$ .

Let  $x = (.0025612)^{\frac{1}{5}}$ ; then

$$\begin{aligned} \log x &= \frac{1}{5} \log (.0025612) = \frac{1}{5} (\bar{3}.4084435) = \frac{1}{5} (\bar{5} + 2.4084435); \\ &= \bar{1}.4816887. \end{aligned}$$

$$\begin{array}{rcl} \log x & = & \bar{1}.4816887 \\ \log .30317 & = & \bar{1}.4816862 \\ \text{diff.} & = & \frac{25}{25} \end{array} \qquad \begin{array}{rcl} \log .30318 & = & \bar{1}.4817005 \\ \log .30317 & = & \bar{1}.4816862 \\ \text{diff. for } .00001 & = & \frac{143}{143} \end{array}$$

$$\therefore \text{proportional increase} = \frac{25}{143} \times .00001 = .00000175.$$

$$\text{Thus} \qquad x = .30317175.$$

$$\begin{array}{r} 143 \ ) \ 250 \ (175 \\ \underline{143} \\ 1070 \\ \underline{1001} \\ 690 \\ \underline{715} \end{array}$$

### EXAMPLES. XV. a.

- Find the value of  $\log 4951634$ , given that  
 $\log 49516 = 4.6947456$ ,  $\log 49517 = 4.6947543$ .
- Find  $\log 3.4713026$ , having given that  
 $\log 347.13 = 2.5404921$ ,  $\log 34714 = 4.5405047$ .
- Find  $\log 2849614$ , having given that  
 $\log 2.8496 = .4547839$ ,  $\log 2.8497 = .4547991$ .



4. Find  $\log 57.63325$ , having given that  
 $\log 576.33 = 2.7606712$ ,  $\log 5763.4 = 3.7606788$ .
5. Given  $\log 60814 = 4.7840036$ , diff. for 1 = 72, find  
 $\log 6081465$ .
6. Find the number whose logarithm is  $4.7461735$ , given  
 $\log 55740 = 4.7461670$ ,  $\log 55741 = 4.7461748$ .
7. Find the number whose logarithm is  $2.8283676$ , given  
 $\log 6.7354 = .8283634$ ,  $\log 67355 = 4.8283698$ .
8. Find the number whose logarithm is  $\bar{2}.0288435$ , given  
 $\log 1068.6 = 3.0288152$ ,  $\log 1.0687 = .0288558$ .
9. Find the number whose logarithm is  $\bar{3}.9184377$ , given  
 $\log 8.2877 = .9184340$ ,  $\log 8287.8 = 3.9184392$ .
10. Given  $\log 253.19 = 2.4034465$ , diff. for 1 = 172, find the  
number whose logarithm is  $\bar{1}.4034508$ .
11. Given  $\log 2.0313 = .3077741$ ,  $\log 2.0314 = .3077954$ ,  
and  $\log 1.4271 = .1544544$ ,  
find the seventh root of  $142.71$ .
12. Find the eighth root of  $13.89492$ , given  
 $\log 13894 = 4.1428273$ ,  $\log 138.95 = 2.1428586$ .
13. Find the value of  $\sqrt[14]{242447}$ , given  
 $\log 2.4244 = .3846043$ , diff. for 1 = 179.
14. Find the twentieth root of  $2069138$ , given  
 $\log 20691 = 4.3157815$ , diff. for 1 = 210.

### Tables of Natural and Logarithmic Functions.

178. Tables have been constructed giving the values of the trigonometrical functions of all angles between  $0^\circ$  and  $90^\circ$  at intervals of  $10''$ . These are called the **Tables of natural sines, cosines, tangents,...** In the smaller Tables, such as Chambers', the interval is  $1'$ .

The logarithms of the functions have also been calculated. Since many of the trigonometrical functions are less than unity



their logarithms are negative, and as the characteristics are not always evident on inspection they cannot be omitted. To avoid the inconvenience of printing the bars over the characteristics, the logarithms are all increased by 10 and are then registered under the name of **tabular logarithmic sines, cosines,...**

The notation used is  $L \cos A$ ,  $L \tan \theta$ ; thus

$$L \sin A \doteq \log \sin A + 10.$$

For instance,

$$\begin{aligned} L \sin 45^\circ &= 10 + \log \sin 45^\circ = 10 + \log \frac{1}{\sqrt{2}} \\ &= 10 - \frac{1}{2} \log 2 = 9.8494850. \end{aligned}$$

**179.** With certain exceptions that need not be here noticed, the rule of proportional parts holds for the natural sines, cosines,... of all angles, and also for their logarithmic sines, cosines,.... In applying this rule it must be remembered that as the angle increases from  $0^\circ$  to  $90^\circ$  the functions sine, tangent, secant increase, while the co-functions cosine, cotangent, cosecant decrease.

*Example 1.* Find the value of  $\sin 29^\circ 37' 42''$ .

From the Tables,	$\sin 29^\circ 38' = .4944476$ $\sin 29^\circ 37' = .4941948$ <div style="border-top: 1px solid black; display: flex; justify-content: space-between; padding-top: 2px;"> <span>diff. for <math>60'' =</math></span> <span>2528</span> </div>
$\therefore$ prop <sup>l</sup> increase for $42'' = \frac{42}{60} \times 2528 =$	1770
	$\sin 29^\circ 37' = .4941948$ $\therefore \sin 29^\circ 37' 42'' = .4943718.$

*Example 2.* Find the angle whose cosine is .7280843.

Let  $A$  be the required angle; then from the Tables,

$\cos 43^\circ 16' = .7281716$ $\cos 43^\circ 17' = .7279722$ <div style="border-top: 1px solid black; display: flex; justify-content: space-between; padding-top: 2px;"> <span>diff. for <math>60'' =</math></span> <span>1994</span> </div>	$\cos 43^\circ 16' = .7281716$ $\cos A = .7280843$ <div style="border-top: 1px solid black; display: flex; justify-content: space-between; padding-top: 2px;"> <span>prop<sup>l</sup> part</span> <span>873</span> </div>
	<div style="border-top: 1px solid black; display: flex; justify-content: space-between; padding-top: 2px;"> <span></span> <span>60</span> </div>

But  $\cos A$  is less than  $\cos 43^\circ 16'$ ;  
hence  $A$  must be greater than  $43^\circ 16'$   
by  $\frac{873}{1994} \times 60''$ , that is by  $26''$  nearly.

Thus the angle is  $43^\circ 16' 26''$ .

1994 )	52380 ( 26
	3988
	12500
	11964

180. In order to illustrate the use of the tabular logarithmic functions we give the following extract from the table of logarithmic sines, cosines,... in Chambers' *Mathematical Tables*.

27 Deg.

,	Sine	Diff.	Cosec.	Secant	D.	Cosine	,
0	9.6570468	2478	10.3429532	10.0501191	644	9.9498809	60
1	9.6572946	2477	10.3427054	10.0501835	644	9.9498165	59
2	9.6575423	2475	10.3424577	10.0502479	645	9.9497521	58
3	9.6577898	2473	10.3422102	10.0503124	646	9.9496876	57
4	9.6580371		10.3419629	10.0503770		9.9496230	56
56	9.6706576	2382	10.3293424	10.0537968	670	9.9462032	4
57	9.6708958	2380	10.3291042	10.0538638	670	9.9461362	3
58	9.6711338	2378	10.3288662	10.0539308	671	9.9460692	2
59	9.6713716	2377	10.3286284	10.0539979	672	9.9460021	1
60	9.6716093		10.3283907	10.0540651		9.9459349	0
'	Cosine	Diff.	Secant	Cosec.	D.	Sine	'

62 Deg.

181. We have quoted here the logarithmic sines, cosecants, secants, and cosines of the angles differing by 1' between 27° 0' and 27° 4', and also between 27° 56' and 27° 60'. The same extract gives the logarithmic functions of the complements of these angles, namely those between 62° 0' and 62° 4', and those between 62° 56' and 62° 60'.

The column of minutes for 27° is given on the left and increases downwards, the column for 62° is on the right and increases upwards.

The names of the functions printed at the top refer to the angle 27°, the names printed at the foot refer to the angle 62°. Thus

$$\begin{aligned}
 L \cos 27^\circ 3' &= 9.9496876, & L \operatorname{cosec} 27^\circ 58' &= 10.3288662, \\
 L \sin 62^\circ 2' &= 9.9460692, & L \cos 62^\circ 59' &= 9.6572946.
 \end{aligned}$$

The first *difference column* gives the differences in the logarithms of the sines and cosecants, the second *difference column* gives the differences in the logarithms of the cosines and secants, each difference corresponding to a difference of 1' in the angle.

*Example 1.* Find  $L \cos 62^\circ 57' 12''$ .

$$\begin{array}{rcl} \text{From the Tables,} & L \cos 62^\circ 57' & = 9.6577898 \\ & L \cos 62^\circ 58' & = 9.6575423 \\ & \text{diff. for } 60'' & \underline{2475} \end{array}$$

$$\therefore \text{proportional decrease for } 12'' = \frac{12}{60} \times 2475 = 495.$$

$$\begin{array}{rcl} & L \cos 60 57' & = 9.6577898 \\ & \text{Subtract for } 12'' & \underline{495} \\ \therefore L \cos 62^\circ 57' 12'' & = & 9.6577403 \end{array}$$

*Example 2.* Given  $L \sec 27^\circ 39' = 10.0526648$ , diff. for  $10'' = 110$ , find  $A$  when  $L \sec A = 10.0527253$ .

$$\begin{array}{rcl} L \sec A & = & 10.0527253 \\ L \sec 27^\circ 39' & = & 10.0526648 \\ & \text{diff.} & \underline{605} \end{array}$$

$$\therefore \text{proportional increase} = \frac{605}{110} \times 10'' = 55''.$$

Thus

$$A = 27^\circ 39' 55''.$$

### EXAMPLES. XV. b.

1. Find  $\sin 38^\circ 3' 35''$ , having given that  
 $\sin 38^\circ 4' = .6165780$ ,  $\sin 38^\circ 3' = .6163489$ .
2. Find  $\tan 38^\circ 24' 37.5''$ , having given that  
 $\tan 38^\circ 25' = .7930640$ ,  $\tan 38^\circ 24' = .7925902$ .
3. Find  $\operatorname{cosec} 55^\circ 21' 28''$ , having given that  
 $\operatorname{cosec} 55^\circ 22' = 1.2153535$ ,  $\operatorname{cosec} 55^\circ 21' = 1.2155978$ .
4. Find the angle whose secant is  $2.1809460$ , given  
 $\sec 62^\circ 43' = 2.1815435$ ,  $\sec 62^\circ 42' = 2.1803139$ .
5. Find the angle whose cosine is  $.8600931$ , given  
 $\cos 30^\circ 41' = .8600007$ ,  $\cos 30^\circ 40' = .8601491$ .
6. Find the angle whose cotangent is  $.8766003$ , given  
 $\cot 48^\circ 46' = .8764620$ ,  $\cot 48^\circ 45' = .8769765$ .
7. Find  $L \sin 44^\circ 17' 33''$ , given  
 $L \sin 44^\circ 18' = 9.8441137$ ,  $L \sin 44^\circ 17' = 9.8439842$ .

8. Find
- $L \cot 36^\circ 26' 16''$
- , given

$$L \cot 36^\circ 27' = 10.1315840, \quad L \cot 36^\circ 26' = 10.1318483.$$

9. Find
- $L \cos 55^\circ 30' 24''$
- , given

$$L \cos 55^\circ 31' = 9.7529442, \quad L \cos 55^\circ 30' = 9.7531280.$$

10. Find the angle whose tabular logarithmic sine is 9.8440018, using the data of example 7.

11. Find the angle whose tabular logarithmic cosine is 9.7530075, using the data of example 9.

12. Given
- $L \tan 24^\circ 50' = 9.6653662$
- , diff. for
- $1' = 3313$
- , find
- 
- $L \tan 24^\circ 50' 52.5''$
- .

13. Given
- $L \operatorname{cosec} 40^\circ 5' = 10.1911808$
- , diff. for
- $1' = 1502$
- , find
- 
- $L \operatorname{cosec} 40^\circ 4' 17.5''$
- .

182. Considerable practice in the use of logarithmic Tables will be required before the quickness and accuracy necessary in all practical calculations can be attained. Experience shews that mistakes frequently arise from incorrect quotation from the Tables, and from clumsy arrangement. The student is reminded that care in taking out the logarithms from the Tables is of the first importance, and that in the course of the work he should learn to leave out all needless steps, making his solutions as concise as possible consistent with accuracy.

**Example 1.** Divide 6.6425693 by .3873007.

From the Tables,

$$\begin{array}{r} \log 6.6425 \quad = \quad .8223316 \\ \quad \quad \quad 6 \quad \quad \quad 40 \\ \quad \quad \quad 9 \quad \quad \quad 59 \\ \quad \quad \quad 3 \quad \quad \quad 20 \\ \hline \log 6.6425693 = .8223362 \\ \log .3873007 = \bar{1}.5880483 \end{array}$$

$$\begin{array}{r} \log .38730 \quad = \quad \bar{1}.5880475 \\ \quad \quad \quad 0 \quad \quad \quad 0 \\ \quad \quad \quad 7 \quad \quad \quad 78 \\ \hline \log .3873007 = \bar{1}.5880483 \end{array}$$

By subtraction, we obtain  $1.2342879$   
 From the Tables,  $\log 17.150 \quad = 1.2342641$

$$\begin{array}{r} 238 \\ 9 \quad 229 \\ \hline 90 \\ 3 \quad 76 \\ \hline \end{array}$$

Thus the quotient is 17.15093.

**Example 2.** The hypotenuse of a right-angled triangle is 3·141024 and one side is 2·593167; find the other side.

Let  $c$  be the hypotenuse,  $a$  the given side, and  $x$  the side required; then

$$\begin{aligned} x^2 &= c^2 - a^2 = (c + a)(c - a); \\ \therefore 2 \log x &= \log(c + a) + \log(c - a). \end{aligned}$$

$$\begin{aligned} c &= 3\cdot141024 \\ a &= 2\cdot593167 \\ c+a &= 5\cdot734191 \\ c-a &= \cdot547853 \end{aligned}$$

$$\begin{array}{rcl} \text{From the Tables, } \log 5\cdot7341 & = & \cdot7584653 \\ & 9 & 68 \\ & 1 & \\ \log \cdot54785 & = & \bar{1}\cdot7386617 \\ & 7 & 55 \\ \text{By addition,} & & \hline & & \cdot4971394 \end{array}$$

Dividing by 2, we have

$$\begin{array}{rcl} \log x & = & \cdot2485697 \\ \log 1\cdot7724 & = & \cdot2485617 \\ & & \hline & & 80 \\ & 3 & 74 \\ & & \hline & & 60 \\ & 2 & 49 \\ & & \hline \end{array}$$

Thus the required side is 1·772432.

### EXAMPLES. XV. c.

[In this exercise the logarithms are to be taken from the Tables.]

1. Multiply 300·2618 by ·0078915194.
2. Find the product of 235·6783 and 357·8438.
3. Find the continued product of  
153·2419, 2·8632503, and ·07583646.
4. Divide 1·0304051 by 27·093524.
5. Divide 357·8364 by ·00318973.
6. Find  $x$  from the equation  
 $\cdot0178345x = 21\cdot85632$ .
7. Find the value of  
 $3\cdot78956 \times \cdot0536872 \div \cdot0072916$ .

8. Find the cube of  $\cdot 83410039$ .
9. Find the fifth root of  $15063\cdot 018$ .
10. Evaluate  $\sqrt[5]{384\cdot 731}$  and  $\sqrt[13]{15\cdot 7324}$ .
11. Find the product of the square root of  $1034\cdot 3963$  and the cube root of  $353246$ .
12. Subtract the square of  $\cdot 7503269$  from the square of  $1\cdot 035627$ .
13. Find the value of

$$\frac{(34\cdot 7326)^{\frac{3}{5}} \times \sqrt[5]{2\cdot 53894}}{\sqrt[5]{4\cdot 39682}}.$$

**Example 3.** Find a third proportional to the cube of  $\cdot 3172564$  and the cube root of  $23\cdot 32873$ .

Let  $x$  be the required third proportional; then

$$(\cdot 3172564)^3 : (23\cdot 32873)^{\frac{1}{3}} = (23\cdot 32873)^{\frac{1}{3}} : x;$$

whence

$$x = (23\cdot 32873)^{\frac{2}{3}} \div (\cdot 3172564)^3;$$

$$\therefore \log x = \frac{2}{3} \log 23\cdot 32873 - 3 \log \cdot 3172564.$$

From the Tables,

$$\begin{array}{r} \log \cdot 31725 = \overline{1}\cdot 5014016 \\ \phantom{\log \cdot 31725 = } \begin{array}{r} 6 \phantom{00} 82 \\ 4 \phantom{00} 55 \end{array} \\ \hline \phantom{\log \cdot 31725 = } \overline{1}\cdot 5014103 \overline{5} \\ \phantom{\log \cdot 31725 = } \phantom{\overline{1}\cdot 5014103} 3 \\ \hline \phantom{\log \cdot 31725 = } 2\cdot 5042311 \end{array}$$

$$\begin{array}{r} \log 23\cdot 328 = 1\cdot 3678775 \\ \phantom{\log 23\cdot 328 = } \begin{array}{r} 7 \phantom{00} 130 \\ 3 \phantom{00} 56 \end{array} \\ \hline \phantom{\log 23\cdot 328 = } 1\cdot 3678910 \overline{6} \\ \phantom{\log 23\cdot 328 = } \phantom{1\cdot 3678910} 2 \\ \hline \phantom{\log 23\cdot 328 = } 3 \overline{2\cdot 7357821} \\ \phantom{\log 23\cdot 328 = } \phantom{3 \overline{2\cdot 7357821}} \cdot 9119274 \\ \hline \phantom{\log 23\cdot 328 = } 2\cdot 5042311 \end{array}$$

$$\begin{array}{rcl} \text{By subtraction, } \log x & = & 2\cdot 4076963 \\ \log 255\cdot 67 & = & 2\cdot 4076798 \\ & & \hline & & 165 \\ & 9 & 153 \\ & & \hline & & 120 \\ & 7 & 119 \\ & & \hline \end{array}$$

Thus the third proportional is  $255\cdot 6797$ .



14. Find a mean proportional between  
 $\cdot 0037258169$  and  $\cdot 56301078$ .
15. Find a third proportional to the square of  $\cdot 43607528$  and the square root of  $\cdot 03751786$ .
16. Find a fourth proportional to  
 $56712\cdot 43$ ,  $29\cdot 302564$ ,  $\cdot 33025107$ .
17. Find the geometric mean between  
 $(\cdot 035689)^{\frac{2}{5}}$  and  $(2\cdot 879432)^{\frac{3}{7}}$ .
18. Find a fourth proportional to  
 $\sqrt[3]{32\cdot 7812}$ ,  $\sqrt[5]{357\cdot 814}$ ,  $\sqrt[4]{7836\cdot 43}$ .
19. Find the value of  
 $\sin 27^{\circ} 13' 12'' \times \cos 46^{\circ} 2' 15''$ .
20. Find the value of  
 $\cot 97^{\circ} 14' 16'' \times \sec 112^{\circ} 13' 5''$ .
21. Evaluate  
 $\sin 20^{\circ} 13' 20'' \times \cot 47^{\circ} 53' 15'' \times \sec 42^{\circ} 15' 30''$ .
22. Find the value of  $ab \sin C$ , when  
 $a=324\cdot 1368$ ,  $b=417\cdot 2431$ ,  $C=113^{\circ} 14' 16''$ .
23. If  $a : b = \sin A : \sin B$ , find  $a$ , given  
 $b=378\cdot 25$ ,  $A=35^{\circ} 15' 33''$ ,  $B=119^{\circ} 14' 18''$ .
24. Find the smallest values of  $\theta$  which satisfy the equations  
 $(1) \tan^3 \theta = \frac{5}{12}$ ;  $(2) 3 \sin^2 \theta + 2 \sin \theta = 1$ .
25. Find  $x$  from the equation  
 $x \times \sec 28^{\circ} 17' 25'' = \sin 23^{\circ} 18' 5'' \times \cot 38^{\circ} 15' 13''$ .
26. Find  $\theta$  from the equation  
 $\sin^3 \theta = \cos^2 \alpha \cot \beta$ ,  
 where  $\alpha=32^{\circ} 47'$  and  $\beta=41^{\circ} 19'$ .

## CHAPTER XVI.

### SOLUTION OF TRIANGLES WITH LOGARITHMS.

**183.** The examples on the solution of triangles in Chap. XIII. furnish a useful exercise on the formulæ connecting the sides and angle of a triangle; but in practical work much of the labour of arithmetical calculation is avoided by the use of logarithms.

We shall now shew how the formulæ of Chap. XIII. may be used or adapted for use in connection with logarithmic Tables.

**184.** *To find the functions of the half-angles in terms of the sides.*

$$\begin{aligned}\text{We have } 2 \sin^2 \frac{A}{2} &= 1 - \cos A \\ &= 1 - \frac{b^2 + c^2 - a^2}{2bc} \\ &= \frac{2bc - b^2 - c^2 + a^2}{2bc} = \frac{a^2 - (b^2 - 2bc + c^2)}{2bc} \\ &= \frac{a^2 - (b - c)^2}{2bc} = \frac{(a + b - c)(a - b + c)}{2bc}.\end{aligned}$$

Let

$$a + b + c = 2s;$$

then

$$a + b - c = 2s - 2c = 2(s - c),$$

and

$$a - b + c = 2s - 2b = 2(s - b).$$

$$\therefore 2 \sin^2 \frac{A}{2} = \frac{4(s - c)(s - b)}{2bc} = \frac{2(s - b)(s - c)}{bc};$$

$$\therefore \sin \frac{A}{2} = \sqrt{\frac{(s - b)(s - c)}{bc}}.$$

$$\begin{aligned}
 \text{Again, } 2 \cos^2 \frac{A}{2} &= 1 + \cos A = 1 + \frac{b^2 + c^2 - a^2}{2bc} \\
 &= \frac{(b+c)^2 - a^2}{2bc} = \frac{(b+c+a)(b+c-a)}{2bc}; \\
 \therefore 2 \cos^2 \frac{A}{2} &= \frac{4s(s-a)}{2bc} = \frac{2s(s-a)}{bc}; \\
 \therefore \cos \frac{A}{2} &= \sqrt{\frac{s(s-a)}{bc}}.
 \end{aligned}$$

$$\begin{aligned}
 \text{Also } \tan \frac{A}{2} &= \sin \frac{A}{2} \div \cos \frac{A}{2} \\
 &= \sqrt{\frac{(s-b)(s-c)}{bc} \times \frac{bc}{s(s-a)}}; \\
 \therefore \tan \frac{A}{2} &= \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}.
 \end{aligned}$$

185. Similarly it may be proved that

$$\begin{aligned}
 \sin \frac{B}{2} &= \sqrt{\frac{(s-c)(s-a)}{ca}}, & \sin \frac{C}{2} &= \sqrt{\frac{(s-a)(s-b)}{ab}}; \\
 \cos \frac{B}{2} &= \sqrt{\frac{s(s-b)}{ca}}, & \cos \frac{C}{2} &= \sqrt{\frac{s(s-c)}{ab}}; \\
 \tan \frac{B}{2} &= \sqrt{\frac{(s-c)(s-a)}{s(s-b)}}, & \tan \frac{C}{2} &= \sqrt{\frac{(s-a)(s-b)}{s(s-c)}}.
 \end{aligned}$$

In each of these formulæ the positive value of the square root must be taken, for each half angle is less than  $90^\circ$ , so that all its functions are positive.

186. *To find  $\sin A$  in terms of the sides.*

$$\begin{aligned}
 \sin A &= 2 \sin \frac{A}{2} \cos \frac{A}{2} \\
 &= 2 \sqrt{\frac{(s-b)(s-c)}{bc} \times \frac{s(s-a)}{bc}}; \\
 \therefore \sin A &= \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)}.
 \end{aligned}$$

We may also obtain this formula in another way which is instructive.

We have

$$\begin{aligned}
 \sin^2 A &= 1 - \cos^2 A = (1 + \cos A)(1 - \cos A) \\
 &= \left(1 + \frac{b^2 + c^2 - a^2}{2bc}\right) \left(1 - \frac{b^2 + c^2 - a^2}{2bc}\right) \\
 &= \frac{(b+c)^2 - a^2}{2bc} \times \frac{a^2 - (b-c)^2}{2bc} \\
 &= \frac{(b+c+a)(b+c-a)(a+b-c)(a-b+c)}{4b^2c^2} \\
 &= \frac{16s(s-a)(s-b)(s-c)}{4b^2c^2}; \\
 \therefore \sin A &= \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)}.
 \end{aligned}$$

The positive value of the square root must be taken, since the *sine* of an angle of any triangle is always positive.

### EXAMPLES. XVI. a.

Prove the following formulæ in any triangle :

1.  $b \cos^2 \frac{A}{2} + a \cos^2 \frac{B}{2} = s.$
2.  $s \tan \frac{B}{2} \tan \frac{C}{2} = s - a.$
3.  $\frac{\text{vers } A}{\text{vers } B} = \frac{a(a+c-b)}{b(b+c-a)}.$
4.  $b \sin^2 \frac{A}{2} + a \sin^2 \frac{B}{2} = s - c.$
5.  $(s-a) \tan \frac{A}{2} = (s-b) \tan \frac{B}{2} = (s-c) \tan \frac{C}{2}.$
6. Find the value of  $\tan \frac{B}{2}$ , when  $a=10$ ,  $b=17$ ,  $c=21$ .
7. Find  $\cot \frac{C}{2}$ , when  $a=13$ ,  $b=14$ ,  $c=15$ .
8. Prove that

$$\frac{1}{a} \cos^2 \frac{A}{2} + \frac{1}{b} \cos^2 \frac{B}{2} + \frac{1}{c} \cos^2 \frac{C}{2} = \frac{s^2}{abc}.$$

9. Prove that

$$\frac{b-c}{a} \cos^2 \frac{A}{2} + \frac{c-a}{b} \cos^2 \frac{B}{2} + \frac{a-b}{c} \cos^2 \frac{C}{2} = 0.$$

187. To solve a triangle when the three sides are given.

From the formula

$$\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}},$$

$$\log \tan \frac{A}{2} = \frac{1}{2} \{ \log (s-b) + \log (s-c) - \log s - \log (s-a) \};$$

whence  $\frac{A}{2}$  may be obtained by the help of the Tables.

Similarly  $B$  can be found from the formula for  $\tan \frac{B}{2}$ , and then  $C$  from the equation  $C = 180^\circ - A - B$ .

In the above solution, we shall require to look out from the tables *four* logarithms only, namely those of  $s, s-a, s-b, s-c$ ; whereas if we were to solve from the sine or cosine formulæ we should require *six* logarithms; for

$$\cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}} \quad \text{and} \quad \cos \frac{B}{2} = \sqrt{\frac{s(s-b)}{ca}},$$

so that we should have to look out the logarithms of the *six* quantities  $s, s-a, s-b, a, b, c$ .

If therefore *all* the angles have to be found by the use of the tables it is best to solve from the tangent formulæ; but if *one* angle only is required it is immaterial whether the sine, cosine, or tangent formula is used.

In cases where a solution has to be obtained from certain given logarithms, the choice of formulæ must depend on the data.

NOTE. We shall always find the angles to the nearest second, so that, on account of the multiplication by 2, the half-angles should be found to the nearest tenth of a second.

188. In Art. 178 we have mentioned that 10 is added to each of the logarithmic functions before they are registered as *tabular* logarithms; but this device is introduced only as a convenience for the purposes of tabulation, and in practice it will be found that the work is more expeditious if the tabular logarithms are not used. The 10 should be subtracted mentally in copying down the logarithms. Thus we should write

$$\log \sin 64^\circ 15' = \bar{1} \cdot 9545793, \quad \log \cot 18^\circ 35' = \cdot 4733850,$$

and in the arrangement of the work care must be taken to keep the mantissæ positive.

*Example 1.* The sides of a triangle are 35, 49, 63; find the greatest angle; given  $\log 2 = \cdot 3010300$ ,  $\log 3 = \cdot 4771213$ ,

$$L \cos 47^\circ 53' = 9.8264910, \text{ diff. for } 60'' = 1397.$$

Since the *angles* of a triangle depend only on the *ratios* of the sides and not on their actual magnitudes, we may substitute for the sides any lengths proportional to them. Thus in the present case we may take  $a=5$ ,  $b=7$ ,  $c=9$ ; then  $C$  is the greatest angle, and

$$\cos \frac{C}{2} = \sqrt{\frac{s(s-c)}{ab}} = \sqrt{\frac{21}{2} \times \frac{3}{2} \times \frac{1}{5 \times 7}} = \sqrt{\frac{9}{20}};$$

$$\therefore \log \cos \frac{C}{2} = \frac{1}{2} (2 \log 3 - \log 2 - 1).$$

$$\begin{array}{r} 2 \log 3 = \cdot 9542426 \\ 1 \cdot 3010300 \\ 2) \cdot 6532126 \\ \hline 1 \cdot 8266063 \end{array}$$

Thus  $\log \cos \frac{C}{2} = 1.8266063$

$$\log \cos 47^\circ 53' = 1.8264910$$

$$\text{diff.} \quad 1153$$

$$\therefore \text{proportional decrease} = \frac{1153}{1397} \times 60'' = 49.5'';$$

$$\therefore \frac{C}{2} = 47^\circ 52' 10.5''.$$

$$\begin{array}{r} 1153 \\ 60 \\ 1397) 69180 \text{ ( } 49.5 \\ \hline 5588 \\ 13300 \\ 12573 \\ \hline 7270 \end{array}$$

Thus the greatest angle is  $95^\circ 44' 21''$ .

*Example 2.* If  $a=283$ ,  $b=317$ ,  $c=428$ , find all the angles.

$$\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}} = \sqrt{\frac{197 \times 86}{514 \times 231}};$$

$$\therefore \log \tan \frac{A}{2} = \frac{1}{2} (\log 197 + \log 86 - \log 514 - \log 231).$$

From the Tables,

$$\log 197 = 2.2944662$$

$$\log 86 = 1.9344985$$

$$4.2289647$$

$$5.0745751$$

$$2) 1.1543896$$

$$\log \tan \frac{A}{2} = 1.5771948$$

$$\log \tan 20^\circ 41' = 1.5769585$$

$$\text{diff.} \quad 2363$$

$$\log 514 = 2.7109631$$

$$\log 231 = 2.3636120$$

$$5.0745751$$

$$\begin{array}{r} 283 \\ 317 \\ 428 \\ 2) 1028 \\ \hline 514 = s \\ 231 = s - a \\ 197 = s - b \\ 86 = s - c \end{array}$$



But diff. for 60'' is 3822,

$$\therefore \text{prop}^l. \text{ increase} = \frac{2363}{3822} \times 60'' = 37.1'';$$

$$\therefore \frac{A}{2} = 20^\circ 41' 37.1'' \text{ and } A = 41^\circ 23' 14''.$$

$$\begin{array}{r} 2363 \\ 60 \\ 3822 \overline{) 141780} \quad (37.1 \\ 11466 \\ \hline 27120 \\ 26754 \\ \hline 3660 \end{array}$$

Again,  $\tan \frac{B}{2} = \sqrt{\frac{(s-c)(s-a)}{s(s-b)}} = \sqrt{\frac{86 \times 231}{514 \times 197}};$

$$\therefore \log \tan \frac{B}{2} = \frac{1}{2} (\log 86 + \log 231 - \log 514 - \log 197).$$

$$\log 86 = 1.9344985$$

$$\log 514 = 2.7109631$$

$$\log 231 = 2.3636120$$

$$\log 197 = 2.2944662$$

$$\underline{4.2981105}$$

$$\underline{5.0054293}$$

$$5.0054293$$

$$2 \overline{) 1.2926812}$$

$$\log \tan \frac{B}{2} = 1.6463406$$

$$\log \tan 23^\circ 53' = 1.6461988$$

$$\text{diff.} \quad \underline{1418}$$

But diff. for 60'' is 3412;

$$\therefore \text{prop}^l. \text{ increase} = \frac{1418}{3412} \times 60'' = 24.9''.$$

$$\therefore \frac{B}{2} = 23^\circ 53' 24.9'' \text{ and } B = 47^\circ 46' 50''.$$

$$\begin{array}{r} 1418 \\ 60 \\ 3412 \overline{) 85080} \quad (24.9 \\ 6824 \\ \hline 16840 \\ 13648 \\ \hline 31920 \end{array}$$

Thus  $A = 41^\circ 23' 14''$ ,  $B = 47^\circ 46' 50''$ ,  $C = 90^\circ 49' 56''$ .

### EXAMPLES. XVI. b.

1. The sides of a triangle are 5, 8, 11; find the greatest angle; given  $\log 7 = .8450980$ ,

$$L \sin 56^\circ 47' = 9.9225205, \quad L \sin 56^\circ 48' = 9.9226032.$$

2. If  $a=40$ ,  $b=51$ ,  $c=43$ , find  $A$ ; given

$$L \tan 24^\circ 44' 13'' = 9.6634464,$$

$$\log 128 = 2.1072100, \quad \log 603 = 2.7803173.$$

3. The sides  $a, b, c$  are as 4 : 5 : 6, find  $B$ ; given  $\log 2$ ,

$$L \cos 27^\circ 53' = 9.9464040, \quad \text{diff. for } 1' = 669.$$

4. Find the greatest angle of the triangle in which the sides are 5, 6, 7; given  $\log 6 = \cdot 7781513$ ,

$$L \cos 39^\circ 14' = 9\cdot 8890644, \quad \text{diff. for } 1' = 1032.$$

5. If  $a=3$ ,  $b=1\cdot 75$ ,  $c=2\cdot 75$ , find  $C$ ; given  $\log 2$ ,

$$L \tan 32^\circ 18' = 9\cdot 8008365, \quad \text{diff. for } 1' = 2796.$$

6. If the sides are 24, 22, 14, find the least angle; given

$$L \tan 17^\circ 33' = 9\cdot 500042, \quad \text{diff. for } 1' = 439.$$

7. Find the greatest angle when the sides are 4, 10, 11; given  $\log 2$ ,  $\log 3$ ,

$$L \cos 46^\circ 47' = 9\cdot 8355378, \quad \text{diff. for } 1' = 1345.$$

8. If  $a : b : c = 15 : 13 : 14$ , find the angles ; given  $\log 2$ ,  $\log 3$ ,  $\log 7$ ,

$$L \tan 26^\circ 33' = 9\cdot 6986847, \quad \text{diff. for } 1' = 3159,$$

$$L \tan 29^\circ 44' = 9\cdot 7567587, \quad \text{diff. for } 1' = 2933.$$

9. If  $a : b : c = 3 : 4 : 2$ , find the angles ; given  $\log 2$ ,  $\log 3$ ,

$$L \tan 14^\circ 28' = 9\cdot 4116146, \quad \text{diff. for } 10'' = 870,$$

$$L \tan 52^\circ 14' = 10\cdot 1108395, \quad \text{diff. for } 10'' = 435.$$

189. To solve a triangle having given two sides and the included angle.

Let the given parts be  $b$ ,  $c$ ,  $A$ , and let

$$k = \frac{\sin B}{b} = \frac{\sin C}{c};$$

then

$$\frac{\sin B - \sin C}{\sin B + \sin C} = \frac{kb - kc}{kb + kc} = \frac{b - c}{b + c};$$

$$\therefore \frac{2 \cos \frac{B+C}{2} \sin \frac{B-C}{2}}{2 \sin \frac{B+C}{2} \cos \frac{B-C}{2}} = \frac{b - c}{b + c};$$

$$\therefore \frac{\tan \frac{B-C}{2}}{\tan \frac{B+C}{2}} = \frac{b - c}{b + c};$$

$$\therefore \tan \frac{B-C}{2} = \frac{b-c}{b+c} \tan \frac{B+C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2},$$

since  $\frac{B+C}{2} = 90^\circ - \frac{A}{2}.$

$$\therefore \log \tan \frac{B-C}{2} = \log (b-c) - \log (b+c) + \log \cot \frac{A}{2},$$

from which equation we can find  $\frac{B-C}{2}.$

Also  $\frac{B+C}{2} = 90^\circ - \frac{A}{2},$  and is therefore known.

By addition and subtraction we obtain  $B$  and  $C.$

From the equation  $\alpha = \frac{b \sin A}{\sin B},$

$$\log \alpha = \log b + \log \sin A - \log \sin B;$$

whence  $\alpha$  may be found.

*Example 1.* If the sides  $a$  and  $b$  are in the ratio of 7 to 3, and the included angle  $C$  is  $60^\circ,$  find  $A$  and  $B;$  given

$$\log 2 = \cdot 3010300, \quad \log 3 = \cdot 4771213,$$

$$L \tan 34^\circ 42' = 9 \cdot 8403776, \text{ diff. for } 1' = 2699.$$

$$\tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2} = \frac{7-3}{7+3} \cot 30^\circ = \frac{4}{10} \sqrt{3};$$

$$\therefore \log \tan \frac{A-B}{2} = 2 \log 2 - 1 + \frac{1}{2} \log 3;$$

$$\therefore \log \tan \frac{A-B}{2} = \bar{1} \cdot 8406207$$

$$\log \tan 34^\circ 42' = \bar{1} \cdot 8403776$$

diff. 2431

$$\therefore \text{prop}^1. \text{ increase} = \frac{2431}{2699} \times 60'' = 54'';$$

$$\therefore \frac{A-B}{2} = 34^\circ 42' 54''.$$

And  $\frac{A+B}{2} = 90^\circ - \frac{C}{2} = 60^\circ.$

By addition,  $A = 94^\circ 42' 54'',$

and by subtraction,  $B = 25^\circ 17' 6''.$

$$\begin{array}{r} 2 \log 2 = \cdot 6020600 \\ \frac{1}{2} \log 3 = \cdot 2385607 \\ \hline \cdot 8406207 \end{array}$$

$$\begin{array}{r} 2431 \\ 60 \\ 2699 \overline{) 145860} \text{ ( 54} \\ \underline{13495} \\ 10910 \\ \underline{10796} \end{array}$$

*Example 2.* If  $a=681$ ,  $c=243$ ,  $B=50^\circ 42'$ , solve the triangle, by the use of Tables.

$$\tan \frac{A-C}{2} = \frac{a-c}{a+c} \cot \frac{B}{2} = \frac{438}{924} \cot 25^\circ 21';$$

$$\therefore \log \tan \frac{A-C}{2} = \log 438 - \log 924 \\ + \log \cot 25^\circ 21'$$

$$\therefore \log \tan \frac{A-C}{2} = \cdot 0002383$$

$$\log \tan 45^\circ = \cdot 0000000 \\ \text{diff.} \quad \underline{2383}$$

And diff. for  $60''$  is 2527;

$$\therefore \text{prop}^1. \text{ increase} = \frac{2383}{2527} \times 60'' = 57'';$$

$$\therefore \frac{A-C}{2} = 45^\circ 0' 57''.$$

$$\text{Also } \frac{A+C}{2} = 90^\circ - \frac{B}{2} = 64^\circ 39'.$$

By addition,

$$A = 109^\circ 39' 57'',$$

and by subtraction,

$$C = 19^\circ 38' 3''.$$

$$\text{Again, } b = \frac{c \sin B}{\sin C};$$

$$\therefore \log b = \log c + \log \sin B - \log \sin C \\ = \log 243 + \log \sin 50^\circ 42' \\ - \log \sin 19^\circ 38' 3''$$

$$\therefore \log b = 2\cdot 7479012 \\ \log 559\cdot 63 = \underline{2\cdot 7479010}$$

$$\therefore b = 559\cdot 63.$$

$$\log 438 = 2\cdot 6414741 \\ \log \cot 25^\circ 21' = \cdot 3244362 \\ \underline{2\cdot 9659103} \\ \log 924 = 2\cdot 9656720 \\ \underline{\cdot 0002383}$$

$$\begin{array}{r} 2383 \\ 60 \\ 2527 \overline{) 142980} \quad ( 56\cdot 6 \\ \underline{12635} \\ 16630 \\ \underline{15162} \\ 14680 \end{array}$$

$$\log \sin 19^\circ 38' = 1\cdot 5263387 \\ \frac{3}{60} \times 3540 = 177 \\ \log \sin 19^\circ 38' 3'' = \underline{1\cdot 5263564} \\ \log 243 = 2\cdot 3856063 \\ \log \sin 50^\circ 42' = 1\cdot 8886513 \\ \underline{2\cdot 2742576} \\ \log \sin 19^\circ 38' 3'' = 1\cdot 5263564 \\ \underline{2\cdot 7479012}$$

Thus  $A = 109^\circ 39' 57''$ ,  $C = 19^\circ 38' 3''$ ,  $b = 559\cdot 63$ .

190. From the formula

$$\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2},$$

it will be seen that if  $b$ ,  $c$ , and  $B - C$  are known  $A$  can be found ; that is, the triangle can be solved when the given parts are two sides and the difference of the angles opposite to them.

### EXAMPLES. XVI. c.

1. If  $a=9$ ,  $b=6$ ,  $C=60^\circ$ , find  $A$  and  $B$ ; given  $\log 2$ ,  $\log 3$ ,  
 $L \tan 19^\circ 6' = 9.5394287$ ,  $L \tan 19^\circ 7' = 9.5398371$ .
2. If  $a=1$ ,  $c=9$ ,  $B=65^\circ$ , find  $A$  and  $C$ ; given  $\log 2$ ,  
 $L \cot 32^\circ 30' = 10.1958127$ ,  
 $L \tan 51^\circ 28' = 10.0988763$ , diff. for  $1' = 2592$ .
3. If  $17a=7b$ ,  $C=60^\circ$ , find  $A$  and  $B$ ; given  $\log 2$ ,  $\log 3$ ,  
 $L \tan 35^\circ 49' = 9.8583357$ , diff. for  $10'' = 2662$ .
4. If  $b=27$ ,  $c=23$ ,  $A=44^\circ 30'$ , find  $B$  and  $C$ ; given  $\log 2$ ,  
 $L \cot 22^\circ 15' = 10.3881591$ ,  
 $L \tan 11^\circ 3' = 9.2906713$ , diff. for  $1' = 6711$ .
5. If  $c=210$ ,  $a=110$ ,  $B=34^\circ 42' 30''$ , find  $C$  and  $A$ ;  
given  $\log 2$ ,  
 $L \cot 17^\circ 21' 15'' = 10.5051500$ .
6. Two sides of a triangle are as  $5 : 3$  and include an angle  
of  $60^\circ 30'$ : find the other angles; given  $\log 2$ ,  
 $L \cot 30^\circ 15' = 10.23420$ ,  
 $L \tan 23^\circ 13' = 9.63240$ , diff. for  $1' = 35$ .
7. If  $a=327$ ,  $c=256$ ,  $B=56^\circ 28'$ , find  $A$  and  $C$ ; given  
 $\log 7.1 = .8512583$ ,  $\log 5.83 = .7656686$ ,  
 $L \tan 61^\circ 46' = 10.2700705$ ,  
 $L \tan 12^\circ 46' = 9.3552267$ , diff. for  $1' = 5859$ .
8. If  $b=4c$ ,  $A=65^\circ$ , find  $B$  and  $C$ ; given  $\log 2$ ,  $\log 3$ ,  
 $L \tan 57^\circ 30' = 10.1958127$ ,  
 $L \tan 43^\circ 18' = 9.9742133$ , diff. for  $1' = 2531$ .
9. If  $a=23031$ ,  $b=7677$ ,  $C=30^\circ 10' 5''$ , find  $A$  and  $B$ ;  
given  $\log 2$ ,  
 $L \tan 15^\circ 5' = 9.4305727$ , diff. for  $10'' = 838$ ,  
 $L \cot 61^\circ 41' = 9.7314436$ , diff. for  $10'' = 504$ .

191. To solve a triangle having given two angles and a side.

Let the given parts be denoted by  $B, C, a$ ; then the third angle  $A$  is found from the equation  $A = 180^\circ - B - C$ ,

and 
$$b = \frac{a \sin B}{\sin A};$$

$$\therefore \log b = \log a + \log \sin B - \log \sin A;$$

whence  $b$  may be found.

Similarly,  $c$  may be obtained from the equation

$$\log c = \log a + \log \sin C - \log \sin A.$$

*Example.* If  $b = 1000$ ,  $A = 45^\circ$ ,  $C = 68^\circ 17' 40''$ , find the least side, having given

$$\log 2 = \cdot 3010300, \log 7 \cdot 6986 = \cdot 8864118, \text{ diff. for } 1 = 57,$$

$$L \sin 66^\circ 42' = 9 \cdot 9630538, \text{ diff. for } 1' = 544.$$

$$B = 180^\circ - 45^\circ - 68^\circ 17' 40'' = 66^\circ 42' 20''.$$

$$\text{The least side} = a = \frac{b \sin A}{\sin B} = \frac{1000 \sin 45^\circ}{\sin 66^\circ 42' 20''};$$

$\begin{aligned} \therefore \log a &= 3 + \log \frac{1}{\sqrt{2}} - \log \sin 66^\circ 42' 20'' \\ &= 3 - \frac{1}{2} \log 2 - \log \sin 66^\circ 42' 20'' \\ &= 3 - \cdot 1135869 \end{aligned}$	$\begin{aligned} \log \sin 66^\circ 42' &= \bar{1} \cdot 9630538 \\ \frac{20}{60} \times 544 &= 181 \\ \frac{1}{2} \log 2 &= \cdot 1505150 \\ \hline &= \cdot 1135869 \end{aligned}$
---	--

$$\begin{aligned} \therefore \log a &= 2 \cdot 8864131 \\ \log 769 \cdot 86 &= 2 \cdot 8864118 \\ \text{diff.} &= \underline{13} \end{aligned}$$

$$\therefore \text{prop}^l. \text{ increase} = \frac{13}{57} = \cdot 22.$$

Thus the least side is 769·8622.

### EXAMPLES. XVI. d.

- If  $B = 60^\circ 15'$ ,  $C = 54^\circ 30'$ ,  $a = 100$ , find  $c$ ; given  
 $L \sin 54^\circ 30' = 9 \cdot 9106860$ ,  $\log 8 \cdot 9646162 = \cdot 9525317$ ,  
 $L \sin 65^\circ 15' = 9 \cdot 9581543$ .



2. If  $A = 55^\circ$ ,  $B = 65^\circ$ ,  $c = 270$ , find  $a$ ; given  $\log 2$ ,  $\log 3$ ,  
 $\log 25538 = 4.4071869$ ,  $L \sin 55^\circ = 9.9133645$ ,  
 $\log 25539 = 4.4072039$ .
3. If  $A = 45^\circ 41'$ ,  $C = 62^\circ 5'$ ,  $b = 100$ , find  $c$ ; given  
 $\log 9.2788 = .96749$ ,  $L \sin 62^\circ 5' = 9.94627$ ,  
 $L \sin 72^\circ 14' = 9.97878$ .
4. If  $B = 70^\circ 30'$ ,  $C = 78^\circ 10'$ ,  $a = 102$ , find  $b$  and  $c$ ; given  
 $\log 2 = .301$ ,  $\log 1.02 = .009$ ,  $\log 1.85 = .267$ ,  $\log 1.92 = .283$ ,  
 $L \sin 70^\circ 30' = 9.974$ ,  $L \sin 78^\circ 10' = 9.990$ ,  
 $L \sin 31^\circ 20' = 9.716$ .
5. If  $a = 123$ ,  $B = 29^\circ 17'$ ,  $C = 135^\circ$ , find  $c$ ; given  $\log 2$ ,  
 $\log 123 = 2.0899051$ ,  $L \sin 15^\circ 43' = 9.4327777$ ,  
 $\log 3211 = 3.5066403$ ,  $D = 135$ .
6. If  $A = 44^\circ$ ,  $C = 70^\circ$ ,  $b = 1006.62$ , find  $a$  and  $c$ ; given  
 $L \sin 44^\circ = 9.8417713$ ,  $\log 100662 = 5.0028656$ ,  
 $L \sin 66^\circ = 9.9607302$ ,  $\log 103543 = 5.0151212$ ,  
 $L \sin 70^\circ = 9.9729858$ ,  $\log 7654321 = 6.8839067$ .
7. If  $a = 1652$ ,  $B = 26^\circ 30'$ ,  $C = 47^\circ 15'$ , find  $b$  and  $c$ ;  
 $L \sin 73^\circ 45' = 9.9822938$ ,  $\log 1.652 = .2180100$ ,  
 $L \sin 26^\circ 30' = 9.6495274$ ,  $\log 7.6780 = .8852481$ ,  $D = 57$ ,  
 $L \sin 47^\circ 15' = 9.8658868$ ,  $\log 1.2636 = .1016096$ ,  $D = 344$ .

192. *To solve a triangle when two sides and the angle opposite to one of them are given.*

Let  $a, b, A$  be given. Then from  $\sin B = \frac{b}{a} \sin A$ , we have

$$\log \sin B = \log b - \log a + \log \sin A;$$

whence  $B$  may be found;

then  $C$  is found from the equation  $C = 180^\circ - A - B$ .

Again, 
$$c = \frac{a \sin C}{\sin A},$$

$$\therefore \log c = \log a + \log \sin C - \log \sin A.$$

If  $a < b$ , and  $A$  is acute the solution is ambiguous and there will be two values of  $B$  supplementary to each other, and also two values of  $C$  and  $c$ . [Art. 147.]

*Example.* If  $b=63$ ,  $c=36$ ,  $C=29^\circ 23' 15''$ , find  $B$ ; given

$$\log 2 = \cdot 3010300, \log 7 = \cdot 8450980.$$

$$L \sin 29^\circ 23' = 9\cdot 6907721, \text{ diff. for } 1' = 2243,$$

$$L \sin 59^\circ 10' = 9\cdot 9338222, \text{ diff. for } 1' = 755.$$

$$\sin B = \frac{b}{c} \sin C = \frac{63}{36} \sin C$$

$$= \frac{7}{4} \sin 29^\circ 23' 15'';$$

$$\therefore \log \sin B = \log 7 - 2 \log 2 \\ + \log \sin 29^\circ 23' 15'';$$

$$\therefore \log \sin B = \bar{1}\cdot 9338662$$

$$\log \sin 59^\circ 10' = \bar{1}\cdot 9338222 \\ \text{diff.} \quad \underline{440}$$

$$\therefore \text{prop}^l. \text{ increase} = \frac{440}{755} \times 60'' = 35'';$$

$$\therefore B = 59^\circ 10' 35''.$$

Also since  $c < b$  there is another value of  $B$  supplementary to the above, namely  $B = 120^\circ 49' 25''$ .

$$\log \sin 29^\circ 23' = \bar{1}\cdot 6907721$$

$$\frac{15}{60} \times 2243 = \quad 561$$

$$\log 7 = \cdot 8450980$$

$$\underline{\cdot 5359262}$$

$$2 \log 2 = \cdot 6020600$$

$$\underline{\bar{1}\cdot 9338662}$$

$$440$$

$$\underline{60}$$

$$755 \mid \begin{array}{r} 26400 \\ 2265 \end{array} ( 35$$

$$\underline{3750}$$

$$\underline{3775}$$

### EXAMPLES. XVI. e.

1. If  $a=145$ ,  $b=178$ ,  $B=41^\circ 10'$ , find  $A$ ; given

$$\log 178 = 2\cdot 2504200, \quad L \sin 41^\circ 10' = 9\cdot 8183919,$$

$$\log 145 = 2\cdot 1613680, \quad L \sin 32^\circ 25' 35'' = 9\cdot 7293399.$$

2. If  $A=26^\circ 26'$ ,  $b=127$ ,  $a=85$ , find  $B$ ; given

$$\log 1\cdot 27 = \cdot 1038037, \quad L \sin 26^\circ 26' = 9\cdot 6485124,$$

$$\log 8\cdot 5 = \cdot 9294189, \quad L \sin 41^\circ 41' 28'' = 9\cdot 8228972.$$

3. If  $c=5$ ,  $b=4$ ,  $C=45^\circ$ , find  $A$  and  $B$ ; given  $\log 2$ ,

$$L \sin 34^\circ 26' = 9\cdot 7525750, \quad L \sin 34^\circ 27' = 9\cdot 7525761.$$

4. If  $a=1405$ ,  $b=1706$ ,  $A=40^\circ$ , find  $B$ ; given

$$\log 1\cdot 405 = \cdot 1476763, \quad \log 1\cdot 706 = \cdot 2319790,$$

$$L \sin 40^\circ = 9\cdot 8080675, \quad L \sin 51^\circ 18' = 9\cdot 8923342,$$

$$\text{diff. for } 1' = 1012.$$

5. If  $B=112^\circ 4'$ ,  $b=573$ ,  $c=394$ , find  $A$  and  $C$ ; given  
 $\log 573=2.7581546$ ,  $\log 394=2.5954962$ ,  
 $L \sin 39^\circ 35'=9.8042757$ , diff. for  $60''=1527$ ,  
 $L \cos 22^\circ 4'=9.9669614$ .

6. If  $b=8.4$ ,  $c=12$ ,  $B=37^\circ 36'$ , find  $A$ ; given  
 $\log 7=.8450980$ ,  $L \sin 37^\circ 36'=9.7854332$ ,  
 $L \sin 60^\circ 39'=9.9403381$ , diff. for  $1'=711$ .

7. Supposing the data for the solution of a triangle to be as in the three following cases, point out whether the solution will be ambiguous or not, and find the third side in the obtuse angled triangle in the ambiguous case:

- (i)  $A=30^\circ$ ,  $a=125$  feet,  $c=250$  feet,  
(ii)  $A=30^\circ$ ,  $a=200$  feet,  $c=250$  feet,  
(iii)  $A=30^\circ$ ,  $a=200$  feet,  $c=125$  feet.

Given  $\log 2$ ,

$$\begin{aligned}\log 6.0389 &= .7809578, & L \sin 38^\circ 41' &= 9.7958800, \\ \log 6.0390 &= .7809650, & L \sin 8^\circ 41' &= 9.1789001.\end{aligned}$$

193. Some formulæ which are not primarily suitable for working with logarithms may be adapted to such work by various artifices.

194. To adapt the formula  $c^2=a^2+b^2$  to logarithmic computation.

We have 
$$c^2=a^2\left(1+\frac{b^2}{a^2}\right).$$

Since an angle can always be found whose tangent is equal to a given numerical quantity, we may put  $\frac{b}{a}=\tan \theta$ , and thus obtain

$$c^2=a^2(1+\tan^2 \theta)=a^2 \sec^2 \theta;$$

$$\therefore c=a \sec \theta;$$

$$\therefore \log c=\log a+\log \sec \theta.$$

The angle  $\theta$  is called a **subsidiary angle** and is found from the equation

$$\log \tan \theta=\log b-\log a.$$

Thus any expression which can be put into the form of the sum of two squares can be readily adapted to logarithmic work.

195. To adapt the formula  $c^2 = a^2 + b^2 - 2ab \cos C$  to logarithmic computation.

From the identities

$$\cos C = \cos^2 \frac{C}{2} - \sin^2 \frac{C}{2}, \text{ and } 1 = \cos^2 \frac{C}{2} + \sin^2 \frac{C}{2},$$

we have

$$\begin{aligned} c^2 &= (a^2 + b^2) \left( \cos^2 \frac{C}{2} + \sin^2 \frac{C}{2} \right) - 2ab \left( \cos^2 \frac{C}{2} - \sin^2 \frac{C}{2} \right) \\ &= (a^2 + b^2 - 2ab) \cos^2 \frac{C}{2} + (a^2 + b^2 + 2ab) \sin^2 \frac{C}{2} \\ &= (a - b)^2 \cos^2 \frac{C}{2} + (a + b)^2 \sin^2 \frac{C}{2} \\ &= (a - b)^2 \cos^2 \frac{C}{2} \left\{ 1 + \left( \frac{a + b}{a - b} \right)^2 \tan^2 \frac{C}{2} \right\}. \end{aligned}$$

Take a subsidiary angle  $\theta$ , such that

$$\tan \theta = \frac{a + b}{a - b} \tan \frac{C}{2},$$

$$\text{then } c^2 = (a - b)^2 \cos^2 \frac{C}{2} (1 + \tan^2 \theta)$$

$$= (a - b)^2 \cos^2 \frac{C}{2} \sec^2 \theta;$$

$$\therefore c = (a - b) \cos \frac{C}{2} \sec \theta;$$

$$\therefore \log c = \log (a - b) + \log \cos \frac{C}{2} + \log \sec \theta,$$

where  $\theta$  is determined from the equation

$$\log \tan \theta = \log (a + b) - \log (a - b) + \log \tan \frac{C}{2}.$$

196. When two sides and the included angle are given, we may solve the triangle by finding the value of the third side first instead of determining the angles first as in Art. 189.

*Example.* If  $a = 3$ ,  $c = 1$ ,  $B = 53^\circ 7' 48''$  find  $b$ ; given

$$\log 2 = .3010300, \log 2.5298 = .4030862, \text{ diff. for } 1 = 172,$$

$$L \cos 26^\circ 33' 54'' = 9.9515452, L \tan 26^\circ 33' 54'' = 9.6989700.$$

We have  $b^2 = c^2 + a^2 - 2ca \cos B$

$$\begin{aligned} &= (a^2 + c^2) \left( \cos^2 \frac{B}{2} + \sin^2 \frac{B}{2} \right) - 2ac \left( \cos^2 \frac{B}{2} - \sin^2 \frac{B}{2} \right) \\ &= (a - c)^2 \cos^2 \frac{B}{2} + (a + c)^2 \sin^2 \frac{B}{2} \\ &= (a - c)^2 \cos^2 \frac{B}{2} \left\{ 1 + \left( \frac{a + c}{a - c} \right)^2 \tan^2 \frac{B}{2} \right\} \\ &= (a - c)^2 \cos^2 \frac{B}{2} (1 + \tan^2 \theta) \dots \dots \dots (1), \end{aligned}$$

where  $\tan \theta = \frac{a + c}{a - c} \tan \frac{B}{2} = 2 \tan 26^\circ 33' 54'';$

$$\begin{aligned} \therefore \log \tan \theta &= \log 2 + \log \tan 26^\circ 33' 54'' \\ &= .3010300 + \bar{1}.6989700 \\ &= 0; \end{aligned}$$

whence  $\tan \theta = 1$ , and  $\theta = 45^\circ$ .

$$\begin{aligned} \text{From (1),} \quad b &= (a - c) \cos \frac{B}{2} \sec \theta \\ &= 2 \sec 45^\circ \cos \frac{B}{2} \\ &= 2\sqrt{2} \cos 26^\circ 33' 54''; \end{aligned}$$

$\therefore \log b = \log 2 + \frac{1}{2} \log 2$ $\quad + \log \cos 26^\circ 33' 54''$	$\log 2 = .3010300$
$\therefore \log b = .4030902$	$\frac{1}{2} \log 2 = .1505150$
$\log 2.5298 = .4030862$	$\log \cos 26^\circ 33' 54'' = \bar{1}.9515452$
$\text{diff.} \quad \underline{\quad 40 \quad}$	$\underline{\quad .4030902 \quad}$

But diff. for 1 is 172;

$$\therefore \text{proportional increase} = \frac{40}{172} = \frac{10}{43} = .23.$$

Thus the third side is 2.529823.

197. The formula  $c^2 = a^2 + b^2 - 2ab \cos C$  may also be adapted to logarithmic computation in two other ways by making use of the identities  $\cos C = 2 \cos^2 \frac{C}{2} - 1$  and  $\cos C = 1 - 2 \sin^2 \frac{C}{2}$ .

We shall take the first of these cases, leaving the other as an exercise.

$$\begin{aligned} c^2 &= a^2 + b^2 - 2ab \cos C \\ &= a^2 + b^2 - 2ab \left( 2 \cos^2 \frac{C}{2} - 1 \right) \\ &= (a+b)^2 - 4ab \cos^2 \frac{C}{2} \\ &= (a+b)^2 \left\{ 1 - \frac{4ab}{(a+b)^2} \cos^2 \frac{C}{2} \right\}. \end{aligned}$$

Let 
$$\frac{4ab}{(a+b)^2} \cos^2 \frac{C}{2} = \cos^2 \theta,$$

then 
$$c^2 = (a+b)^2 (1 - \cos^2 \theta) = (a+b)^2 \sin^2 \theta;$$

$$\therefore c = (a+b) \sin \theta;$$

$$\therefore \log c = \log (a+b) + \log \sin \theta.$$

To determine  $\theta$  we have the equation

$$\cos \theta = \frac{2\sqrt{ab}}{a+b} \cos \frac{C}{2};$$

$$\therefore \log \cos \theta = \log 2 + \frac{1}{2} (\log a + \log b) - \log (a+b) + \log \cos \frac{C}{2}.$$

Since  $2\sqrt{ab}$  is never greater than  $a+b$  and  $\cos \frac{C}{2}$  is positive and less than unity,  $\cos \theta$  is positive and less than unity, and thus  $\theta$  is an acute angle.

### EXAMPLES. XVI. f.

1. If  $a=8$ ,  $b=7$ ,  $c=9$ , find the angles ; given  $\log 2$ ,  $\log 3$ ,  
 $L \tan 24^\circ 5' = 9.6502809$ , diff. for  $60'' = 3390$ ,  
 $L \tan 36^\circ 41' = 9.8721123$ , diff. for  $60'' = 2637$ .

2. The difference between the angles at the base of a triangle is  $24^\circ$ , and the sides opposite these angles are 175 and 337 : find all the angles ; given  $\log 2$ ,  $\log 3$ ,

$$L \tan 12^\circ = 9.3274745, \quad L \cot 56^\circ 6' 27'' = 9.8272293.$$



3. One of the sides of a right-angled triangle is two-sevenths of the hypotenuse: find the greater of the two acute angles; given  $\log 2$ ,  $\log 7$ ,  $L \sin 14^\circ 11' = 9.455921$ ,  $L \sin 14^\circ 12' = 9.456031$ .

4. Find the greatest side when two of the angles are  $78^\circ 14'$  and  $71^\circ 24'$  and the sides joining them is 2183; given

$$\log 2.183 = .3390537, \quad \log 4.2274 = .6260733, \quad D = 103, \\ L \sin 78^\circ 14' = 9.9907766, \quad L \sin 30^\circ 22' = 9.7037486.$$

5. If  $b = 2$  ft. 6 in.,  $c = 2$  ft.,  $A = 22^\circ 20'$ , find the other angles; and then shew that the side  $a$  is very approximately 1 foot. Given  $\log 2$ ,  $\log 3$ ,

$$L \cot 11^\circ 10' = 10.70465, \quad L \sin 49^\circ 27' 34'' = 9.88079, \\ L \sin 22^\circ 20' = 9.57977, \quad L \tan 29^\circ 22' 26'' = 9.75041.$$

6. If  $a = 1.56234$ ,  $b = .43766$ ,  $C = 58^\circ 42' 6''$ , find  $A$  and  $B$ ; given  $\log 56234 = 4.75$ ,

$$\log \cot 29^\circ 21' = .250015, \quad \log \cot 29^\circ 22' = .249715.$$

7. If  $a = 9$ ,  $b = 12$ ,  $A = 30^\circ$ , find the values of  $c$ , having given

$$\log 12 = 1.07918, \quad L \sin 30^\circ = 9.69897, \\ \log 9 = .95424, \quad L \sin 11^\circ 48' 39'' = 9.31108, \\ \log 171 = 2.23301, \quad L \sin 41^\circ 48' 39'' = 9.82391, \\ \log 368 = 2.56635, \quad L \sin 108^\circ 11' 21'' = 9.97774.$$

8. The sides of a triangle are 9 and 3, and the difference of the angles opposite to them is  $90^\circ$ : find the angles; having given  $\log 2$ ,

$$L \tan 26^\circ 33' = 9.6986847, \quad L \tan 26^\circ 34' = 9.6990006.$$

9. Two sides of a triangle are 1404 and 960 respectively, and an angle opposite to one of them is  $32^\circ 15'$ : find the angle contained by the two sides; having given  $\log 2$ ,  $\log 3$ ,

$$\log 13 = 1.1139434, \quad L \operatorname{cosec} 32^\circ 15' = 10.2727724, \\ L \sin 21^\circ 23' = 9.5621316, \quad L \sin 51^\circ 18' = 9.8923236.$$

10. If  $b : c = 11 : 10$  and  $A = 35^\circ 25'$ , use the formula

$$\tan \frac{1}{2}(B - C) = \tan^2 \frac{\phi}{2} \cot \frac{A}{2} \text{ to find } B \text{ and } C;$$

given  $\log 1.1 = .041393$ ,  $L \tan 12^\circ 18' 36'' = 9.338891$ ,  
 $L \cos 24^\circ 37' 12'' = 9.958607$ ,  $L \cot 17^\circ 42' 30'' = 10.495800$ ,  
 $L \tan 8^\circ 28' 56.5'' = 9.173582$ .

11. If  $A = 50^\circ$ ,  $b = 1071$ ,  $a = 873$ , find  $B$ ; given  
 $\log 1.071 = .029789$ ,  $\log 8.73 = .941014$ ,  
 $L \sin 50^\circ = 9.884254$ ,  $L \sin 70^\circ = 9.972986$ ,  
 $L \sin 70^\circ 1' = 9.973032$ .

12. If  $a = 3$ ,  $b = 1$ ,  $C = 53^\circ 7' 48''$ , find  $c$  without determining  $A$  and  $B$ ; given  $\log 2$ ,  
 $\log 25298 = 4.4030862$ ,  $L \cos 26^\circ 33' 54'' = 9.9515452$ ,  
 $\log 25299 = 4.4031034$ ,  $L \tan 26^\circ 33' 54'' = 9.6989700$ .

*(In the following Examples the necessary Logarithms must be taken from the Tables.)*

13. Given  $a = 1000$ ,  $b = 840$ ,  $c = 1258$ , find  $B$ .
14. Solve the triangle in which  $a = 525$ ,  $b = 650$ ,  $c = 777$ .
15. Find the least angle when the sides are proportional to 4, 5, and 6.
16. If  $B = 90^\circ$ ,  $AC = 57.321$ ,  $AB = 28.58$ , find  $A$  and  $C$ .
17. Find the hypotenuse of a right-angled triangle in which the smallest angle is  $18^\circ 37' 29''$  and the side opposite to it is 284 feet.
18. The sides of a triangle are 9 and 7 and the angle between them is  $60^\circ$ : find the other angles.
19. How long must a ladder be so that when inclined to the ground at an angle of  $72^\circ 15'$  it may just reach a window 42.37 feet from the ground?
20. If  $a = 31.95$ ,  $b = 21.96$ ,  $C = 35^\circ$ , find  $A$  and  $B$ .
21. Find  $B$ ,  $C$ ,  $a$  when  $b = 25.12$ ,  $c = 13.83$ ,  $A = 47^\circ 15''$ .
22. Find the greatest angle of the triangle whose sides are 1837.2, 2385.6, 2173.84.
23. When  $a = 21.352$ ,  $b = 45.6843$ ,  $c = 37.2134$ , find  $A$ ,  $B$ , and  $C$ .
24. If  $b = 647.324$ ,  $c = 850.273$ ,  $A = 103^\circ 12' 54''$ , find the remaining parts.
25. If  $b = 23.2783$ ,  $A = 37^\circ 57'$ ,  $B = 43^\circ 13'$ , find the remaining sides.
26. Find  $a$  and  $b$  when  $B = 72^\circ 43' 25''$ ,  $C = 47^\circ 12' 17''$ ,  $c = 2484.3$ .

27. If  $AB=4517$ ,  $AC=150$ ,  $A=31^{\circ} 30''$ , find the remaining parts.

28. Find  $A$ ,  $B$ , and  $b$  when

$$a=324\cdot68, c=421\cdot73, C=35^{\circ} 17' 12''.$$

29. Given  $a=321\cdot7$ ,  $c=435\cdot6$ ,  $A=36^{\circ} 18' 27''$ , find  $C$ .

30. If  $b=1625$ ,  $c=1665$ ,  $B=52^{\circ} 19'$ , solve the obtuse-angled triangle to which the data belong.

31. If  $a=3795$ ,  $B=73^{\circ} 15' 15''$ ,  $C=42^{\circ} 18' 30''$ , find the other sides.

32. Find the angles of the two triangles which have  $b=17$ ,  $c=12$ , and  $C=43^{\circ} 12' 12''$ .

33. Two sides of a triangle are  $2\cdot7402$  ft. and  $\cdot7401$  ft. respectively, and contain an angle  $59^{\circ} 27' 5''$ : find the base and altitude of the triangle.

34. The difference between the angles at the base of a triangle is  $17^{\circ} 48'$  and the sides subtending these angles are  $105\cdot25$  ft. and  $76\cdot75$  ft.: find the angle included by the given sides.

35. From the following data :

$$(1) \quad A=43^{\circ} 15', \quad AB=36\cdot5, \quad BC=20,$$

$$(2) \quad A=43^{\circ} 15', \quad AB=36\cdot5, \quad BC=30,$$

$$(3) \quad A=43^{\circ} 15', \quad AB=36\cdot5, \quad BC=45,$$

point out which solution is impossible and which ambiguous. Find the third side for the triangle the solution of which is neither impossible nor ambiguous.

36. In any triangle prove that  $c=(a-b)\sec\theta$ , where

$$\tan\theta=\frac{2\sqrt{ab}}{a-b}\sin\frac{C}{2}.$$

If  $a=17\cdot32$ ,  $b=13\cdot47$ ,  $C=47^{\circ} 13'$ , find  $c$  without finding  $A$  and  $B$ .

37. If  $\tan\phi=\frac{a+b}{a-b}\tan\frac{C}{2}$ , prove that  $c=(a-b)\cos\frac{C}{2}\sec\phi$ .

If  $a=27\cdot3$ ,  $b=16\cdot8$ ,  $C=45^{\circ} 12'$ , find  $\theta$ , and thence find  $c$ .

## CHAPTER XVII.

### HEIGHTS AND DISTANCES.


198. Some easy cases of heights and distances depending only on the solution of right-angled triangles have been already dealt with in Chap. VI. The problems in the present chapter are of a more general character, and require for their solution some geometrical skill as well as a ready use of trigonometrical formulæ.

#### Measurements in one plane.

199. *To find the height and distance of an inaccessible object on a horizontal plane.*

Let  $A$  be the position of the observer,  $CP$  the object; from  $P$  draw  $PC$  perpendicular to the horizontal plane; then it is required to find  $PC$  and  $AC$ .

At  $A$  observe the angle of elevation  $PAC$ . Measure a base line  $AB$  in a direct line from  $A$  towards the object, and at  $B$  observe the angle of elevation  $PBC$ .



Let  $\angle PAC = \alpha$ ,  $\angle PBC = \beta$ ,  $AB = a$ .

From  $\triangle PBC$ ,  $PC = PB \sin \beta$ .

From  $\triangle PAB$ ,

$$PB = \frac{AB \sin PAB}{\sin APB} = \frac{a \sin \alpha}{\sin (\beta - \alpha)};$$

$$\therefore PC = a \sin \alpha \sin \beta \operatorname{cosec} (\beta - \alpha).$$

Also  $AC = PC \cot \alpha = a \cos \alpha \sin \beta \operatorname{cosec} (\beta - \alpha)$ .

Each of the above expressions is adapted to logarithmic work; thus if  $PC = x$ , we have

$$\log x = \log a + \log \sin \alpha + \log \sin \beta + \log \operatorname{cosec} (\beta - \alpha).$$

**NOTE.** Unless the contrary is stated, it will be supposed that the observer's height is disregarded, and that the angles of elevation are measured from the ground.

**Example I.** A person walking along a straight road observes that at two consecutive milestones the angles of elevation of a hill in front of him are  $30^\circ$  and  $75^\circ$ : find the height of the hill.

In the adjoining figure,

$\angle PAC = 30^\circ$ ,  $\angle PBC = 75^\circ$ ,  $AB = 1$  mile;

$\angle APB = 75^\circ - 30^\circ = 45^\circ$ .

Let  $x$  be the height in yards; then


$$x = PB \sin 75^\circ;$$

$$\text{but } PB = \frac{AB \sin PAB}{\sin APB} = \frac{1760 \sin 30^\circ}{\sin 45^\circ};$$

$$\therefore x = \frac{1760 \sin 30^\circ \sin 75^\circ}{\sin 45^\circ}$$

$$= 1760 \times \frac{1}{2} \times \frac{\sqrt{2}}{2} \times \frac{\sqrt{3}+1}{2\sqrt{2}}$$

$$= 440(\sqrt{3}+1).$$



If we take  $\sqrt{3} = 1.732$  and reduce to feet, we find that the height is 3606.24 ft.

### EXAMPLES. XVII. a.

1. From the top of a cliff 200 ft. above the sea-level the angles of depression of two boats in the same vertical plane as the observer are  $45^\circ$  and  $30^\circ$ : find their distance apart.

2. A person observes the elevation of a mountain top to be  $15^\circ$ , and after walking a mile directly towards it on level ground the elevation is  $75^\circ$ : find the height of the mountain in feet.

3. From a ship at sea the angle subtended by two forts  $A$  and  $B$  is  $30^\circ$ . The ship sails 4 miles towards  $A$  and the angle is then  $48^\circ$ : prove that the distance of  $B$  at the second observation is 6.472 miles.

4. From the top of a tower  $h$  ft. high the angles of depression of two objects on the horizontal plane and in a line passing through the foot of the tower are  $45^\circ - A$  and  $45^\circ + A$ . Shew that the distance between them is  $2h \tan 2A$ .



5. An observer finds that the angular elevation of a tower is  $A$ . On advancing  $a$  feet towards the tower the elevation is  $45^\circ$  and on advancing  $b$  feet nearer the elevation is  $90^\circ - A$ : find the height of the tower.

6. A person observes that two objects  $A$  and  $B$  bear due N. and N.  $30^\circ$  W. respectively. On walking a mile in the direction N.W. he finds that the bearings of  $A$  and  $B$  are N.E. and due E. respectively: find the distance between  $A$  and  $B$ .

7. A tower stands at the foot of a hill whose inclination to the horizon is  $9^\circ$ ; from a point 40 ft. up the hill the tower subtends an angle of  $54^\circ$ : find its height.

8. At a point on a level plane a tower subtends an angle  $\alpha$  and a flagstaff  $c$  ft. in length at the top of the tower subtends an angle  $\beta$ : shew that the height of the tower is

$$c \sin \alpha \operatorname{cosec} \beta \cos (\alpha + \beta).$$


**Example II.** The upper three-fourths of a ship's mast subtends at a point on the deck an angle whose tangent is  $\cdot 6$ ; find the tangent of the angle subtended by the whole mast at the same point.

Let  $C$  be the point of observation, and let  $APB$  be the mast,  $AP$  being the lower fourth of it.

Let  $AB = 4a$ , so that  $AP = a$ ;  
also let  $AC = b$ ,  $\angle ACB = \theta$ ,  $\angle BCP = \beta$ ,  
so that  $\tan \beta = \cdot 6$ .

$$\text{From } \triangle PCA, \quad \tan (\theta - \beta) = \frac{a}{b};$$

$$\text{from } \triangle BCA, \quad \tan \theta = \frac{4a}{b};$$



$$\therefore \tan \theta = 4 \tan (\theta - \beta) = \frac{4 (\tan \theta - \tan \beta)}{1 + \tan \theta \tan \beta};$$

$$\therefore \tan \theta = \frac{4 \left( \tan \theta - \frac{3}{5} \right)}{1 + \frac{3}{5} \tan \theta} = \frac{4 (5 \tan \theta - 3)}{5 + 3 \tan \theta}.$$

$$\text{On reduction,} \quad \tan^2 \theta - 5 \tan \theta + 4 = 0;$$

$$\text{whence} \quad \tan \theta = 1 \text{ or } 4.$$

**NOTE.** The student should observe that in examples of this class we make use of right-angled triangles in which the horizontal base line forms one side.



**Example III.** A tower  $BCD$  surmounted by a spire  $DE$  stands on a horizontal plane. From the extremity  $A$  of a horizontal line  $BA$ , it is found that  $BC$  and  $DE$  subtend equal angles. If  $BC=9$  ft.,  $CD=72$  ft., and  $DE=36$  ft., find  $BA$ .

Let  $\angle BAC = \angle DAE = \theta$ ,  
 $\angle DAB = \alpha$ ,  $AB = x$  ft.

Now  $BC=9$  ft.,  $BD=81$  ft.,  $BE=117$  ft.

$$\therefore \tan(\alpha + \theta) = \frac{BE}{AB} = \frac{117}{x};$$

$$\tan \alpha = \frac{BD}{AB} = \frac{81}{x};$$

$$\tan \theta = \frac{BC}{AB} = \frac{9}{x}.$$

But  $\tan(\alpha + \theta) = \frac{\tan \alpha + \tan \theta}{1 - \tan \alpha \tan \theta};$

$$\therefore \frac{117}{x} = \frac{\frac{81}{x} + \frac{9}{x}}{1 - \frac{81}{x} \cdot \frac{9}{x}} = \frac{90}{x} \cdot \frac{x^2}{x^2 - 81 \times 9}.$$

$$117x^2 - 81 \times 9 \times 117 = 90x^2;$$


$$\therefore 27x^2 = 81 \times 9 \times 117;$$

$$\therefore x^2 = 81 \times 39;$$

$$\therefore x = 9\sqrt{39}.$$

But  $\sqrt{39} = 6.245$  nearly;  $\therefore x = 56.205$  nearly.

Thus  $AB = 56.2$  ft. nearly.



9. A flagstaff 20 ft. long standing on a wall 10 ft. high subtends an angle whose tangent is  $\cdot 5$  at a point on the ground: find the tangent of the angle subtended by the wall at this point.

10. A statue standing on the top of a pillar 25 feet high subtends an angle whose tangent is  $\cdot 125$  at a point 60 feet from the foot of the pillar: find the height of the statue.

11. A tower  $BCD$  surmounted by a spire  $DE$  stands on a horizontal plane. From the extremity  $A$  of a horizontal line  $BA$  it is found that  $BC$  and  $DE$  subtend equal angles.

If  $BC=9$  ft.,  $CD=280$  ft., and  $DE=35$  ft., prove that  $BA=180$  ft. nearly.

12. On the bank of a river there is a column 192 ft. high supporting a statue 24 ft. high. At a point on the opposite bank directly facing the column the statue subtends the same angle as a man 6 ft. high standing at the base of the column: find the breadth of the river.

13. A monument  $ABCDE$  stands on level ground. At a point  $P$  on the ground the portions  $AB, AC, AD$  subtend angles  $\alpha, \beta, \gamma$  respectively. Supposing that  $AB=a, AC=b, AD=c, AP=x$ , and  $\alpha+\beta+\gamma=180^\circ$ , shew that  $(a+b+c)x^2=abc$ .

**Example IV.** The altitude of a rock is observed to be  $47^\circ$ ; after walking 1000 ft. towards it up a slope inclined at  $32^\circ$  to the horizon the altitude is  $77^\circ$ . Find the vertical height of the rock above the first point of observation, given  $\sin 47^\circ = .731$ .

Let  $P$  be the top of the rock,  $A$  and  $B$  the points of observation; then in the figure  $\angle PAC=47^\circ, \angle BAC=32^\circ$ ,

$\angle PDC = \angle PBE = 77^\circ, AB=1000$  ft.

Let  $x$  ft. be the height; then

$$x = PA \sin PAC = PA \sin 47^\circ.$$

We have therefore to find  $PA$  in terms of  $AB$ .

In  $\triangle PAB$ ,  $\angle PAB = 47^\circ - 32^\circ = 15^\circ$ ;

$$\angle APB = 77^\circ - 47^\circ = 30^\circ;$$

$$\therefore \angle ABP = 135^\circ;$$

$$\therefore PA = \frac{AB \sin ABP}{\sin APB}$$


$$= \frac{1000 \sin 135^\circ}{\sin 30^\circ}$$

$$= 1000 \sqrt{2};$$

$$\therefore x = PA \sin 47^\circ = 1000 \sqrt{2} \times .731$$

$$= 731 \sqrt{2}.$$

If we take  $\sqrt{2} = 1.414$ , we find that the height is 1034 ft. nearly.



14. From a point on the horizontal plane, the elevation of the top of a hill is  $45^\circ$ . After walking 500 yards towards its summit up a slope inclined at an angle of  $15^\circ$  to the horizon the elevation is  $75^\circ$ : find the height of the hill in feet.

15. From a station  $B$  at the base of a mountain its summit  $A$  is seen at an elevation of  $60^\circ$ ; after walking one mile towards the summit up a plane making an angle of  $30^\circ$  with the horizon to another station  $C$ , the angle  $BCA$  is observed to be  $135^\circ$ : find the height of the mountain in feet.

16. The elevation of the summit of a hill from a station  $A$  is  $\alpha$ . After walking  $c$  feet towards the summit up a slope inclined at an angle  $\beta$  to the horizon the elevation is  $\gamma$ : shew that the height of the hill is  $c \sin \alpha \sin (\gamma - \beta) \operatorname{cosec} (\gamma - \alpha)$  feet.

17. From a point  $A$  an observer finds that the elevation of Ben Nevis is  $60^\circ$ ; he then walks 800 ft. on a level plane towards the summit and then 800 ft. further up a slope of  $30^\circ$  and finds the elevation to be  $75^\circ$ : shew that the height of Ben Nevis above  $A$  is 4478 ft. approximately.

200. In many of the problems which follow, the solution depends upon the knowledge of some geometrical proposition.

**Example I.** A tower stands on a horizontal plane. From a mound 14 ft. above the plane and at a horizontal distance of 48 ft. from the tower an observer notices a loophole, and finds that the portions of the tower above and below the loophole subtend equal angles. If the height of the loophole is 30 ft., find the height of the tower.

Let  $AB$  be the tower,  $C$  the point of observation,  $L$  the loophole. Draw  $CD$  vertical and  $CE$  horizontal. Let  $AB = x$ . We have

$$CD = 14, AD = EC = 48, BE = x - 14.$$

$$\text{From } \triangle ADC, AC^2 = (14)^2 + (48)^2 = 2500;$$

$$\therefore AC = 50.$$

$$\begin{aligned} \text{From } \triangle CEB, CB^2 &= (x - 14)^2 + (48)^2 \\ &= x^2 - 28x + 2500. \end{aligned}$$

$$\text{Now } \angle BCL = \angle ACL;$$


$$\text{hence by Euc. vi. 3, } \frac{BC}{AC} = \frac{BL}{AL};$$

$$\therefore \frac{\sqrt{x^2 - 28x + 2500}}{50} = \frac{x - 30}{30}.$$

$$\text{By squaring, } 9(x^2 - 28x + 2500) = 25(x^2 - 60x + 900).$$

$$\text{On reduction, we obtain } 16x^2 - 1248x = 0; \text{ whence } x = 78.$$

Thus the tower is 78 ft. high.



## EXAMPLES. XVII. b.

1. At one side of a road is a flagstaff 25 ft. high fixed on the top of a wall 15 ft. high. On the other side of the road at a point on the ground directly opposite the flagstaff and wall subtend equal angles: find the width of the road.

2. A statue  $a$  feet high stands on a column  $3a$  feet high. To an observer on a level with the top of the statue, the column and statue subtend equal angles: find the distance of the observer from the top of the statue.

3. A flagstaff  $a$  feet high placed on the top of a tower  $b$  feet high subtends the same angle as the tower to an observer  $h$  feet high standing on the horizontal plane at a distance  $d$  feet from the foot of the tower: shew that

$$(a-b)d^2 = (a+b)b^2 - 2b^2h - (a-b)h^2.$$

**Example II.** A flagstaff is fixed on the top of a wall standing upon a horizontal plane. An observer finds that the angles subtended at a point on this plane by the wall and the flagstaff are  $\alpha$  and  $\beta$ . He then walks a distance  $c$  directly towards the wall and finds that the flagstaff again subtends an angle  $\beta$ . Find the heights of the wall and flagstaff.

Let  $ED$  be the wall,  $DC$  the flagstaff,  $A$  and  $B$  the points of observation.

Then  $\angle CAD = \beta = \angle CBD$ , so that the four points  $C, A, B, D$  are concyclic.

$$\begin{aligned}\therefore \angle ABD &= \text{suppl. of } \angle ACE \\ &= 90^\circ + (\alpha + \beta), \text{ from } \triangle CAE.\end{aligned}$$

Hence in  $\triangle ADB$ ,

$$\begin{aligned}\angle ADB &= 180^\circ - \alpha - \{90^\circ + (\alpha + \beta)\} \\ &= 90^\circ - (2\alpha + \beta).\end{aligned}$$


$$\therefore AD = \frac{AB \sin \angle ABD}{\sin \angle ADB} = \frac{c \cos (\alpha + \beta)}{\cos (2\alpha + \beta)}.$$

Hence in  $\triangle ADE$ ,

$$DE = AD \sin \alpha = \frac{c \sin \alpha \cos (\alpha + \beta)}{\cos (2\alpha + \beta)}.$$

And in  $\triangle CAD$ ,

$$CD = \frac{AD \sin \angle CAD}{\sin \angle ACD} = \frac{AD \sin \beta}{\cos (\alpha + \beta)} = \frac{c \sin \beta}{\cos (2\alpha + \beta)}.$$



4. A tower standing on a cliff subtends an angle  $a$  at each of two stations in the same horizontal line passing through the base of the cliff and at distances of  $a$  feet and  $b$  feet from the cliff. Prove that the height of the tower is  $(a+b) \tan a$  feet.

5. A column placed on a pedestal 20 feet high subtends an angle of  $45^\circ$  at a point on the ground, and it also subtends an angle of  $45^\circ$  at a point which is 20 feet nearer the pedestal: find the height of the column.

6. A flagstaff on a tower subtends the same angle at each of two places  $A$  and  $B$  on the ground. The elevations of the top of the flagstaff as seen from  $A$  and  $B$  are  $a$  and  $\beta$  respectively. If  $AB=a$ , shew that the length of the flagstaff is


$$a \sin(a+\beta-90^\circ) \operatorname{cosec}(a-\beta).$$

**Example III.** A man walking towards a tower  $AB$  on which a flagstaff  $BC$  is fixed observes that when he is at a point  $E$ , distant  $c$  ft. from the tower, the flagstaff subtends its greatest angle. If  $\angle BEC=a$ , prove that the heights of the tower and flagstaff are  $c \tan\left(\frac{\pi}{4}-\frac{a}{2}\right)$  and  $2c \tan a$  ft. respectively.

Since  $E$  is the point in the horizontal line  $AE$  at which  $BC$  subtends a maximum angle, it can easily be proved that  $AE$  touches the circle passing round the triangle  $CBE$ .

[See Hall and Stevens' *Euclid*, p. 242.]

The centre  $D$  of this circle lies in the vertical line through  $E$ . Draw  $DF$  perpendicular to  $BC$ , then  $DF$  bisects  $BC$  and also  $\angle CDB$ .



By Euc. III. 20,

$$\angle CDB = 2 \angle CEB = 2a;$$

$$\therefore \angle CDF = \angle BDF = a.$$

$$\therefore CB = 2CF = 2DF \tan a = 2c \tan a.$$

Again,  $\angle AEB = \angle ECB$  in alternate segment

$$= \frac{1}{2} \angle EDB \text{ at centre}$$

$$= \frac{1}{2} \left( \frac{\pi}{2} - a \right).$$

$$\therefore AB = c \tan AEB = c \tan \left( \frac{\pi}{4} - \frac{a}{2} \right).$$



7. A pillar stands on a pedestal. At a distance of 60 feet from the base of the pedestal the pillar subtends its greatest angle  $30^\circ$ : shew that the length of the pillar is  $40\sqrt{3}$  feet, and that the pedestal also subtends  $30^\circ$  at the point of observation.

8. A person walking along a canal observes that two objects are in the same line which is inclined at an angle  $\alpha$  to the canal. He walks a distance  $c$  further and observes that the objects subtend their greatest angle  $\beta$ : shew that their distance apart is

$$2c \sin \alpha \sin \beta / (\cos \alpha + \cos \beta).$$

9. A tower with a flagstaff stands on a horizontal plane. Shew that the distances from the base at which the flagstaff subtends the same angle and that at which it subtends the greatest possible angle are in geometrical progression.

10. The line joining two stations  $A$  and  $B$  subtends equal angles at two other stations  $C$  and  $D$ : prove that

$$AB \sin CBD = CD \sin ADB.$$

11. Two straight lines  $ABC$ ,  $DEC$  meet at  $C$ . If

$$\angle DAE = \angle DBE = \alpha, \text{ and } \angle EAB = \beta, \angle EBC = \gamma,$$

shew that

$$BC = \frac{AB \sin \beta \sin (\alpha + \beta)}{\sin (\gamma - \beta) \sin (\alpha + \beta + \gamma)}.$$

12. Two objects  $P$  and  $Q$  subtend an angle of  $30^\circ$  at  $A$ . Lengths of 20 feet and 10 feet are measured from  $A$  at right angles to  $AP$  and  $AQ$  respectively to points  $R$  and  $S$  at each of which  $PQ$  subtends angles of  $30^\circ$ : find the length of  $PQ$ .

13. A ship sailing N.E. is in a line with two beacons which are 5 miles apart, and of which one is due N. of the other. In 3 minutes and also in 21 minutes the beacons are found to subtend a right angle at the ship. Prove that the ship is sailing at the rate of 10 miles an hour, and that the beacons subtend their greatest angle at the ship at the end of  $3\sqrt{7}$  minutes.

14. A man walking along a straight road notes when he is in the line of a long straight fence, and observes that 78 yards from this point the fence subtends an angle of  $60^\circ$ , and that 260 yards further on this angle is increased to  $120^\circ$ . When he has walked 260 yards still further, he finds that the fence again subtends an angle of  $60^\circ$ . If  $\alpha$  be the angle which the direction of the fence makes with the road, shew that  $13 \sin \alpha = 5$ . Also shew that the middle point of the fence is 120 yards distant from the road.



### Measurements in more than one plane.

201. In Art. 199 the base line  $AB$  was measured *directly towards* the object. If this is not possible we may proceed as follows.

From  $A$  measure a base line  $AB$  in any convenient direction in the horizontal plane. At  $A$  observe the two angles  $PAB$  and  $PAC$ ; and at  $B$  observe the angle  $PBA$ .

Let  $\angle PAB = \alpha$ ,  $\angle PAC = \beta$ ,

$\angle PBA = \gamma$ ,

$AB = a$ ,  $PC = x$ .


From  $\triangle PAC$ ,

$$x = PA \sin \beta.$$

From  $\triangle PAB$ ,

$$PA = \frac{AB \sin PBA}{\sin APB} = \frac{a \sin \gamma}{\sin (a + \gamma)};$$

$$\therefore x = a \sin \beta \sin \gamma \operatorname{cosec} (a + \gamma).$$



202. To shew how to find the distance between two inaccessible objects.

Let  $P$  and  $Q$  be the objects.

Take any two convenient stations  $A$  and  $B$  in the same horizontal plane, and measure the distance between them.


At  $A$  observe the angles  $PAQ$  and  $QAB$ . Also if  $AP$ ,  $AQ$ ,  $AB$  are not in the same plane, measure the angle  $PAB$ .

At  $B$  observe the angles  $ABP$  and  $ABQ$ .


In  $\triangle PAB$ , we know  $\angle PAB$ ,  $\angle PBA$ , and  $AB$ ;  
so that  $AP$  may be found.

In  $\triangle QAB$ , we know  $\angle QAB$ ,  $\angle QBA$ , and  $AB$ ;  
so that  $AQ$  may be found.

In  $\triangle PAQ$ , we know  $AP$ ,  $AQ$ , and  $\angle PAQ$ ;  
so that  $PQ$  may be found.



*Example 1.* The angular elevation of a tower  $CD$  at a place  $A$  due South of it is  $30^\circ$ , and at a place  $B$  due West of  $A$  the elevation is  $18^\circ$ . If  $AB=a$ , shew that the height of the tower is  $\frac{a}{\sqrt{2+2\sqrt{5}}}$ .



Let  $CD=x$ .

From the right-angled triangle  $DCA$ ,  $AC=x \cot 30^\circ$ .

From the right-angled triangle  $DCB$ ,  $BC=x \cot 18^\circ$ .

But  $\angle BAC$  is a right angle,

$$\therefore BC^2 - AC^2 = a^2;$$

$$\therefore x^2 (\cot^2 18^\circ - \cot^2 30^\circ) = a^2;$$

$$\therefore x^2 (\operatorname{cosec}^2 18^\circ - \operatorname{cosec}^2 30^\circ) = a^2;$$

$$\therefore x^2 \left\{ \left( \frac{4}{\sqrt{5}-1} \right)^2 - 4 \right\} = a^2;$$

$$\therefore x^2 \{ (\sqrt{5}+1)^2 - 4 \} = a^2;$$


$$\therefore x^2 (2 \times 2\sqrt{5}) = a^2,$$

which gives the height required.

*Example 2.* A hill of inclination 1 in 5 faces South. Shew that a road on it which takes a N.E. direction has an inclination 1 in 7.

Let  $AD$  running East and West be the ridge of the hill, and let  $ABFD$  be a vertical plane through  $AD$ . Let  $C$  be a point at the foot of the hill, and  $ABC$  a section made by a vertical plane running North and South. Draw  $CG$  in a N.E. direction in the horizontal plane and let it meet  $BF$  in  $G$ ; draw  $GH$  parallel to  $BA$ ; then if  $CH$  is joined it will represent the direction of the road.

Since the inclination of  $CA$  is 1 in 5, we may take  $AB=a$ , and  $AC=5a$ , so that  $BC^2=24a^2$ .



Since  $CBG$  is a right-angled isosceles triangle,

$$CG^2 = 2CB^2 = 48a^2.$$

Hence in the right-angled triangle  $CGH$ ,

$$CH^2 = 48a^2 + a^2 = 49a^2;$$

$$\therefore CH = 7a = 7GH.$$

Thus the slope of the road is 1 in 7.

### EXAMPLES. XVII. c.

1. The elevation of a hill at a place  $P$  due East of it is  $45^\circ$ , and at a place  $Q$  due South of  $P$  the elevation is  $30^\circ$ . If the distance from  $P$  to  $Q$  is 500 yards, find the height of the hill in feet.

2. The elevation of a spire at a point  $A$  due West of it is  $60^\circ$ , and at point  $B$  due South of  $A$  the elevation is  $30^\circ$ . If the spire is 250 feet high, find the distance between  $A$  and  $B$ .

3. A river flows due North, and a tower stands on its left bank. From a point  $A$  up-stream and on the same bank as the tower the elevation of the tower is  $60^\circ$ , and from a point  $B$  just opposite on the other bank the elevation is  $45^\circ$ . If the tower is 360 feet high, find the breadth of the river.

4. The elevation of a steeple at a place  $A$  due S. of it is  $45^\circ$ , and at a place  $B$  due W. of  $A$  the elevation is  $15^\circ$ . If  $AB=2a$ , shew that the height of the steeple is  $a(3^{\frac{1}{2}} - 3^{-\frac{1}{2}})$ .

5. A person due S. of a lighthouse observes that his shadow cast by the light at the top is 24 feet long. On walking 100 yards due E. he finds his shadow to be 30 feet long. Supposing him to be 6 feet high, find the height of the light from the ground.

6. The angles of elevation of a balloon from two stations a mile apart and from a point halfway between them are observed to be  $60^\circ$ ,  $30^\circ$ , and  $45^\circ$  respectively. Prove that the height of the balloon is  $440\sqrt{6}$  yards.

[If  $AD$  is a median of the triangle  $ABC$ ,  
then  $2AD^2 + 2BD^2 = AB^2 + AC^2$ .]

7. At each end of a base of length  $2a$ , the angular elevation of a mountain is  $\theta$ , and at the middle point of the base the elevation is  $\phi$ . Prove that the height of the mountain is

$$a \sin \theta \sin \phi \sqrt{\operatorname{cosec}(\phi + \theta) \operatorname{cosec}(\phi - \theta)}.$$

8. Two vertical poles, whose heights are  $a$  and  $b$ , subtend the same angle  $\alpha$  at a point in the line joining their feet. If they subtend angles  $\beta$  and  $\gamma$  at any point in the horizontal plane at which the line joining their feet subtends a right angle, prove that

$$(a+b)^2 \cot^2 \alpha = a^2 \cot^2 \beta + b^2 \cot^2 \gamma.$$

9. From the top of a hill a person finds that the angles of depression of three consecutive milestones on a straight level road are  $\alpha$ ,  $\beta$ ,  $\gamma$ . Shew that the height of the hill is

$$5280\sqrt{2} / \sqrt{\cot^2 \alpha - 2 \cot^2 \beta + \cot^2 \gamma} \text{ feet.}$$

10. Two chimneys  $AB$  and  $CD$  are of equal height. A person standing between them in the line  $AC$  joining their bases observes the elevation of the one nearer to him to be  $60^\circ$ . After walking 80 feet in a direction at right angles to  $AC$  he observes their elevations to be  $45^\circ$  and  $30^\circ$ : find their height and distance apart.

11. Two persons who are 500 yards apart observe the bearing and angular elevation of a balloon at the same instant. One finds the elevation  $60^\circ$  and the bearing S.W., the other finds the elevation  $45^\circ$  and the bearing W. Find the height of the balloon.

12. The side of a hill faces due S. and is inclined to the horizon at an angle  $\alpha$ . A straight railway upon it is inclined at an angle  $\beta$  to the horizon: if the bearing of the railway be  $x$  degrees E. of N., shew that  $\cos x = \cot \alpha \tan \beta$ .

## EXAMPLES. XVII. d.

[In the following examples the logarithms are to be taken from the Tables.]

1. A man in a balloon observes that two churches which he knows to be one mile apart subtend an angle of  $11^{\circ} 25' 20''$  when he is exactly over the middle point between them : find the height of the balloon in miles.

2. There are three points  $A, B, C$  in a straight line on a level piece of ground. A vertical pole erected at  $C$  has an elevation of  $5^{\circ} 30'$  from  $A$  and  $10^{\circ} 45'$  from  $B$ . If  $AB$  is 100 yards, find the height of the pole and the distance  $BC$ .

3. The angular altitude of a lighthouse seen from a point on the shore is  $12^{\circ} 31' 46''$ , and from a point 500 feet nearer the altitude is  $26^{\circ} 33' 55''$  : find its height above the sea-level.

4. From a boat the angles of elevation of the highest and lowest points of a flagstaff 30 ft. high on the edge of a cliff are  $46^{\circ} 12'$  and  $44^{\circ} 13'$  : find the height and distance of the cliff.

5. From the top of a hill the angles of depression of two successive milestones on level ground, and in the same vertical plane as the observer, are  $5^{\circ}$  and  $10^{\circ}$ . Find the height of the hill in feet and the distance of the nearer milestone in miles.

6. An observer whose eye is 15 feet above the roadway finds that the angle of elevation of the top of a telegraph post is  $17^{\circ} 18' 35''$ , and that the angle of depression of the foot of the post is  $8^{\circ} 32' 15''$  : find the height of the post and its distance from the observer.

7. Two straight railroads are inclined at an angle of  $20^{\circ} 16'$ . At the same instant two engines start from the point of intersection, one along each line ; one travels at the rate of 20 miles an hour : at what rate must the other travel so that after 3 hours the distance between them shall be 30 miles ?

8. An observer finds that from the doorstep of his house the elevation of the top of a spire is  $5a$ , and that from the roof above the doorstep it is  $4a$ . If  $h$  be the height of the roof above the doorstep, prove that the height of the spire above the doorstep and the horizontal distance of the spire from the house are respectively

$$h \operatorname{cosec} a \cos 4a \sin 5a \text{ and } h \operatorname{cosec} a \cos 4a \cos 5a.$$

If  $h=39$  feet, and  $a=7^{\circ} 17' 39''$ , calculate the height and the distance.




## CHAPTER XVIII.

### PROPERTIES OF TRIANGLES AND POLYGONS.

203. *To find the area of a triangle.*

Let  $\Delta$  denote the area of the triangle  $ABC$ . Draw  $AD$  perpendicular to  $BC$ .

By Euc. I. 41, the area of a triangle is half the area of a rectangle on the same base and of the same altitude.



$$\begin{aligned}\therefore \Delta &= \frac{1}{2} (\text{base} \times \text{altitude}) \\ &= \frac{1}{2} BC \cdot AD = \frac{1}{2} BC \cdot AB \sin B \\ &= \frac{1}{2} ca \sin B.\end{aligned}$$

Similarly, it may be proved that

$$\Delta = \frac{1}{2} ab \sin C, \text{ and } \Delta = \frac{1}{2} bc \sin A.$$

These three expressions for the area are comprised in the single statement

$$\Delta = \frac{1}{2} (\text{product of two sides}) \times (\text{sine of included angle}).$$

$$\text{Again, } \Delta = \frac{1}{2} bc \sin A = bc \sin \frac{A}{2} \cos \frac{A}{2}$$

$$\begin{aligned}&= bc \sqrt{\frac{(s-b)(s-c)}{bc}} \sqrt{\frac{s(s-a)}{bc}} \\ &= \sqrt{s(s-a)(s-b)(s-c)},\end{aligned}$$

which gives the area in terms of the sides.



$$\begin{aligned}
 \text{Again, } \Delta &= \frac{1}{2} bc \sin A = \frac{1}{2} \sin A \cdot \frac{a \sin B}{\sin A} \cdot \frac{a \sin C}{\sin A} \\
 &= \frac{a^2 \sin B \sin C}{2 \sin A} \\
 &= \frac{a^2 \sin B \sin C}{2 \sin (B+C)},
 \end{aligned}$$

which gives the area in terms of one side and the functions of the adjacent angles.

NOTE. Many writers use the symbol  $S$  for the area of a triangle, but to avoid confusion between  $S$  and  $s$  in manuscript work the symbol  $\Delta$  is preferable.

*Example 1.* The sides of a triangle are 17, 25, 28: find the lengths of the perpendiculars from the angles upon the opposite sides.

$$\text{From the formula } \Delta = \frac{1}{2} (\text{base} \times \text{altitude}),$$

it is evident that the three perpendiculars are found by dividing  $2\Delta$  by the three sides in turn.

$$\begin{aligned}
 \text{Now } \Delta &= \sqrt{s(s-a)(s-b)(s-c)} = \sqrt{35 \times 18 \times 10 \times 7} \\
 &= 5 \times 7 \times 6 = 210.
 \end{aligned}$$

Thus the perpendiculars are  $\frac{420}{17}$ ,  $\frac{420}{25}$ ,  $\frac{420}{28}$ , or  $\frac{420}{17}$ ,  $\frac{84}{5}$ , 15.

*Example 2.* Two angles of a triangular field are  $22.5^\circ$  and  $45^\circ$ , and the length of the side opposite to the latter is one furlong: find the area.

Let  $A = 22\frac{1}{2}^\circ$ ,  $B = 45^\circ$ , then  $b = 220$  yds., and  $C = 112\frac{1}{2}^\circ$ .

$$\text{From the formula } \Delta = \frac{b^2 \sin A \sin C}{2 \sin B},$$

$$\begin{aligned}
 \text{the area in sq. yds.} &= \frac{220 \times 220 \times \sin 22\frac{1}{2}^\circ \times \sin 112\frac{1}{2}^\circ}{2 \sin 45^\circ} \\
 &= \frac{220 \times 220 \times \sin 22\frac{1}{2}^\circ \times \cos 22\frac{1}{2}^\circ}{2 \times 2 \sin 22\frac{1}{2}^\circ \cos 22\frac{1}{2}^\circ} \\
 &= 110 \times 110.
 \end{aligned}$$


$$\text{Expressed in acres, the area} = \frac{110 \times 110}{4840} = 2\frac{1}{2}.$$

204. To find the radius of the circle circumscribing a triangle.

Let  $S$  be the centre of the circle circumscribing the triangle  $ABC$ , and  $R$  its radius.

Bisect  $\angle BSC$  by  $SD$ , which will also bisect  $BC$  at right angles.

Now by Euc. III. 20,  
 $\angle BSC$  at centre  
 $= \text{twice } \angle BAC$   
 $= 2A$ ;



and  $\frac{a}{2} = BD = BS \sin BSD = R \sin A$ ;

$$\therefore R = \frac{a}{2 \sin A}.$$

Thus  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$ ,

or  $a = 2R \sin A$ ,  $b = 2R \sin B$ ,  $c = 2R \sin C$ .

*Example.* Shew that  $2R^2 \sin A \sin B \sin C = \Delta$ .

$$\begin{aligned} \text{The first side} &= \frac{1}{2} \cdot 2R \sin A \cdot 2R \sin B \cdot \sin C \\ &= \frac{1}{2} ab \sin C \\ &= \Delta. \end{aligned}$$

205. From the result of the last article we deduce the following important theorem:

*If a chord of length  $l$  subtend an angle  $\theta$  at the circumference of a circle whose radius is  $R$ , then  $l = 2R \sin \theta$ .*


206. For shortness, the circle circumscribing a triangle may be called the *Circum-circle*, its centre the *Circum-centre*, and its radius the *Circum-radius*.

The circum-radius may be expressed in a form not involving the angles, for

$$R = \frac{a}{2 \sin A} = \frac{abc}{2bc \sin A} = \frac{abc}{4\Delta}.$$

207. *To find the radius of the circle inscribed in a triangle.*

Let  $I$  be the centre of the circle inscribed in the triangle  $ABC$ , and  $D, E, F$  the points of contact; then  $ID, IE, IF$  are perpendicular to the sides.



Now  $\Delta$  = sum of the areas of the triangles  $BIC, CIA, AIB$

$$= \frac{1}{2} ar + \frac{1}{2} br + \frac{1}{2} cr = \frac{1}{2} (a+b+c) r$$

$$= sr ;$$

whence

$$r = \frac{\Delta}{s}.$$

208. *To express the radius of the inscribed circle in terms of one side and the functions of the half-angles.*

In the figure of the previous article, we know from Euc. iv. 4 that  $I$  is the point of intersection of the lines bisecting the angles, so that

$$\angle IBD = \frac{B}{2}, \quad \angle ICD = \frac{C}{2}.$$

$$\therefore BD = r \cot \frac{B}{2}, \quad CD = r \cot \frac{C}{2}.$$

$$\therefore r \left( \cot \frac{B}{2} + \cot \frac{C}{2} \right) = a ;$$

$$\therefore r \sin \frac{B+C}{2} = a \sin \frac{B}{2} \sin \frac{C}{2} ;$$

$$\therefore r = \frac{a \sin \frac{B}{2} \sin \frac{C}{2}}{\cos \frac{A}{2}}.$$

**209. DEFINITION.** A circle which touches one side of a triangle and the other two sides produced is said to be an **escribed circle** of the triangle.

Thus the triangle  $ABC$  has *three* escribed circles, one touching  $BC$ , and  $AB$ ,  $AC$  produced; a second touching  $CA$ , and  $BC$ ,  $BA$  produced; a third touching  $AB$ , and  $CA$ ,  $CB$  produced.

We shall assume that the student is familiar with the construction of the escribed circles.

[See Hall and Stevens' *Euclid*, p. 255.]

For shortness, we shall call the circle inscribed in a triangle the *In-circle*, its centre the *In-centre*, and its radius the *In-radius*; and similarly the escribed circles may be called the *Ex-circles*, their centres the *Ex-centres*, and their radii the *Ex-radii*.

**210.** To find the radius of an escribed circle of a triangle.

Let  $I_1$  be the centre of the circle touching the side  $BC$  and the two sides  $AB$  and  $AC$  produced. Let  $D_1$ ,  $E_1$ ,  $F_1$  be the points of contact; then the lines joining  $I_1$  to these points are perpendicular to the sides.


Let  $r_1$  be the radius; then

$$\begin{aligned}\Delta &= \text{area } ABC \\ &= \text{area } ABI_1C - \text{area } BI_1C \\ &= \text{area } BI_1A + \text{area } CI_1A \\ &\quad - \text{area } BI_1C \\ &= \frac{1}{2} cr_1 + \frac{1}{2} br_1 - \frac{1}{2} ar_1 \\ &= \frac{1}{2} (c + b - a) r_1 \\ &= (s - a) r_1;\end{aligned}$$

$$\therefore r_1 = \frac{\Delta}{s - a}.$$

Similarly, if  $r_2$ ,  $r_3$  be the radii of the escribed circles opposite to the angles  $B$  and  $C$  respectively,

$$r_2 = \frac{\Delta}{s - b}, \quad r_3 = \frac{\Delta}{s - c}.$$



**211.** *To find the radii of the escribed circles in terms of one side and the functions of the half-angles.*

In the figure of the last article,  $I_1$  is the point of intersection of the lines bisecting the angles  $B$  and  $C$  externally; so that

$$\angle I_1 B D_1 = 90^\circ - \frac{B}{2}, \quad \angle I_1 C D_1 = 90^\circ - \frac{C}{2}.$$

$$\therefore BD_1 = r_1 \cot \left( 90^\circ - \frac{B}{2} \right) = r_1 \tan \frac{B}{2},$$

$$CD_1 = r_1 \cot \left( 90^\circ - \frac{C}{2} \right) = r_1 \tan \frac{C}{2};$$

$$\therefore r_1 \left( \tan \frac{B}{2} + \tan \frac{C}{2} \right) = a;$$

$$\therefore r_1 \sin \frac{B+C}{2} = a \cos \frac{B}{2} \cos \frac{C}{2};$$

$$\therefore r_1 = \frac{a \cos \frac{B}{2} \cos \frac{C}{2}}{\cos \frac{A}{2}}.$$

Similarly,

$$r_2 = \frac{b \cos \frac{C}{2} \cos \frac{A}{2}}{\cos \frac{B}{2}}, \quad r_3 = \frac{c \cos \frac{A}{2} \cos \frac{B}{2}}{\cos \frac{C}{2}}.$$

**212.** By substituting

$$a = 2R \sin A, \quad b = 2R \sin B, \quad c = 2R \sin C,$$

in the formulæ of Art. 208 and Art. 211, we have

$$r = 4R \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2},$$

$$r_1 = 4R \sin \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2},$$

$$r_2 = 4R \cos \frac{A}{2} \sin \frac{B}{2} \cos \frac{C}{2},$$

$$r_3 = 4R \cos \frac{A}{2} \cos \frac{B}{2} \sin \frac{C}{2}.$$

*Example 1.* Shew that  $\frac{r_1 - r}{a} + \frac{r_2 - r}{b} = \frac{c}{r_3}$ .

$$\begin{aligned} \text{The first side} &= \frac{1}{a} \left( \frac{\Delta}{s-a} - \frac{\Delta}{s} \right) + \frac{1}{b} \left( \frac{\Delta}{s-b} - \frac{\Delta}{s} \right) \\ &= \frac{\Delta}{s(s-a)} + \frac{\Delta}{s(s-b)} = \frac{\Delta(2s-a-b)}{s(s-a)(s-b)} \\ &= \frac{\Delta c}{s(s-a)(s-b)} = \frac{\Delta(s-c)}{s(s-a)(s-b)(s-c)} \\ &= \frac{\Delta c(s-c)}{\Delta^2} = \frac{c(s-c)}{\Delta} \\ &= \frac{c}{r_3}. \end{aligned}$$

*Example 2.* If  $r_1 = r_2 + r_3 + r$ , prove that the triangle is right-angled.

By transposition,  $r_1 - r = r_2 + r_3$ ;

$$\begin{aligned} \therefore 4R \sin \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2} - 4R \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} \\ &= 4R \cos \frac{A}{2} \sin \frac{B}{2} \cos \frac{C}{2} + 4R \cos \frac{A}{2} \cos \frac{B}{2} \sin \frac{C}{2}; \\ \therefore \sin \frac{A}{2} \left( \cos \frac{B}{2} \cos \frac{C}{2} - \sin \frac{B}{2} \sin \frac{C}{2} \right) \\ &= \cos \frac{A}{2} \left( \sin \frac{B}{2} \cos \frac{C}{2} + \cos \frac{B}{2} \sin \frac{C}{2} \right); \\ \therefore \sin \frac{A}{2} \cos \frac{B+C}{2} &= \cos \frac{A}{2} \sin \frac{B+C}{2}; \\ \therefore \sin^2 \frac{A}{2} &= \cos^2 \frac{A}{2}; \end{aligned}$$

whence  $\frac{A}{2} = 45^\circ$ , and  $A = 90^\circ$ .

**213.** Many important relations connecting a triangle and its circles may be established by elementary geometry.

With the notation of previous articles, since tangents to a circle from the same point are equal,




we have  $AF = AE$ ,  $BD = BF$ ,  $CD = CE$ ;

$\therefore AF + (BD + CD) = \text{half the sum of the sides};$

$$\therefore AF + a = s.$$

$$\therefore AF = s - a = AE.$$

Similarly,  $BD = BF = s - b$ ,  $CD = CE = s - c$ .



Also 
$$r = AF \tan \frac{A}{2} = (s - a) \tan \frac{A}{2}.$$

Similarly, 
$$r = (s - b) \tan \frac{B}{2}, \quad r = (s - c) \tan \frac{C}{2}.$$

Again,  $AF_1 = AE_1$ ,  $BF_1 = BD_1$ ,  $CE_1 = CD_1$ ;

$$\therefore 2AF_1 = AF_1 + AE_1 = (AB + BD_1) + (AC + CD_1) \\ = \text{sum of the sides};$$

$$\therefore AF_1 = s = AE_1.$$

$$\therefore BD_1 = BF_1 = s - c, \quad CD_1 = CE_1 = s - b.$$

Also 
$$r_1 = AF_1 \tan \frac{A}{2} = s \tan \frac{A}{2}.$$

Similarly, 
$$r_2 = s \tan \frac{B}{2}, \quad r_3 = s \tan \frac{C}{2}.$$

**EXAMPLES. XVIII. a.**

1. Two sides of a triangle are 300 ft. and 120 ft., and the included angle is  $150^\circ$ ; find the area.
2. Find the area of the triangle whose sides are 171, 204, 195.
3. Find the sine of the greatest angle of a triangle whose sides are 70, 147, and 119.
4. If the sides of a triangle are 39, 40, 25, find the lengths of the three perpendiculars from the angular points on the opposite sides.
5. One side of a triangle is 30 ft. and the adjacent angles are  $22\frac{1}{2}^\circ$  and  $112\frac{1}{2}^\circ$ , find the area.
6. Find the area of a parallelogram two of whose adjacent sides are 42 and 32 ft., and include an angle of  $30^\circ$ .
7. The area of a rhombus is 648 sq. yds. and one of the angles is  $150^\circ$ : find the length of each side.
8. In a triangle if  $a=13$ ,  $b=14$ ,  $c=15$ , find  $r$  and  $R$ .
9. Find  $r_1$ ,  $r_2$ ,  $r_3$  in the case of a triangle whose sides are 17, 10, 21.
10. If the area of a triangle is 96, and the radii of the escribed circles are 8, 12, 24, find the sides.

Prove the following formulæ :

- |  |  |
|--|--|
| 11. $\sqrt{rr_1r_2r_3} = \Delta.$            | 12. $s(s-a) \tan \frac{A}{2} = \Delta.$              |
| 13. $rr_1 \cot \frac{A}{2} = \Delta.$        | 14. $4Rrs = abc.$                                    |
| 15. $r_1r_2r_3 = rs^2.$                      | 16. $r \cot \frac{B}{2} \cot \frac{C}{2} = r_1.$     |
| 17. $Rr(\sin A + \sin B + \sin C) = \Delta.$ |  |
| 18. $r_1r_2 + rr_3 = ab.$                    | 19. $\cos \frac{A}{2} \sqrt{bc(s-b)(s-c)} = \Delta.$ |
| 20. $r_1 + r_2 = c \cot \frac{C}{2}.$        | 21. $(r_1 - r)(r_2 + r_3) = a^2.$                    |

$$22. \quad r_1 \cot \frac{A}{2} = r_2 \cot \frac{B}{2} = r_3 \cot \frac{C}{2} = r \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2}.$$

$$23. \quad \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = \frac{1}{r}. \qquad 24. \quad r_2 r_3 + r_3 r_1 + r_1 r_2 = s^2.$$

$$25. \quad r_1 + r_2 + r_3 - r = 4R. \qquad 26. \quad r + r_1 + r_2 - r_3 = 4R \cos C.$$

$$27. \quad b^2 \sin 2C + c^2 \sin 2B = 4\Delta.$$

$$28. \quad 4R \cos \frac{C}{2} = (a+b) \sec \frac{A-B}{2}.$$

$$29. \quad a^2 - b^2 = 2Rc \sin (A-B).$$

$$30. \quad \frac{a^2 - b^2}{2} \cdot \frac{\sin A \sin B}{\sin (A-B)} = \Delta.$$

31. If the perpendiculars from  $A$ ,  $B$ ,  $C$  to the opposite sides are  $p_1$ ,  $p_2$ ,  $p_3$  respectively, prove that

$$(1) \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} = \frac{1}{r}; \qquad (2) \quad \frac{1}{p_1} + \frac{1}{p_2} - \frac{1}{p_3} = \frac{1}{r_3}.$$

Prove the following identities :

$$32. \quad (r_1 - r)(r_2 - r)(r_3 - r) = 4Rr^2.$$

$$33. \quad \left(\frac{1}{r} - \frac{1}{r_1}\right) \left(\frac{1}{r} - \frac{1}{r_2}\right) \left(\frac{1}{r} - \frac{1}{r_3}\right) = \frac{4R}{r^2 s^2}.$$

$$34. \quad 4\Delta (\cot A + \cot B + \cot C) = a^2 + b^2 + c^2.$$

$$35. \quad \frac{b-c}{r_1} + \frac{c-a}{r_2} + \frac{a-b}{r_3} = 0.$$

$$36. \quad a^2 b^2 c^2 (\sin 2A + \sin 2B + \sin 2C) = 32\Delta^3.$$

$$37. \quad a \cos A + b \cos B + c \cos C = 4R \sin A \sin B \sin C.$$

$$38. \quad a \cot A + b \cot B + c \cot C = 2(R + r).$$

$$39. \quad (b+c) \tan \frac{A}{2} + (c+a) \tan \frac{B}{2} + (a+b) \tan \frac{C}{2} \\ = 4R (\cos A + \cos B + \cos C).$$

$$40. \quad r (\sin A + \sin B + \sin C) = 2R \sin A \sin B \sin C.$$

$$41. \quad \cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} + \cos^2 \frac{C}{2} = 2 + \frac{r}{2R}.$$


### Inscribed and circumscribed Polygons.

**214.** *To find the perimeter and area of a regular polygon of  $n$  sides inscribed in a circle.*

Let  $r$  be the radius of the circle, and  $AB$  a side of the polygon.

Join  $OA$ ,  $OB$ , and draw  $OD$  bisecting  $\angle AOB$ ; then  $AB$  is bisected at right angles in  $D$ .

$$\begin{aligned}\text{And } \angle AOB &= \frac{1}{n} \text{ (four right angles)} \\ &= \frac{2\pi}{n}.\end{aligned}$$



$$\text{Perimeter of polygon} = nAB = 2nAD = 2nOA \sin AOD$$

$$= 2nr \sin \frac{\pi}{n}.$$

$$\text{Area of polygon} = n (\text{area of triangle } AOB)$$

$$= \frac{1}{2} nr^2 \sin \frac{2\pi}{n}.$$


**215.** *To find the perimeter and area of a regular polygon of  $n$  sides circumscribed about a given circle.*

Let  $r$  be the radius of the circle, and  $AB$  a side of the polygon. Let  $AB$  touch the circle at  $D$ . Join  $OA$ ,  $OB$ ,  $OD$ ; then  $OD$  bisects  $AB$  at right angles, and also bisects  $\angle AOB$ .

$$\text{Perimeter of polygon}$$

$$= nAB = 2nAD = 2nOD \tan AOD$$

$$= 2nr \tan \frac{\pi}{n}.$$



$$\text{Area of polygon} = n (\text{area of triangle } AOB)$$

$$= nOD \cdot AD$$

$$= nr^2 \tan \frac{\pi}{n}.$$

216. There is no need to burden the memory with the formulæ of the last two articles, as in any particular instance they are very readily obtained.

*Example 1.* The side of a regular dodecagon is 2 ft., find the radius of the circumscribed circle.

Let  $r$  be the required radius. In the adjoining figure we have


$$AB=2, \angle AOB=\frac{2\pi}{12}.$$

$$AB=2AD=2r \sin \frac{\pi}{12};$$

$$\therefore 2r \sin 15^\circ=2;$$

$$\therefore r=\frac{1}{\sin 15^\circ}=\frac{2\sqrt{2}}{\sqrt{3}-1}=\sqrt{2}(\sqrt{3}+1).$$

Thus the radius is  $\sqrt{6}+\sqrt{2}$  feet.



*Example 2.* A regular pentagon and a regular decagon have the same perimeter, prove that their areas are as 2 to  $\sqrt{5}$ .

Let  $AB$  be one of the  $n$  sides of a regular polygon,  $O$  the centre of the circumscribed circle,  $OD$  perpendicular to  $AB$ .

Then if  $AB=a$ ,

area of polygon  $=nAD \cdot OD$


$$=nAD \cdot AD \cot \frac{\pi}{n}$$

$$=\frac{na^2}{4} \cot \frac{\pi}{n}.$$

Denote the perimeter of the pentagon and decagon by  $10c$ . Then each side of the pentagon is  $2c$ , and its area is  $5c^2 \cot \frac{\pi}{5}$ .

Each side of the decagon is  $c$ , and its area is  $\frac{5}{2}c^2 \cot \frac{\pi}{10}$ .

$$\begin{aligned} \therefore \frac{\text{Area of pentagon}}{\text{Area of decagon}} &= \frac{2 \cot 36^\circ}{\cot 18^\circ} = \frac{2 \cos 36^\circ \sin 18^\circ}{\sin 36^\circ \cos 18^\circ} = \frac{2 \cos 36^\circ}{2 \cos^2 18^\circ} \\ &= \frac{2 \cos 36^\circ}{1 + \cos 36^\circ} = \frac{2(\sqrt{5}+1)}{4} \div \left(1 + \frac{\sqrt{5}+1}{4}\right) \\ &= \frac{2(\sqrt{5}+1)}{5+\sqrt{5}} = \frac{2}{\sqrt{5}}. \end{aligned}$$



217. *To find the area of a circle.*


Let  $r$  be the radius of the circle, and let a regular polygon of  $n$  sides be described about it. Then from the adjoining figure, we have

area of polygon  $= n$  (area of triangle  $AOB$ )

$$= n \left( \frac{1}{2} AB \cdot OD \right)$$

$$= \frac{1}{2} OD \cdot nAB$$

$$= \frac{r}{2} \times \text{perimeter of polygon.}$$



By increasing the number of sides without limit, the area and the perimeter of the polygon may be made to differ as little as we please from the area and the circumference of the circle. Hence

$$\begin{aligned} \text{area of a circle} &= \frac{r}{2} \times \text{circumference} \\ &= \frac{r}{2} \times 2\pi r \quad [\text{Art. 59.}] \\ &= \pi r^2. \end{aligned}$$

218. *To find the area of the sector of a circle.*

Let  $\theta$  be the circular measure of the angle of the sector; then by Euc. VI. 33,

$$\frac{\text{area of sector}}{\text{area of circle}} = \frac{\theta}{2\pi};$$

$$\therefore \text{area of sector} = \frac{\theta}{2\pi} \times \pi r^2 = \frac{1}{2} r^2 \theta.$$

**EXAMPLES. XVIII. b.**

$$\left[ \text{In this Exercise take } \pi = \frac{22}{7}. \right]$$

1. Find the area of a regular decagon inscribed in a circle whose radius is 3 feet; given  $\sin 36^\circ = .588$ .



2. Find the perimeter and area of a regular quindecagon described about a circle whose diameter is 3 yards ; given

$$\tan 12^\circ = \cdot 213.$$

3. Shew that the areas of the inscribed and circumscribed circles of a regular hexagon are in the ratio of 3 to 4.

4. Find the area of a circle inscribed in a regular pentagon whose area is 250 sq. ft. ; given  $\cot 36^\circ = 1\cdot376$ .

5. Find the perimeter of a regular octagon inscribed in a circle whose area is 1386 sq. inches ; given  $\sin 22^\circ 30' = \cdot382$ .

6. Find the perimeter of a regular pentagon described about a circle whose area is 616 sq. ft. ; given  $\tan 36^\circ = \cdot727$ .

7. Find the diameter of the circle circumscribing a regular quindecagon, whose inscribed circle has an area of 2464 sq. ft. ; given  $\sec 12^\circ = 1\cdot022$ .

8. Find the area of a regular dodecagon inscribed in a circle whose regular inscribed pentagon has an area of 50 sq. ft.

9. A regular pentagon and a regular decagon have the same area, prove that the ratio of their perimeters is  $\sqrt{2} : \sqrt[4]{5}$ .

10. Two regular polygons of  $n$  sides and  $2n$  sides have the same perimeter ; shew that the ratio of their areas is

$$2 \cos \frac{\pi}{n} : 1 + \cos \frac{\pi}{n}.$$

11. If  $2a$  be the side of a regular polygon of  $n$  sides,  $R$  and  $r$  the radii of the circumscribed and inscribed circles, prove that

$$R + r = a \cot \frac{\pi}{2n}.$$

12. Prove that the square of the side of a regular pentagon inscribed in a circle is equal to the sum of the squares of the sides of a regular hexagon and decagon inscribed in the same circle.


13. With reference to a given circle,  $A_1$  and  $B_1$  are the areas of the inscribed and circumscribed regular polygons of  $n$  sides,  $A_2$  and  $B_2$  are corresponding quantities for regular polygons of  $2n$  sides : prove that

(1)  $A_2$  is a geometric mean between  $A_1$  and  $B_1$  ;

(2)  $B_2$  is a harmonic mean between  $A_2$  and  $B_1$ .

### The Ex-central Triangle.

\*219. Let  $ABC$  be a triangle,  $I_1, I_2, I_3$  its ex-centres ; then  $I_1I_2I_3$  is called the **Ex-central triangle** of  $ABC$ .



Let  $I$  be the in-centre ; then from the construction for finding the positions of the in-centre and ex-centres, it follows that :

(i) The points  $I, I_1$  lie on the line bisecting the angle  $BAC$ ; the points  $I, I_2$  lie on the line bisecting the angle  $ABC$ ; the points  $I, I_3$  lie on the line bisecting the angle  $ACB$ .

(ii) The points  $I_2, I_3$  lie on the line bisecting the angle  $BAC$  externally ; the points  $I_3, I_1$  lie on the line bisecting the angle  $ABC$  externally ; the points  $I_1, I_2$  lie on the line bisecting the angle  $ACB$  externally.

(iii) The line  $AI_1$  is perpendicular to  $I_2I_3$ ; the line  $BI_2$  is perpendicular to  $I_3I_1$ ; the line  $CI_3$  is perpendicular to  $I_1I_2$ . Thus the triangle  $ABC$  is the *Pedal triangle* of its ex-central triangle  $I_1I_2I_3$ . [See Art. 223.]

(iv) The angles  $IBI_1$  and  $ICI_1$  are right angles ; hence the points  $B, I, C, I_1$  are concyclic. Similarly, the points  $C, I, A, I_2$ , and the points  $A, I, B, I_3$  are concyclic.

(v) The lines  $AI_1, BI_2, CI_3$  meet at the in-centre  $I$ , which is therefore the *Orthocentre* of the ex-central triangle  $I_1I_2I_3$ .

(vi) Each of the four points  $I, I_1, I_2, I_3$  is the orthocentre of the triangle formed by joining the other three points.

**\*220.** *To find the distances between the in-centre and ex-centres.*

With the figure of the last article,

$$\begin{aligned} II_1 &= AI_1 - AI = r_1 \operatorname{cosec} \frac{A}{2} - r \operatorname{cosec} \frac{A}{2} = (r_1 - r) \operatorname{cosec} \frac{A}{2} \\ &= 4R \left( \sin \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2} - \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} \right) \operatorname{cosec} \frac{A}{2} \\ &= 4R \cos \frac{B+C}{2} = 4R \sin \frac{A}{2}. \end{aligned}$$

Thus the distances are

$$4R \sin \frac{A}{2}, \quad 4R \sin \frac{B}{2}, \quad 4R \sin \frac{C}{2}.$$

**\*221.** *To find the sides and angles of the ex-central triangle.*

With the figure of Art. 219,

$$\begin{aligned} \angle BI_1C &= \angle BI_1I + \angle CI_1I \\ &= \angle BCI + \angle CBI && [\text{Euc. III. 21}] \\ &= \frac{C}{2} + \frac{B}{2} = 90^\circ - \frac{A}{2}. \end{aligned}$$

Thus the angles are

$$90^\circ - \frac{A}{2}, \quad 90^\circ - \frac{B}{2}, \quad 90^\circ - \frac{C}{2}.$$

Again,

$$\begin{aligned} I_1I_2 &= I_1C + I_2C = r_1 \operatorname{cosec} \left( 90^\circ - \frac{C}{2} \right) + r_2 \operatorname{cosec} \left( 90^\circ - \frac{C}{2} \right) \\ &= 4R \left( \sin \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2} + \cos \frac{A}{2} \sin \frac{B}{2} \cos \frac{C}{2} \right) \sec \frac{C}{2} \\ &= 4R \sin \frac{A+B}{2} = 4R \cos \frac{C}{2}. \end{aligned}$$

Thus the sides are

$$4R \cos \frac{A}{2}, \quad 4R \cos \frac{B}{2}, \quad 4R \cos \frac{C}{2}.$$

**\*222.** To find the area and circum-radius of the ex-central triangle.

The area =  $\frac{1}{2}$  (product of two sides)  $\times$  (sine of included angle)

$$= \frac{1}{2} \times 4R \cos \frac{B}{2} \times 4R \cos \frac{C}{2} \times \sin \left( 90^\circ - \frac{A}{2} \right)$$


$$= 8R^2 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}.$$

$$\text{The circum-radius} = \frac{I_2 I_3}{2 \sin I_2 I_1 I_3} = \frac{4R \cos \frac{A}{2}}{2 \sin \left( 90^\circ - \frac{A}{2} \right)} = 2R.$$

### The Pedal Triangle.

**\*223.** Let  $G, H, K$  be the feet of the perpendiculars from the angular points on the opposite sides of the triangle  $ABC$ ; then  $GHK$  is called the **Pedal triangle** of  $ABC$ .

The three perpendiculars  $AG, BH, CK$  meet in a point  $O$  which is called the **Orthocentre** of the triangle  $ABC$ .



**\*224.** To find the sides and angles of the pedal triangle.

In the figure of the last article, the points  $K, O, G, B$  are concyclic;

$$\therefore \angle OGK = \angle OBK = 90^\circ - A.$$

Also the points  $H, O, G, C$  are concyclic;

$$\therefore \angle OGH = \angle OCH = 90^\circ - A;$$

$$\therefore \angle KGH = 180^\circ - 2A.$$

Thus the angles of the pedal triangle are

$$180^\circ - 2A, \quad 180^\circ - 2B, \quad 180^\circ - 2C.$$

Again,  $\angle AKH = 180^\circ - \angle BKH = \angle BCH$ ,

since the points  $B, K, H, C$  are concyclic;

$$\therefore \angle AKH = C.$$

$$\therefore \frac{HK}{AH} = \frac{\sin A}{\sin AKH} = \frac{\sin A}{\sin C} = \frac{a}{c};$$

$$\therefore HK = \frac{a}{c} \cdot AH = \frac{a}{c} \cdot c \cos A = a \cos A.$$

Thus the sides of the pedal triangle are

$$a \cos A, \quad b \cos B, \quad c \cos C.$$

In terms of  $R$ , the equivalent forms become

$$R \sin 2A, \quad R \sin 2B, \quad R \sin 2C.$$

If the angle  $ACB$  of the given triangle is obtuse, the expressions  $180^\circ - 2C$  and  $c \cos C$  are both negative, and the values we have obtained require some modification. We leave the student to shew that in this case the angles are  $2A, 2B, 2C - 180^\circ$ , and the sides  $a \cos A, b \cos B, -c \cos C$ .

**\*225.** To find the area and circum-radius of the pedal triangle.

$$\text{The area} = \frac{1}{2} (\text{product of two sides}) \times (\text{sine of included angle})$$

$$= \frac{1}{2} R \sin 2B \cdot R \sin 2C \cdot \sin (180^\circ - 2A)$$

$$= \frac{1}{2} R^2 \sin 2A \sin 2B \sin 2C.$$


$$\text{The circum-radius} = \frac{HK}{2 \sin HGK} = \frac{R \sin 2A}{2 \sin (180^\circ - 2A)} = \frac{R}{2}.$$

NOTE. The circum-circle of the pedal triangle is the nine points circle of the triangle  $ABC$ . Thus the radius of the nine points circle of the triangle  $ABC$  is  $\frac{R}{2}$ . [See Hall and Stevens' *Euclid*, p. 281.]

**\*226.** In Art. 224, we have proved that  $OG, OH, OK$  bisect the angles  $HGK, KHG, GKH$  respectively, so that  $O$  is the in-centre of the triangle  $GHK$ . Thus the orthocentre of a triangle is the in-centre of the pedal triangle.

Again, the line  $CGB$  which is at right angles to  $OG$  bisects  $\angle HGK$  externally. Similarly the lines  $AHC$  and  $BKA$  bisect  $\angle KHG$  and  $\angle GKH$  externally, so that  $ABC$  is the ex-central triangle of its pedal triangle  $GHK$ .

\*227. In Art. 219, we have seen that  $ABC$  is the pedal triangle of its ex-central triangle  $I_1I_2I_3$ . Certain theorems depending on this connection are more evident from the adjoining figure, in which the fact that  $ABC$  is the pedal triangle of  $I_1I_2I_3$  is brought more prominently into view. For instance, the circum-circle of the triangle  $ABC$  is the nine points circle of the triangle  $I_1I_2I_3$ , and passes through the middle points of  $II_1$ ,  $II_2$ ,  $II_3$  and of  $I_2I_3$ ,  $I_3I_1$ ,  $I_1I_2$ .



\*228. To find the distance between the in-centre and circum-centre.

Let  $S$  be the circum-centre and  $I$  the in-centre. Produce  $AI$  to meet the circum-circle in  $H$ ; join  $CH$  and  $CI$ .

Draw  $IE$  perpendicular to  $AC$ . Produce  $HS$  to meet the circumference in  $L$ , and join  $CL$ . Then

$$\begin{aligned}\angle HIC &= \angle IAC + \angle ICA \\ &= \frac{A}{2} + \frac{C}{2};\end{aligned}$$


$$\begin{aligned}\angle HCI &= \angle ICB + \angle BCH \\ &= \frac{C}{2} + \angle BAH \\ &= \frac{C}{2} + \frac{A}{2};\end{aligned}$$

$$\therefore \angle HCI = \angle HIC;$$

$$\therefore HI = HC = 2R \sin \frac{A}{2}.$$

Also  $AI = IE \operatorname{cosec} \frac{A}{2} = r \operatorname{cosec} \frac{A}{2};$

$$\therefore AI \cdot IH = 2Rr.$$







**\*230.** *To find the distance of the orthocentre from the circumcentre.*

With the usual notation, we have


$$SO^2 = SA^2 + AO^2 - 2SA \cdot AO \cos SAO.$$

Now  $AS = R$ ;

$$\begin{aligned} AO &= AH \operatorname{cosec} C \\ &= c \cos A \operatorname{cosec} C \\ &= 2R \sin C \cos A \operatorname{cosec} C \\ &= 2R \cos A; \end{aligned}$$

$$\begin{aligned} \angle SAO &= \angle SAC - \angle OAC \\ &= (90^\circ - B) - (90^\circ - C) \\ &= C - B. \end{aligned}$$

$$\begin{aligned} \therefore SO^2 &= R^2 + 4R^2 \cos^2 A - 4R^2 \cos A \cos (C - B) \\ &= R^2 - 4R^2 \cos A \{ \cos (B + C) + \cos (C - B) \} \\ &= R^2 - 8R^2 \cos A \cos B \cos C. \end{aligned}$$



The student may apply a similar method to establish the results of the last two articles.

### \*EXAMPLES. XVIII. c.

1. Shew that the distance of the in-centre from  $A$  is

$$4R \sin \frac{B}{2} \sin \frac{C}{2}.$$

2. Shew that the distances of the ex-centre  $I_1$  from the angular points  $A, B, C$  are

$$4R \cos \frac{B}{2} \cos \frac{C}{2}, \quad 4R \sin \frac{A}{2} \cos \frac{C}{2}, \quad 4R \sin \frac{A}{2} \cos \frac{B}{2}.$$

3. Prove that the area of the ex-central triangle is equal to

$$(1) \ 2Rs; \quad (2) \ \frac{1}{2} \Delta \operatorname{cosec} \frac{A}{2} \operatorname{cosec} \frac{B}{2} \operatorname{cosec} \frac{C}{2}.$$

4. Shew that

$$r \cdot II_1 \cdot II_2 \cdot II_3 = 4R \cdot IA \cdot IB \cdot IC.$$

5. Shew that the perimeter and in-radius of the pedal triangle are respectively

$$4R \sin A \sin B \sin C \quad \text{and} \quad 2R \cos A \cos B \cos C.$$

6. If  $g, h, k$  denote the sides of the pedal triangle, prove that

$$(1) \quad \frac{g}{a^2} + \frac{h}{b^2} + \frac{k}{c^2} = \frac{a^2 + b^2 + c^2}{2abc};$$

$$(2) \quad \frac{(b^2 - c^2)g}{a^2} + \frac{(c^2 - a^2)h}{b^2} + \frac{(a^2 - b^2)k}{c^2} = 0.$$

7. Prove that the ex-radii of the pedal triangle are  $2R \sin A \cos B \cos C$ ,  $2R \cos A \sin B \cos C$ ,  $2R \cos A \cos B \sin C$ .

8. Prove that any formula which connects the sides and angles of a triangle holds if we replace

$$(1) \quad a, b, c \text{ by } a \cos A, \quad b \cos B, \quad c \cos C, \\ \text{and } A, B, C \text{ by } 180^\circ - 2A, \quad 180^\circ - 2B, \quad 180^\circ - 2C;$$

$$(2) \quad a, b, c \text{ by } a \operatorname{cosec} \frac{A}{2}, \quad b \operatorname{cosec} \frac{B}{2}, \quad c \operatorname{cosec} \frac{C}{2},$$

$$\text{and } A, B, C \text{ by } 90^\circ - \frac{A}{2}, \quad 90^\circ - \frac{B}{2}, \quad 90^\circ - \frac{C}{2}.$$

9. Prove that the radius of the circum-circle is never less than the diameter of the in-circle.

10. If  $R = 2r$ , shew that the triangle is equilateral.

11. Prove that

$$SI^2 + SI_1^2 + SI_2^2 + SI_3^2 = 12R^2.$$

12. Prove that

$$(1) \quad a \cdot AI^2 + b \cdot BI^2 + c \cdot CI^2 = abc;$$

$$(2) \quad a \cdot AI_1^2 - b \cdot BI_1^2 - c \cdot CI_1^2 = abc.$$

13. If  $GHK$  be the pedal triangle, and  $O$  the orthocentre, prove that

$$(1) \quad \frac{OG}{AG} + \frac{OH}{BH} + \frac{OK}{CK} = 1;$$

$$(2) \quad \frac{OG}{OG + a \cot A} + \frac{OH}{OH + b \cot B} + \frac{OK}{OK + c \cot C} = 1.$$

14. If  $GHK$  be the pedal triangle, shew that the sum of the circum-radii of the triangles  $AHK$ ,  $BKG$ ,  $CGH$  is equal to  $R + r$ .

15. If  $A_1B_1C_1$  is the ex-central triangle of  $ABC$ , and  $A_2B_2C_2$  the ex-central triangle of  $A_1B_1C_1$ , and  $A_3B_3C_3$  the ex-central triangle of  $A_2B_2C_2$ , and so on: find the angles of the triangle  $A_nB_nC_n$ , and prove that when  $n$  is indefinitely increased the triangle becomes equilateral.

16. Prove that

$$(1) \quad OS^2 = 9R^2 - a^2 - b^2 - c^2;$$

$$(2) \quad OI^2 = 2r^2 - 4R^2 \cos A \cos B \cos C;$$

$$(3) \quad OI_1^2 = 2r_1^2 - 4R^2 \cos A \cos B \cos C.$$

17. If  $f, g, h$  denote the distances of the circum-centre of the pedal triangle from the angular points of the original triangle, shew that

$$4(f^2 + g^2 + h^2) = 11R^2 + 8R^2 \cos A \cos B \cos C.$$

### Quadrilaterals.

\*231. To prove that the area of a quadrilateral is equal to

$$\frac{1}{2} (\text{product of the diagonals}) \times (\text{sine of included angle}).$$


Let the diagonals  $AC, BD$  intersect at  $P$ , and let  $\angle DPA = a$ , and let  $S$  denote the area of the quadrilateral.

$$\Delta DAC = \Delta APD + \Delta CPD$$

$$\begin{aligned} &= \frac{1}{2} DP \cdot AP \sin a \\ &\quad + \frac{1}{2} DP \cdot PC \sin (\pi - a) \\ &= \frac{1}{2} DP (AP + PC) \sin a \\ &= \frac{1}{2} DP \cdot AC \sin a. \end{aligned}$$

$$\text{Similarly} \quad \Delta ABC = \frac{1}{2} BP \cdot AC \sin a.$$

$$\begin{aligned} \therefore S &= \frac{1}{2} (DP + BP) AC \sin a \\ &= \frac{1}{2} DB \cdot AC \sin a. \end{aligned}$$



**\*232.** To find the area of a quadrilateral in terms of the sides and the sum of two opposite angles.

Let  $ABCD$  be the quadrilateral, and let  $a, b, c, d$  be the lengths of its sides,  $S$  its area.

By equating the two values of  $BD^2$  found from the triangles  $BAD, BCD$ , we have

$$\begin{aligned} a^2 + d^2 - 2ad \cos A &= b^2 + c^2 - 2bc \cos C; \\ \therefore a^2 + d^2 - b^2 - c^2 &= 2ad \cos A - 2bc \cos C \dots\dots\dots(1). \end{aligned}$$

Also  $S = \text{sum of areas of triangles } BAD, BCD$

$$\begin{aligned} &= \frac{1}{2} ad \sin A + \frac{1}{2} bc \sin C; \\ \therefore 4S &= 2ad \sin A + 2bc \sin C \dots\dots\dots(2). \end{aligned}$$

Square (2) and add to the square of (1);

$$\therefore 16S^2 + (a^2 + d^2 - b^2 - c^2)^2 = 4a^2d^2 + 4b^2c^2 - 8abcd \cos(A + C).$$

Let  $A + C = 2a$ ; then

$$\begin{aligned} \cos(A + C) &= \cos 2a = 2 \cos^2 a - 1; \\ \therefore 16S^2 &= 4(ad + bc)^2 - (a^2 + d^2 - b^2 - c^2)^2 - 16abcd \cos^2 a. \end{aligned}$$

But the first two terms on the right

$$\begin{aligned} &= (2ad + 2bc + a^2 + d^2 - b^2 - c^2)(2ad + 2bc - a^2 - d^2 + b^2 + c^2) \\ &= \{(a + d)^2 - (b - c)^2\} \{(b + c)^2 - (a - d)^2\} \\ &= (a + d + b - c)(a + d - b + c)(b + c + a - d)(b + c - a + d) \\ &= (2\sigma - 2c)(2\sigma - 2b)(2\sigma - 2d)(2\sigma - 2a), \\ &\qquad\qquad\qquad \text{where } a + b + c + d = 2\sigma, \\ &= 16(\sigma - a)(\sigma - b)(\sigma - c)(\sigma - d). \end{aligned}$$

Thus  $S^2 = (\sigma - a)(\sigma - b)(\sigma - c)(\sigma - d) - abcd \cos^2 a$ ,  
where  $2\sigma$  denotes the sum of the sides,  $2a$  the sum of either pair of opposite angles.

**\*233.** In the case of a *cyclic quadrilateral*,  $A + C = 180^\circ$ , so that  $a = 90^\circ$ ; hence

$$S = \sqrt{(\sigma - a)(\sigma - b)(\sigma - c)(\sigma - d)}.$$

This formula may be obtained directly as in the last article

by making use of the condition  $A + C = 180^\circ$  during the course of the work. In this case  $\cos C = -\cos A$ , and  $\sin C = \sin A$ , so that the expressions (1) and (2) become

$$a^2 + d^2 - b^2 - c^2 = 2(ad + bc) \cos A,$$

and

$$4S = 2(ad + bc) \sin A;$$

whence by eliminating  $A$  we obtain

$$16S^2 + (a^2 + d^2 - b^2 - c^2)^2 = 4(ad + bc)^2.$$

**\*234.** *To find the diagonals and the circum-radius of a cyclic quadrilateral.*

If  $ABCD$  is a cyclic quadrilateral, we have just proved that

$$2(ad + bc) \cos A = a^2 + d^2 - b^2 - c^2.$$

Now  $BD^2 = a^2 + d^2 - 2ad \cos A$

$$\begin{aligned} &= a^2 + d^2 - \frac{ad(a^2 + d^2 - b^2 - c^2)}{ad + bc} \\ &= \frac{bc(a^2 + d^2) + ad(b^2 + c^2)}{ad + bc} \\ &= \frac{(ab + cd)(ac + bd)}{ad + bc}. \end{aligned}$$

Similarly, we may prove that

$$AC^2 = \frac{(ad + bc)(ac + bd)}{ab + cd}.$$

Thus

$$AC \cdot BD = ac + bd, \quad [\text{Compare Euc. VI. D.}]$$

and

$$\frac{AC}{BD} = \frac{ad + bc}{ab + cd}.$$

The circle passing round the quadrilateral circumscribes the triangle  $ABD$ ; hence

$$\text{the circum-radius} = \frac{BD}{2 \sin A}$$

$$\begin{aligned} &= \frac{(ad + bc) BD}{2(ad + bc) \sin A} = \frac{(ad + bc) BD}{4S} \\ &= \frac{1}{4S} \sqrt{(ab + cd)(ac + bd)(ad + bc)}. \end{aligned}$$



*Example.* A quadrilateral  $ABCD$  is such that one circle can be inscribed in it and another circle circumscribed about it; shew that

$$\tan^2 \frac{A}{2} = \frac{bc}{ad}.$$

If a circle can be inscribed in a quadrilateral, the sum of one pair of the opposite sides is equal to that of the other pair;

$$\therefore a+c=b+d.$$

Since the quadrilateral is cyclic,

$$\cos A = \frac{a^2 + d^2 - b^2 - c^2}{2(ad+bc)}. \quad [\text{Art. 233.}]$$

But  $a-d=b-c$ , so that  $a^2-2ad+d^2=b^2-2bc+c^2$ ;

$$\therefore a^2 + d^2 - b^2 - c^2 = 2(ad-bc);$$

$$\therefore \cos A = \frac{ad-bc}{ad+bc};$$

$$\therefore \tan^2 \frac{A}{2} = \frac{1-\cos A}{1+\cos A} = \frac{bc}{ad}.$$

### \*EXAMPLES. XVIII. d.

1. If a circle can be inscribed in a quadrilateral, shew that its radius is  $S/\sigma$  where  $S$  is the area and  $2\sigma$  the sum of the sides of the quadrilateral.

2. If the sides of a cyclic quadrilateral be 3, 3, 4, 4, shew that a circle can be inscribed in it, and find the radii of the inscribed and circumscribed circles.

3. If the sides of a cyclic quadrilateral be 1, 2, 4, 3, shew that the cosine of the angle between the two greatest sides is  $\frac{5}{7}$ , and that the radius of the inscribed circle is  $\cdot 98$  nearly.

4. The sides of a cyclic quadrilateral are 60, 25, 52, 39: shew that two of the angles are right angles, and find the diagonals and the area.

5. The sides of a quadrilateral are 4, 5, 8, 9, and one diagonal is 9: find the area.

6. If a circle can be inscribed in a cyclic quadrilateral, shew that the area of the quadrilateral is  $\sqrt{abcd}$ , and that the radius of the circle is

$$2\sqrt{abcd}/(a+b+c+d).$$

7. If the sides of a quadrilateral are given, shew that the area is a maximum when the quadrilateral can be inscribed in a circle.

8. If the sides of a quadrilateral are 23, 29, 37, 41 inches, prove that the maximum area is 7 sq. ft.

9. If  $ABCD$  is a cyclic quadrilateral, prove that

$$\tan^2 \frac{B}{2} = \frac{(\sigma - a)(\sigma - b)}{(\sigma - c)(\sigma - d)}.$$

10. If  $f, g$  denote the diagonals of a quadrilateral and  $\beta$  the angle between them, prove that

$$2fg \cos \beta = (a^2 + c^2) - (b^2 + d^2).$$

11. If  $\beta$  is the angle between the diagonals of any quadrilateral, prove that the area is

$$\frac{1}{4} \{ (a^2 + c^2) - (b^2 + d^2) \} \tan \beta.$$

12. Prove that the area of a quadrilateral in which a circle can be inscribed is

$$\sqrt{abcd} \sin \frac{A + C}{2}.$$

13. If a circle can be inscribed in a quadrilateral whose diagonals are  $f$  and  $g$ , prove that

$$4S^2 = f^2 g^2 - (ac - bd)^2.$$

14. If  $\beta$  is the angle between the diagonals of a cyclic quadrilateral, prove that

$$(1) \quad (ac + bd) \sin \beta = (ad + bc) \sin A;$$

$$(2) \quad \cos \beta = \frac{(a^2 + c^2) - (b^2 + d^2)}{2(ac + bd)};$$

$$(3) \quad \tan^2 \frac{\beta}{2} = \frac{(\sigma - b)(\sigma - d)}{(\sigma - a)(\sigma - c)} \text{ or } \frac{(\sigma - a)(\sigma - c)}{(\sigma - b)(\sigma - d)}.$$

15. If  $f, g$  are the diagonals of a quadrilateral, shew that

$$S = \frac{1}{4} \sqrt{4f^2 g^2 - (a^2 + c^2 - b^2 - d^2)^2}.$$

16. In a cyclic quadrilateral, prove that the product of the segments of a diagonal is

$$abcd(ac + bd)/(ab + cd)(ad + bc).$$

235. The following exercise consists of miscellaneous questions involving the properties of triangles.

### EXAMPLES. XVIII. e.

1. If the sides of a triangle are 242, 1212, 1450 yards, shew that the area is 6 acres.

2. One of the sides of a triangle is 200 yards and the adjacent angles are  $22.5^\circ$  and  $67.5^\circ$ ; find the area.

3. If  $r_1 = 2r_2 = 2r_3$ , shew that  $3a = 4b$ .

4. If  $a, b, c$  are in A. P., shew that  $r_1, r_2, r_3$  are in H. P.

5. Find the area of a triangle whose sides are

$$\frac{y}{z} + \frac{z}{x}, \quad \frac{z}{x} + \frac{x}{y}, \quad \frac{x}{y} + \frac{y}{z}.$$

6. If  $\sin A : \sin C = \sin(A - B) : \sin(B - C)$ , shew that  $a^2, b^2, c^2$  are in A. P.

Prove that

$$7. \quad \frac{a \sin A + b \sin B + c \sin C}{4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}} = \frac{a^2 + b^2 + c^2}{2s}.$$

$$8. \quad \left( \frac{a^2}{\sin A} + \frac{b^2}{\sin B} + \frac{c^2}{\sin C} \right) \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} = \Delta.$$

$$9. \quad (r_2 + r_3)(r_3 + r_1)(r_1 + r_2) = 4R(r_2 r_3 + r_3 r_1 + r_1 r_2).$$

$$10. \quad \tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2} = \frac{r_1 + r_2 + r_3}{(r_2 r_3 + r_3 r_1 + r_1 r_2)^{\frac{1}{2}}}.$$

$$11. \quad bc \cot \frac{A}{2} + ca \cot \frac{B}{2} + ab \cot \frac{C}{2} = 4Rs^2 \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} - \frac{3}{s} \right).$$

$$12. \quad \left( \frac{1}{r} + \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} \right)^2 = \frac{4}{r} \left( \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} \right).$$

13. The perimeter of a right-angled triangle is 70, and the in-radius is 6: find the sides.

14. If  $f, g, h$  are the perpendiculars from the circum-centre on the sides, prove that

$$\frac{a}{f} + \frac{b}{g} + \frac{c}{h} = \frac{abc}{4fgh}.$$

15. An equilateral triangle and a regular hexagon have the same perimeter: shew that the areas of their inscribed circles are as 4 to 9.

\*16. Shew that the perimeter of the pedal triangle is equal to  $abc/2R^2$ .

\*17. Shew that the area of the ex-central triangle is equal to  $abc(a+b+c)/4\Delta$ .

18. In the ambiguous case, if  $A$ ,  $a$ ,  $b$  are the given parts, and  $c_1, c_2$  the two values of the third side, shew that the distance between the circum-centres of the two triangles is  $\frac{c_1 \sim c_2}{2 \sin A}$ .

\*19. If  $\beta$  be the angle between the diagonals of a cyclic quadrilateral, shew that

$$\sin \beta = \frac{2S}{ac + bd}.$$

\*20. Shew that

$$r^3 \cdot II_1 \cdot II_2 \cdot II_3 = IA^2 \cdot IB^2 \cdot IC^2.$$

\*21. Shew that the sum of the squares of the sides of the ex-central triangle is equal to  $8R(4R+r)$ .

\*22. If circles can be inscribed in and circumscribed about a quadrilateral, and if  $\beta$  be the angle between the diagonals, shew that

$$\cos \beta = (ac \sim bd)/(ac + bd).$$

23. If  $l, m, n$  are the lengths of the medians of a triangle, prove that

- (1)  $4(l^2 + m^2 + n^2) = 3(a^2 + b^2 + c^2)$ ;
- (2)  $(b^2 - c^2)l^2 + (c^2 - a^2)m^2 + (a^2 - b^2)n^2 = 0$ ;
- (3)  $16(l^4 + m^4 + n^4) = 9(a^4 + b^4 + c^4)$ .

24. Shew that the radii of the escribed circles are the roots of the equation

$$x^3 - (4R + r)x^2 + s^2x - s^2r = 0.$$

25. If  $\Delta_1, \Delta_2, \Delta_3$  be the areas of the triangles cut off by tangents to the in-circle parallel to the sides of a triangle, prove that

$$\frac{\Delta_1}{(s-a)^2} = \frac{\Delta_2}{(s-b)^2} = \frac{\Delta_3}{(s-c)^2} = \frac{\Delta}{s^2}.$$

\*26. The triangle  $LMN$  is formed by joining the points of contact of the in-circle; shew that it is similar to the ex-central triangle, and that their areas are as  $r^2$  to  $4R^2$ .

27. In the triangle  $PQR$  formed by drawing tangents at  $A, B, C$  to the circum-circle, prove that the angles and sides are

$$180^\circ - 2A, \quad 180^\circ - 2B, \quad 180^\circ - 2C;$$

and 
$$\frac{a}{2 \cos B \cos C}, \quad \frac{b}{2 \cos C \cos A}, \quad \frac{c}{2 \cos A \cos B}.$$

28. If  $p, q, r$  be the lengths of the bisectors of the angles of a triangle, prove that

$$(1) \quad \frac{1}{p} \cos \frac{A}{2} + \frac{1}{q} \cos \frac{B}{2} + \frac{1}{r} \cos \frac{C}{2} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c};$$

$$(2) \quad \frac{pqr}{4\Delta} = \frac{abc(\alpha + b + c)}{(b+c)(c+a)(a+b)}.$$

29. If the perpendiculars  $AG, BH, CK$  are produced to meet the circum-circle in  $L, M, N$ , prove that

$$(1) \quad \text{area of triangle } LMN = 8\Delta \cos A \cos B \cos C;$$

$$(2) \quad AL \sin A + BM \sin B + CN \sin C = 8R \sin A \sin B \sin C.$$

30. If  $r_a, r_b, r_c$  be the radii of the circles inscribed between the in-circle and the sides containing the angles  $A, B, C$  respectively, shew that

$$(1) \quad r_a = r \tan^2 \frac{\pi - A}{4}; \quad (2) \quad \sqrt{r_b r_c} + \sqrt{r_c r_a} + \sqrt{r_a r_b} = r.$$

\*31. Lines drawn through the angular points of a triangle  $ABC$  parallel to the sides of the pedal triangle form a triangle  $XYZ$ : shew that the perimeter and area of  $XYZ$  are respectively

$$2R \tan A \tan B \tan C \quad \text{and} \quad R^2 \tan A \tan B \tan C.$$

\*32. A straight line cuts three concentric circles in  $A, B, C$  and passes at a distance  $p$  from their centre: shew that the area of the triangle formed by the tangents at  $A, B, C$  is

$$\frac{BC \cdot CA \cdot AB}{2p}.$$

### MISCELLANEOUS EXAMPLES. F.

1. If  $a + \beta + \gamma + \delta = 180^\circ$ , shew that

$$\cos a \cos \beta + \cos \gamma \cos \delta = \sin a \sin \beta + \sin \gamma \sin \delta.$$

2. Prove that

$$\cos(15^\circ - A) \sec 15^\circ - \sin(15^\circ - A) \operatorname{cosec} 15^\circ = 4 \sin A.$$

3. Shew that in a triangle

$$\cot A + \sin A \operatorname{cosec} B \operatorname{cosec} C$$

retains the same value if any two of the angles  $A, B, C$  are interchanged.

4. If  $a = 2$ ,  $b = \sqrt{8}$ ,  $A = 30^\circ$ , solve the triangle.

5. Shew that

$$(1) \quad \cot 18^\circ = \sqrt{5} \cot 36^\circ;$$

$$(2) \quad 16 \sin 36^\circ \sin 72^\circ \sin 108^\circ \sin 144^\circ = 5.$$

6. Find the number of ciphers before the first significant digit in  $(.0396)^{90}$ , given

$$\log 2 = .30103, \log 3 = .47712, \log 11 = 1.04139.$$

7. An observer finds that the angle subtended by the line joining two points  $A$  and  $B$  on the horizontal plane is  $30^\circ$ . On walking 50 yards directly towards  $A$  the angle increases to  $75^\circ$ : find his distance from  $B$  at each observation.

8. Prove that  $\cos^2 a + \cos^2 \beta + \cos^2 \gamma + \cos^2(a + \beta + \gamma)$   
 $= 2 + 2 \cos(\beta + \gamma) \cos(\gamma + a) \cos(a + \beta).$

9. Shew that

$$(1) \quad \tan 40^\circ + \cot 40^\circ = 2 \sec 10^\circ;$$

$$(2) \quad \tan 70^\circ + \tan 20^\circ = 2 \operatorname{cosec} 40^\circ.$$

10. Prove that

$$(1) \quad 2 \sin 4a - \sin 10a + \sin 2a = 16 \sin a \cos a \cos 2a \sin^2 3a;$$

$$(2) \quad \sin \frac{2\pi}{7} + \sin \frac{4\pi}{7} - \sin \frac{6\pi}{7} = 4 \sin \frac{\pi}{7} \sin \frac{3\pi}{7} \sin \frac{5\pi}{7}.$$



11. If  $B=30^\circ$ ,  $b=3\sqrt{2}-\sqrt{6}$ ,  $c=6-2\sqrt{3}$ , solve the triangle.

12. From a ship which is sailing N.E., the bearing of a rock is N.N.W. After the ship has sailed 10 miles the rock bears due W.: find the distance of the ship from the rock at each observation.

13. Shew that in any triangle

$$\frac{b^2 - c^2}{\cos B + \cos C} + \frac{c^2 - a^2}{\cos C + \cos A} + \frac{a^2 - b^2}{\cos A + \cos B} = 0.$$

14. If  $\cos(\theta - a)$ ,  $\cos \theta$ ,  $\cos(\theta + a)$  are in harmonical progression, shew that

$$\cos \theta = \sqrt{2} \cos \frac{a}{2}.$$

15. If  $\sin \beta$  be the geometric mean between  $\sin a$  and  $\cos a$ , prove that

$$\cos 2\beta = 2 \cos^2 \left( \frac{\pi}{4} + a \right).$$

16. Shew that the distances of the orthocentre from the sides are  $2R \cos B \cos C$ ,  $2R \cos C \cos A$ ,  $2R \cos A \cos B$ .

17. If  $\cos \theta = \frac{\cos u - e}{1 - e \cos u}$ ,

prove that  $\tan \frac{\theta}{2} = \sqrt{\frac{1+e}{1-e}} \tan \frac{u}{2}.$

18. If the sides of a right-angled triangle are

$$2(1 + \sin \theta) + \cos \theta \quad \text{and} \quad 2(1 + \cos \theta) + \sin \theta,$$

prove that the hypotenuse is

$$3 + 2(\cos \theta + \sin \theta).$$

\*19. Prove that the distances of the in-centre of the ex-central triangle  $I_1 I_2 I_3$  from its ex-centres are

$$8R \sin \frac{B+C}{4}, \quad 8R \sin \frac{C+A}{4}, \quad 8R \sin \frac{A+B}{4}.$$

\*20. Prove that the distances between the ex-centres of the ex-central triangle  $I_1 I_2 I_3$  are

$$8R \cos \frac{B+C}{4}, \quad 8R \cos \frac{C+A}{4}, \quad 8R \cos \frac{A+B}{4}.$$

21. If

$(1 + \cos \alpha)(1 + \cos \beta)(1 + \cos \gamma) = (1 - \cos \alpha)(1 - \cos \beta)(1 - \cos \gamma)$ ,  
shew that each expression is equal to  $\pm \sin \alpha \sin \beta \sin \gamma$ .

22. If the sum of four angles is  $180^\circ$ , shew that the sum of the products of their sines taken two together is equal to the sum of the products of their cosines taken two together.

\*23. In a triangle, shew that

$$(1) II_1 \cdot II_2 \cdot II_3 = 16R^2r; \quad (2) II_1^2 + II_2^2 + II_3^2 = 16R^2.$$

24. Find the angles of a triangle whose sides are proportional to

$$(1) \cos \frac{A}{2}, \quad \cos \frac{B}{2}, \quad \cos \frac{C}{2};$$

$$(2) \sin 2A, \quad \sin 2B, \quad \sin 2C.$$

25. Prove that the expression

$$\sin^2(\theta + \alpha) + \sin^2(\theta + \beta) - 2 \cos(\alpha - \beta) \sin(\theta + \alpha) \sin(\theta + \beta)$$

is independent of  $\theta$ .

\*26. If  $a, b, c, d$  are the sides of a quadrilateral described about a circle, prove that

$$ab \sin^2 \frac{A}{2} = cd \sin^2 \frac{C}{2}.$$

27. Tangents parallel to the three sides are drawn to the in-circle. If  $p, q, r$  be the lengths of the parts of the tangents within the triangle, prove that  $\frac{p}{a} + \frac{q}{b} + \frac{r}{c} = 1$ .

[The Tables will be required for Examples 28 and 29.]

28. From the top of a cliff 1566 ft. in height a train, which is travelling at a uniform speed in a straight line to a tunnel immediately below the observer, is seen to pass two consecutive stations at an interval of 3 minutes. The angles of depression of the two stations are  $13^\circ 14' 12''$  and  $56^\circ 24' 36''$  respectively; how fast is the train travelling?

29. A harbour lies in a direction  $46^\circ 8' 8''$  South of West from a fort, and at a distance of 27.23 miles from it. A ship sets out from the harbour at noon and sails due East at 10 miles an hour; when will the ship be 20 miles from the fort?

## CHAPTER XIX.

### GENERAL VALUES AND INVERSE FUNCTIONS.

236. THE equation  $\sin \theta = \frac{1}{2}$  is satisfied by  $\theta = \frac{\pi}{6}$ , and by  $\theta = \pi - \frac{\pi}{6}$ , and all angles coterminal with these will have the same sine. This example shews that there are an infinite number of angles whose sine is equal to a given quantity. Similar remarks apply to the other functions.


We proceed to shew how to express by a single formula all angles which have a given sine, cosine, or tangent.

237. From the results proved in Chap. IX., it is easily seen that in going once through the four quadrants, there are two and only two positions of the boundary line which give angles with the same sine, cosine, or tangent.


Thus if  $\sin a$  has a given value, the positions of the radius vector are  $OP$  and  $OP'$  bounding the angles  $a$  and  $\pi - a$ . [Art. 92.]



If  $\cos a$  has a given value, the positions of the radius vector are  $OP$  and  $OP'$  bounding the angles  $a$  and  $2\pi - a$ . [Art. 105.]




If  $\tan a$  has a given value, the positions of the radius vector are  $OP$  and  $OP'$  bounding the angles  $a$  and  $\pi + a$ . [Art. 97.]



**238.** To find a formula for all the angles which have a given sine.

Let  $\alpha$  be the smallest positive angle which has a given sine. Draw  $OP$  and  $OP'$  bounding the angles  $\alpha$  and  $\pi - \alpha$ ; then the required angles are those coterminal with  $OP$  and  $OP'$ .



The positive angles are

$$2p\pi + \alpha \text{ and } 2p\pi + (\pi - \alpha),$$

where  $p$  is zero, or any positive integer.

The negative angles are

$$-(\pi + \alpha) \text{ and } -(2\pi - \alpha),$$

and those which may be obtained from them by the addition of any negative multiple of  $2\pi$ ; that is, angles denoted by

$$2q\pi - (\pi + \alpha) \text{ and } 2q\pi - (2\pi - \alpha),$$

where  $q$  is zero, or any negative integer.

These angles may be grouped as follows :

$$\left. \begin{array}{l} 2p\pi + \alpha, \\ (2q - 2)\pi + \alpha, \end{array} \right\} \text{ and } \left\{ \begin{array}{l} (2p + 1)\pi - \alpha, \\ (2q - 1)\pi - \alpha, \end{array} \right.$$

and it will be noticed that even multiples of  $\pi$  are followed by  $+\alpha$ , and odd multiples of  $\pi$  by  $-\alpha$ .

Thus all angles equi-sinal with  $\alpha$  are included in the formula

$$n\pi + (-1)^n \alpha,$$

where  $n$  is zero, or any integer positive or negative.

This is also the formula for all angles which have the same cosecant as  $\alpha$ .

*Example 1.* Write down the general solution of  $\sin \theta = \frac{\sqrt{3}}{2}$ .

The least value of  $\theta$  which satisfies the equation is  $\frac{\pi}{3}$ ; therefore the general solution is  $n\pi + (-1)^n \frac{\pi}{3}$ .

*Example 2.* Find the general solution of  $\sin^2 \theta = \sin^2 \alpha$ .


This equation gives either  $\sin \theta = +\sin \alpha \dots\dots\dots(1)$ ,  
or  $\sin \theta = -\sin \alpha = \sin(-\alpha) \dots\dots\dots(2)$ .

From (1),  $\theta = n\pi + (-1)^n a$ ;  
 and from (2),  $\theta = n\pi + (-1)^n (-a)$ .

Both values are included in the formula  $\theta = n\pi \pm a$ .

**239.** To find a formula for all the angles which have a given cosine.

Let  $a$  be the smallest positive angle which has a given cosine. Draw  $OP$  and  $OP'$  bounding the angles  $a$  and  $2\pi - a$ ; then the required angles are those coterminal with  $OP$  and  $OP'$ .



The positive angles are

$$2p\pi + a \text{ and } 2p\pi + (2\pi - a),$$

where  $p$  is zero, or any positive integer.

The negative angles are

$$-a \text{ and } -(2\pi - a),$$

and those which may be obtained from them by the addition of any negative multiple of  $2\pi$ ; that is, angles denoted by

$$2q\pi - a \text{ and } 2q\pi - (2\pi - a),$$

where  $q$  is zero, or any negative integer.

The angles may be grouped as follows :

$$\left. \begin{array}{l} 2p\pi + a, \\ 2q\pi - a, \end{array} \right\} \text{ and } \left\{ \begin{array}{l} (2p+2)\pi - a, \\ (2q-2)\pi + a, \end{array} \right.$$

and it will be noticed that the multiples of  $\pi$  are always even, but may be followed by  $+a$  or by  $-a$ .

Thus all angles equi-cosinal with  $a$  are included in the formula

$$2n\pi \pm a,$$

where  $n$  is zero, or any integer positive or negative.


This is also the formula for all angles which have the same secant as  $a$ .

*Example 1.* Find the general solution of  $\cos \theta = -\frac{1}{2}$ .

The least value of  $\theta$  is  $\pi - \frac{\pi}{3}$ , or  $\frac{2\pi}{3}$ ; hence the general solution is  $2n\pi \pm \frac{2\pi}{3}$ .

**240.** To find a formula for all the angles which have a given tangent.

Let  $a$  be the smallest positive angle which has a given tangent. Draw  $OP$  and  $OP'$  bounding the angles  $a$  and  $\pi + a$ ; then the required angles are those coterminal with  $OP$  and  $OP'$ .



The positive angles are

$$2p\pi + a \text{ and } 2p\pi + (\pi + a),$$

where  $p$  is zero, or any positive integer.

The negative angles are

$$-(\pi - a) \text{ and } -(2\pi - a),$$

and those which may be obtained from them by the addition of any negative multiple of  $2\pi$ ; that is, angles denoted by

$$2q\pi - (\pi - a) \text{ and } 2q\pi - (2\pi - a),$$

where  $q$  is zero, or any negative integer.

The angles may be grouped as follows :

$$\left. \begin{array}{l} 2p\pi + a, \\ (2q - 2)\pi + a, \end{array} \right\} \text{ and } \left\{ \begin{array}{l} (2p + 1)\pi + a, \\ (2q - 1)\pi + a, \end{array} \right.$$

and it will be noticed that whether the multiple of  $\pi$  is even or odd, it is always followed by  $+a$ . Thus all angles equi-tangential with  $a$  are included in the formula

$$\theta = n\pi + a.$$

This is also the formula for all the angles which have the same cotangent as  $a$ .

*Example.* Solve the equation  $\cot 4\theta = \cot \theta$ .

The general solution is  $4\theta = n\pi + \theta$ ;

whence  $3\theta = n\pi$ , or  $\theta = \frac{n\pi}{3}$ .

**241.** All angles which are both equi-sinal and equi-cosinal with  $a$  are included in the formula  $2n\pi + a$ .

All angles equi-cosinal with  $a$  are included in the formula  $2n\pi \pm a$ ; so that the multiple of  $\pi$  is even. But in the formula  $n\pi + (-1)^n a$ , which includes all angles equi-sinal with  $a$ , when the multiple of  $\pi$  is even,  $a$  must be preceded by the  $+$  sign. Thus the formula is  $2n\pi + a$ .



**242.** In the solution of equations, the general value of the angle should always be given.

*Example.* Solve the equation  $\cos 9\theta = \cos 5\theta - \cos \theta$ .

By transposition,  $(\cos 9\theta + \cos \theta) - \cos 5\theta = 0$ ;

$$\therefore 2 \cos 5\theta \cos 4\theta - \cos 5\theta = 0;$$

$$\therefore \cos 5\theta (2 \cos 4\theta - 1) = 0;$$

$\therefore$  either  $\cos 5\theta = 0$ , or  $2 \cos 4\theta - 1 = 0$ .

From the first equation,  $5\theta = 2n\pi \pm \frac{\pi}{2}$ , or  $\theta = \frac{(4n \pm 1)\pi}{10}$ ;

and from the second,  $4\theta = 2n\pi \pm \frac{\pi}{3}$ , or  $\theta = \frac{(6n \pm 1)\pi}{12}$ .

### EXAMPLES. XIX. a.

Find the general solution of the equations :

$$1. \sin \theta = \frac{1}{2}. \quad 2. \sin \theta = \frac{1}{\sqrt{2}}. \quad 3. \cos \theta = \frac{1}{2}.$$

$$4. \tan \theta = \sqrt{3}. \quad 5. \cot \theta = -\sqrt{3}. \quad 6. \sec \theta = -\sqrt{2}.$$

$$7. \cos^2 \theta = \frac{1}{2}. \quad 8. \tan^2 \theta = \frac{1}{3}. \quad 9. \operatorname{cosec}^2 \theta = \frac{4}{3}.$$

$$10. \cos \theta = \cos a. \quad 11. \tan^2 \theta = \tan^2 a.$$

$$12. \sec^2 \theta = \sec^2 a. \quad 13. \tan 2\theta = \tan \theta.$$

$$14. \operatorname{cosec} 3\theta = \operatorname{cosec} 3a. \quad 15. \cos 3\theta = \cos 2\theta.$$

$$16. \sin 5\theta + \sin \theta = \sin 3\theta. \quad 17. \cos \theta - \cos 7\theta = \sin 4\theta.$$

$$18. \sin 4\theta - \sin 3\theta + \sin 2\theta - \sin \theta = 0.$$

$$19. \cos \theta + \cos 3\theta + \cos 5\theta + \cos 7\theta = 0.$$

$$20. \sin 5\theta \cos \theta = \sin 6\theta \cos 2\theta.$$

$$21. \sin 11\theta \sin 4\theta + \sin 5\theta \sin 2\theta = 0.$$

$$22. \sqrt{2} \cos 3\theta - \cos \theta = \cos 5\theta.$$

$$23. \sin 7\theta - \sqrt{3} \cos 4\theta = \sin \theta.$$

$$24. 1 + \cos \theta = 2 \sin^2 \theta. \quad 25. \tan^2 \theta + \sec \theta = 1.$$

$$26. \cot^2 \theta - 1 = \operatorname{cosec} \theta. \quad 27. \cot \theta - \tan \theta = 2.$$

$$28. \text{If } 2 \cos \theta = -1 \text{ and } 2 \sin \theta = \sqrt{3}, \text{ find } \theta.$$

$$29. \text{If } \sec \theta = \sqrt{2} \text{ and } \tan \theta = -1, \text{ find } \theta.$$

**243.** In the following examples, the solution is simplified by the use of some particular artifice.

*Example 1.* Solve the equation  $\cos m\theta = \sin n\theta$ .

$$\text{Here} \quad \cos m\theta = \cos \left( \frac{\pi}{2} - n\theta \right);$$

$$\therefore m\theta = 2k\pi \pm \left( \frac{\pi}{2} - n\theta \right),$$

where  $k$  is zero, or any integer.

By transposition, we obtain

$$(m+n)\theta = \left( 2k + \frac{1}{2} \right) \pi, \text{ or } (m-n)\theta = \left( 2k - \frac{1}{2} \right) \pi.$$

This equation may also be solved through the medium of the sine. For we have

$$\sin \left( \frac{\pi}{2} - m\theta \right) = \sin n\theta;$$

$$\therefore \frac{\pi}{2} - m\theta = p\pi + (-1)^p n\theta,$$

where  $p$  is zero or any integer;

$$\therefore \{m + (-1)^p n\} \theta = \left( \frac{1}{2} - p \right) \pi.$$

**NOTE.** The general solution can frequently be obtained in several ways. The various forms which the result takes are merely different modes of expressing the same series of angles.

*Example 2.* Solve  $\sqrt{3} \cos \theta + \sin \theta = 1$ .

Multiply every term by  $\frac{1}{2}$ , then

$$\frac{\sqrt{3}}{2} \cos \theta + \frac{1}{2} \sin \theta = \frac{1}{2},$$

$$\therefore \cos \frac{\pi}{6} \cos \theta + \sin \frac{\pi}{6} \sin \theta = \frac{1}{2};$$

$$\therefore \cos \left( \theta - \frac{\pi}{6} \right) = \frac{1}{2};$$

$$\therefore \theta - \frac{\pi}{6} = 2n\pi \pm \frac{\pi}{3};$$

$$\therefore \theta = 2n\pi + \frac{\pi}{2} \text{ or } 2n\pi - \frac{\pi}{6}.$$

NOTE. In examples of this type, it is a common mistake to square the equation; but this process is objectionable, because it introduces solutions which do not belong to the given equation. Thus in the present instance,

$$\sqrt{3} \cos \theta = 1 - \sin \theta;$$

by squaring,  $3 \cos^2 \theta = (1 - \sin \theta)^2.$

But the solutions of this equation include the solutions of

$$-\sqrt{3} \cos \theta = 1 - \sin \theta,$$

as well as those of the given equation.

*Example 3.* Solve  $\cos 2\theta = \cos \theta + \sin \theta.$

From this equation,  $\cos^2 \theta - \sin^2 \theta = \cos \theta + \sin \theta;$

$$\therefore (\cos \theta + \sin \theta)(\cos \theta - \sin \theta) = \cos \theta + \sin \theta;$$

$$\therefore \text{either} \quad \cos \theta + \sin \theta = 0 \dots \dots \dots (1),$$

$$\text{or} \quad \cos \theta - \sin \theta = 1 \dots \dots \dots (2).$$

From (1),  $\tan \theta = -1,$

$$\therefore \theta = n\pi - \frac{\pi}{4}.$$

From (2),  $\frac{1}{\sqrt{2}} \cos \theta - \frac{1}{\sqrt{2}} \sin \theta = \frac{1}{\sqrt{2}};$

$$\therefore \cos \theta \cos \frac{\pi}{4} - \sin \theta \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}.$$

$$\therefore \cos \left( \theta + \frac{\pi}{4} \right) = \frac{1}{\sqrt{2}};$$

$$\therefore \theta + \frac{\pi}{4} = 2n\pi \pm \frac{\pi}{4};$$

$$\therefore \theta = 2n\pi \text{ or } 2n\pi - \frac{\pi}{2}.$$

### EXAMPLES. XIX. b.

Find the general solution of the equations :

- |  |   |
|--|---|
| 1. $\tan p\theta = \cot q\theta.$              | 2. $\sin m\theta + \cos n\theta = 0.$               |
| 3. $\cos \theta - \sqrt{3} \sin \theta = 1.$   | 4. $\sin \theta - \sqrt{3} \cos \theta = 1.$        |
| 5. $\cos \theta = \sqrt{3} (1 - \sin \theta).$ | 6. $\sin \theta + \sqrt{3} \cos \theta = \sqrt{2}.$ |

Find the general solution of the equations :

7.  $\cos \theta - \sin \theta = \frac{1}{\sqrt{2}}$ .
8.  $\cos \theta + \sin \theta + \sqrt{2} = 0$ .
9.  $\operatorname{cosec} \theta + \cot \theta = \sqrt{3}$ .
10.  $\cot \theta - \cot 2\theta = 2$ .
11.  $2 \sin \theta \sin 3\theta = 1$ .
12.  $\sin 3\theta = 8 \sin^3 \theta$ .
13.  $\tan \theta + \tan 3\theta = 2 \tan 2\theta$ .
14.  $\cos \theta - \sin \theta = \cos 2\theta$ .
15.  $\operatorname{cosec} \theta + \sec \theta = 2\sqrt{2}$ .
16.  $\sec \theta - \operatorname{cosec} \theta = 2\sqrt{2}$ .
17.  $\sec 4\theta - \sec 2\theta = 2$ .
18.  $\cos 3\theta + 8 \cos^3 \theta = 0$ .
19.  $1 + \sqrt{3} \tan^2 \theta = (1 + \sqrt{3}) \tan \theta$ .
20.  $\tan^3 \theta + \cot^3 \theta = 8 \operatorname{cosec}^3 2\theta + 12$ .
21.  $\sin \theta = \sqrt{2} \sin \phi, \quad \sqrt{3} \cos \theta = \sqrt{2} \cos \phi$ .
22.  $\operatorname{cosec} \theta = \sqrt{3} \operatorname{cosec} \phi, \quad \cot \theta = 3 \cot \phi$ .
23.  $\sec \phi = \sqrt{2} \sec \theta, \quad \cot \theta = \sqrt{3} \cot \phi$ .

24. Explain why the same two series of angles are given by the equations

$$\theta + \frac{\pi}{4} = n\pi + (-1)^n \frac{\pi}{6} \quad \text{and} \quad \theta - \frac{\pi}{4} = 2n\pi \pm \frac{\pi}{3}.$$

25. Shew that the formulæ

$$\left(2n + \frac{1}{4}\right)\pi \pm a \quad \text{and} \quad \left(n - \frac{1}{4}\right)\pi + (-1)^n \left(\frac{\pi}{2} - a\right)$$

comprise the same angles, and illustrate by a figure.

### Inverse Circular Functions.

244. If  $\sin \theta = s$ , we know that  $\theta$  may be *any* angle whose sine is  $s$ . It is often convenient to express this statement *inversely* by writing  $\theta = \sin^{-1} s$ .

In this *inverse notation*  $\theta$  stands alone on one side of the equation, and may be regarded as an angle whose value is only known through the medium of its sine. Similarly,  $\tan^{-1} \sqrt{3}$  indicates in a concise form any one of the angles whose tangent is  $\sqrt{3}$ . But all these angles are given by the formula  $n\pi + \frac{\pi}{3}$ . Thus

$$\theta = \tan^{-1} \sqrt{3} \quad \text{and} \quad \theta = n\pi + \frac{\pi}{3}$$

are equivalent statements expressed in different forms.

**245.** Expressions of the form  $\cos^{-1}x$ ,  $\sin^{-1}a$ ,  $\tan^{-1}b$  are called **Inverse Circular Functions**.

It must be remembered that these expressions denote angles, and that  $-1$  is not an algebraical index; that is,

$$\sin^{-1}x \text{ is not the same as } (\sin x)^{-1} \text{ or } \frac{1}{\sin x}.$$

**246.** From Art. 244, we see that an inverse function has an infinite number of values.

If  $f$  denote any one of the circular functions, and  $f^{-1}(x)=A$ , the **principal value** of  $f^{-1}(x)$  is the smallest numerical value of  $A$ . Thus the principal values of

$$\begin{array}{cccc} \cos^{-1}\frac{1}{2}, & \sin^{-1}\left(-\frac{1}{2}\right), & \cos^{-1}\left(-\frac{1}{\sqrt{2}}\right), & \tan^{-1}(-1) \\ \text{are } 60^\circ, & -30^\circ, & 135^\circ, & -45^\circ. \end{array}$$

Hence if  $x$  be positive, the principal values of  $\sin^{-1}x$ ,  $\cos^{-1}x$ ,  $\tan^{-1}x$  all lie between  $0$  and  $90^\circ$ .

If  $x$  be negative, the principal values of  $\sin^{-1}x$  and  $\tan^{-1}x$  lie between  $0$  and  $-90^\circ$ , and the principal value of  $\cos^{-1}x$  lies between  $90^\circ$  and  $180^\circ$ .

In numerical instances we shall usually suppose that the *principal value* is selected.

**247.** If  $\sin \theta = x$ , we have  $\cos \theta = \sqrt{1-x^2}$ .

Expressed in the inverse notation, these equations become

$$\theta = \sin^{-1}x, \quad \theta = \cos^{-1}\sqrt{1-x^2}.$$

In each of these two statements,  $\theta$  has an infinite number of values; but, as the formulæ for the general values of the sine and cosine are not identical, we cannot assert that the equation

$$\sin^{-1}x = \cos^{-1}\sqrt{1-x^2}$$

is identically true. This will be seen more clearly from a numerical instance. If  $x = \frac{1}{2}$ , then  $\sqrt{1-x^2} = \frac{\sqrt{3}}{2}$ .

Here  $\sin^{-1}x$  may be any one of the angles

$$30^\circ, 150^\circ, 390^\circ, 510^\circ, \dots;$$

and  $\cos^{-1}\sqrt{1-x^2}$  may be any one of the angles

$$30^\circ, 330^\circ, 390^\circ, 690^\circ, \dots$$

**248.** From the relations established in the previous chapters, we may deduce corresponding relations connecting the inverse functions. Thus in the identity

$$\cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta},$$

let  $\tan \theta = a$ , so that  $\theta = \tan^{-1} a$ ; then

$$\cos (2 \tan^{-1} a) = \frac{1 - a^2}{1 + a^2};$$

$$\therefore 2 \tan^{-1} a = \cos^{-1} \frac{1 - a^2}{1 + a^2}.$$

Similarly, the formula

$$\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$$

when expressed in the inverse notation becomes

$$3 \cos^{-1} a = \cos^{-1} (4a^3 - 3a).$$

**249.** *To prove that*

$$\tan^{-1} x + \tan^{-1} y = \tan^{-1} \frac{x+y}{1-xy}.$$

Let  $\tan^{-1} x = \alpha$ , so that  $\tan \alpha = x$ ;  
and  $\tan^{-1} y = \beta$ , so that  $\tan \beta = y$ .

We require  $\alpha + \beta$  in the form of an inverse tangent.

$$\begin{aligned} \text{Now } \tan (\alpha + \beta) &= \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} \\ &= \frac{x + y}{1 - xy}; \end{aligned}$$

$$\therefore \alpha + \beta = \tan^{-1} \frac{x+y}{1-xy};$$

that is,  $\tan^{-1} x + \tan^{-1} y = \tan^{-1} \frac{x+y}{1-xy}.$

By putting  $y = x$ , we obtain

$$2 \tan^{-1} x = \tan^{-1} \frac{2x}{1-x^2}.$$

**NOTE.** It is useful to remember that

$$\tan (\tan^{-1} x + \tan^{-1} y) = \frac{x+y}{1-xy}.$$



*Example 1.* Prove that

$$\tan^{-1}5 - \tan^{-1}3 + \tan^{-1}\frac{7}{9} = n\pi + \frac{\pi}{4}.$$

$$\begin{aligned} \text{The first side} &= \tan^{-1}\frac{5-3}{1+15} + \tan^{-1}\frac{7}{9} \\ &= \tan^{-1}\frac{1}{8} + \tan^{-1}\frac{7}{9} \\ &= \tan^{-1}\frac{\frac{1}{8} + \frac{7}{9}}{1 - \frac{7}{72}} = \tan^{-1}1 \\ &= n\pi + \frac{\pi}{4}. \end{aligned}$$

NOTE. The value of  $n$  cannot be assigned until we have selected some particular values for the angles  $\tan^{-1}5$ ,  $\tan^{-1}3$ ,  $\tan^{-1}\frac{7}{9}$ . If we choose the *principal values*, then  $n=0$ .

*Example 2.* Prove that

$$\sin^{-1}\frac{4}{5} + \cos^{-1}\frac{12}{13} + \sin^{-1}\frac{16}{65} = \frac{\pi}{2}.$$

We may write this identity in the form

$$\sin^{-1}\frac{4}{5} + \cos^{-1}\frac{12}{13} = \frac{\pi}{2} - \sin^{-1}\frac{16}{65} = \cos^{-1}\frac{16}{65}.$$

Let  $\alpha = \sin^{-1}\frac{4}{5}$ , so that  $\sin \alpha = \frac{4}{5}$ ;



and  $\beta = \cos^{-1}\frac{12}{13}$ , so that  $\cos \beta = \frac{12}{13}$ .

We have to express  $\alpha + \beta$  as an inverse cosine.

Now  $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$ ;  
whence by reading off the values of the functions from the figures in the margin, we have

$$\begin{aligned} \cos(\alpha + \beta) &= \frac{3}{5} \cdot \frac{12}{13} - \frac{4}{5} \cdot \frac{5}{13} \\ &= \frac{16}{65}; \end{aligned}$$

$$\therefore \alpha + \beta = \cos^{-1}\frac{16}{65}.$$



It is sometimes convenient to work entirely in terms of the tangent or cotangent.

*Example 3.* Prove that



$$2 \cot^{-1} 7 + \cos^{-1} \frac{3}{5} = \operatorname{cosec}^{-1} \frac{125}{117}.$$

$$\text{The first side} = \cot^{-1} \frac{7^2 - 1}{2 \times 7} + \cot^{-1} \frac{3}{4}$$

$$= \cot^{-1} \frac{24}{7} + \cot^{-1} \frac{3}{4}$$

$$= \cot^{-1} \frac{\frac{24}{7} \times \frac{3}{4} - 1}{\frac{24}{7} + \frac{3}{4}}$$

$$= \cot^{-1} \frac{44}{117} = \operatorname{cosec}^{-1} \frac{125}{117}.$$



### EXAMPLES. XIX. c.

Prove the following statements :

1.  $\sin^{-1} \frac{12}{13} = \cot^{-1} \frac{5}{12}.$

2.  $\operatorname{cosec}^{-1} \frac{17}{8} = \tan^{-1} \frac{8}{15}.$

3.  $\sec(\tan^{-1} x) = \sqrt{1+x^2}.$

4.  $2 \tan^{-1} \frac{1}{3} = \tan^{-1} \frac{3}{4}.$

5.  $\tan^{-1} \frac{4}{3} - \tan^{-1} 1 = \tan^{-1} \frac{1}{7}.$

6.  $\tan^{-1} \frac{2}{11} + \cot^{-1} \frac{24}{7} = \tan^{-1} \frac{1}{2}.$

7.  $\cot^{-1} \frac{4}{3} - \cot^{-1} \frac{15}{8} = \cot^{-1} \frac{84}{13}.$

8.  $2 \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{4} = \tan^{-1} \frac{32}{43}.$

9.  $\tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3} = \tan^{-1} \frac{5}{6} + \tan^{-1} \frac{1}{11}.$

10.  $\tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} + \tan^{-1} \frac{1}{18} = \cot^{-1} 3.$

$$11. \tan^{-1} \frac{3}{5} + \sin^{-1} \frac{3}{5} = \tan^{-1} \frac{27}{11}.$$

$$12. 2 \cot^{-1} \frac{5}{4} = \tan^{-1} \frac{40}{9}. \quad 13. 2 \tan^{-1} \frac{8}{15} = \sin^{-1} \frac{240}{289}.$$

$$14. \sin(2 \sin^{-1} x) = 2x \sqrt{1-x^2}.$$

$$15. \cos^{-1} x = 2 \sin^{-1} \sqrt{\frac{1-x}{2}}.$$

$$16. 2 \tan^{-1} \sqrt{\frac{x}{a}} = \cos^{-1} \frac{a-x}{a+x}.$$

$$17. 2 \tan^{-1} \frac{1}{8} + \tan^{-1} \frac{1}{7} + 2 \tan^{-1} \frac{1}{5} = \frac{\pi}{4}.$$

$$18. \sin^{-1} a - \cos^{-1} b = \cos^{-1} \{b \sqrt{1-a^2} + a \sqrt{1-b^2}\}.$$

$$19. \sin^{-1} \frac{4}{5} + \cos^{-1} \frac{2}{\sqrt{5}} = \cot^{-1} \frac{2}{11}.$$

$$20. \cos^{-1} \frac{63}{65} + 2 \tan^{-1} \frac{1}{5} = \sin^{-1} \frac{3}{5}.$$

$$21. \tan^{-1} m + \tan^{-1} n = \cos^{-1} \frac{1-mn}{\sqrt{(1+m^2)(1+n^2)}}.$$

$$22. \cos^{-1} \frac{20}{29} - \tan^{-1} \frac{16}{63} = \cos^{-1} \frac{1596}{1885}.$$

$$23. \cos^{-1} \sqrt{\frac{2}{3}} - \cos^{-1} \frac{\sqrt{6+1}}{2\sqrt{3}} = \frac{\pi}{6}.$$

$$24. \tan(2 \tan^{-1} x) = 2 \tan(\tan^{-1} x + \tan^{-1} x^3).$$

$$25. \tan^{-1} a = \tan^{-1} \frac{a-b}{1+ab} + \tan^{-1} \frac{b-c}{1+bc} + \tan^{-1} c.$$

$$26. \text{ If } \tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \pi, \text{ prove that } \\ x + y + z = xyz.$$

$$27. \text{ If } u = \cot^{-1} \sqrt{\cos a} - \tan^{-1} \sqrt{\cos a}, \text{ prove that } \\ \sin u = \tan^2 \frac{a}{2}.$$

250. We shall now shew how to solve equations expressed in the inverse notation.

*Example 1.* Solve  $\tan^{-1} 2x + \tan^{-1} 3x = n\pi + \frac{3\pi}{4}$ .

We have  $\tan^{-1} \frac{2x+3x}{1-6x^2} = n\pi + \frac{3\pi}{4}$ ;

$$\therefore \frac{2x+3x}{1-6x^2} = \tan \left( n\pi + \frac{3\pi}{4} \right) = -1;$$

$$\therefore 6x^2 - 5x - 1 = 0, \text{ or } (6x+1)(x-1) = 0;$$

$$\therefore x = 1, \text{ or } -\frac{1}{6}.$$

*Example 2.* Solve  $\sin^{-1} x + \sin^{-1} (1-x) = \cos^{-1} x$ .

By transposition,  $\sin^{-1} (1-x) = \cos^{-1} x - \sin^{-1} x$ .

Let  $\cos^{-1} x = \alpha$ , and  $\sin^{-1} x = \beta$ ; then

$$\sin^{-1} (1-x) = \alpha - \beta;$$

$$\therefore 1-x = \sin (\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta.$$

But  $\cos \alpha = x$ , and therefore  $\sin \alpha = \sqrt{1-x^2}$ ;

also  $\sin \beta = x$ , and therefore  $\cos \beta = \sqrt{1-x^2}$ ;

$$\therefore 1-x = (1-x^2) - x^2 = 1-2x^2;$$

$$\therefore 2x^2 - x = 0;$$

whence  $x = 0$ , or  $\frac{1}{2}$ .

### EXAMPLES. XIX. d.

Solve the equations :

1.  $\sin^{-1} x = \cos^{-1} x$ .                      2.  $\tan^{-1} x = \cot^{-1} x$ .

3.  $\tan^{-1} (x+1) - \tan^{-1} (x-1) = \cot^{-1} 2$ .

4.  $\cot^{-1} x + \cot^{-1} 2x = \frac{3\pi}{4}$ .

5.  $\sin^{-1} x - \cos^{-1} x = \sin^{-1} (3x-2)$ .

6.  $\cos^{-1} x - \sin^{-1} x = \cos^{-1} x\sqrt{3}$ .

$$7. \quad \tan^{-1} \frac{x-1}{x-2} + \tan^{-1} \frac{x+1}{x+2} = \frac{\pi}{4}.$$

$$8. \quad 2 \cot^{-1} 2 + \cos^{-1} \frac{3}{5} = \operatorname{cosec}^{-1} x.$$

$$9. \quad \tan^{-1} x + \tan^{-1} (1-x) = 2 \tan^{-1} \sqrt{x-x^2}.$$

$$10. \quad \cos^{-1} \frac{1-a^2}{1+a^2} - \cos^{-1} \frac{1-b^2}{1+b^2} = 2 \tan^{-1} x.$$

$$11. \quad \sin^{-1} \frac{2a}{1+a^2} + \tan^{-1} \frac{2x}{1-x^2} = \cos^{-1} \frac{1-b^2}{1+b^2}.$$

$$12. \quad \cot^{-1} \frac{x^2-1}{2x} + \tan^{-1} \frac{2x}{x^2-1} + \frac{4\pi}{3} = 0.$$

13. Shew that we can express

$$\sin^{-1} \frac{2ab}{a^2+b^2} + \sin^{-1} \frac{2cd}{c^2+d^2} \text{ in the form } \sin^{-1} \frac{2xy}{x^2+y^2}$$

where  $x$  and  $y$  are rational functions of  $a, b, c, d$ .

$$14. \quad \text{If } \sin [2 \cos^{-1} \{\cot (2 \tan^{-1} x)\}] = 0, \text{ find } x.$$

$$15. \quad \text{If } 2 \tan^{-1} (\cos \theta) = \tan^{-1} (2 \operatorname{cosec} \theta), \text{ find } \theta.$$

$$16. \quad \text{If } \sin (\pi \cos \theta) = \cos (\pi \sin \theta), \text{ shew that}$$

$$2\theta = \pm \sin^{-1} \frac{3}{4}.$$

17. If  $\sin (\pi \cot \theta) = \cos (\pi \tan \theta)$ , and  $n$  is any integer, shew that either  $\cot 2\theta$  or  $\operatorname{cosec} 2\theta$  is of the form  $\frac{4n+1}{4}$ .

18. If  $\tan (\pi \cot \theta) = \cot (\pi \tan \theta)$ , and  $n$  is any integer, shew that

$$\tan \theta = \frac{2n+1}{4} \pm \frac{\sqrt{4n^2+4n-15}}{4}.$$

19. Find all the positive integral solutions of

$$\tan^{-1} x + \cot^{-1} y = \tan^{-1} 3.$$

### MISCELLANEOUS EXAMPLES. G.

1. If the sines of the angles of a triangle are in the ratio of 4 : 5 : 6, shew that the cosines are in the ratio of 12 : 9 : 2.

2. Solve the equations :

$$(1) \quad 2 \cos^3 \theta + \sin^2 \theta - 1 = 0; \quad (2) \quad \sec^3 \theta - 2 \tan^2 \theta = 2.$$

3. If  $\tan \beta = 2 \sin \alpha \sin \gamma \operatorname{cosec} (\alpha + \gamma)$ , prove that  $\cot \alpha$ ,  $\cot \beta$ ,  $\cot \gamma$  are in arithmetical progression.

4. In a triangle shew that

$$4r(r_1 + r_2 + r_3) = 2(bc + ca + ab) - (a^2 + b^2 + c^2).$$

5. Prove that

$$(1) \quad \tan^{-1} \frac{1}{3} - \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} = \tan^{-1} \frac{3}{11};$$

$$(2) \quad \sin^{-1} \frac{3}{5} + \sin^{-1} \frac{8}{17} + \sin^{-1} \frac{36}{85} = \frac{\pi}{2}.$$

6. Find the greatest angle of the triangle whose sides are 185, 222, 259; given  $\log 6 = .7781513$ ,

$$L \cos 39^\circ 14' = 9.8890644, \text{ diff. for } 1' = 1032.$$

7. If  $\tan(\alpha + \theta) = n \tan(\alpha - \theta)$ , prove that  $\frac{\sin 2\theta}{\sin 2\alpha} = \frac{n-1}{n+1}$ .

8. If in a triangle  $8R^2 = a^2 + b^2 + c^2$ , prove that one of the angles is a right angle.

9. The area of a regular polygon of  $n$  sides inscribed in a circle is three-fourths of the area of the circumscribed regular polygon with the same number of sides: find  $n$ .

10.  $ABCD$  is a straight sea-wall. From  $B$  the straight lines drawn to two boats are each inclined at  $45^\circ$  to the direction of the wall, and from  $C$  the angles of inclination are  $15^\circ$  and  $75^\circ$ . If  $BC = 400$  yards, find the distance between the boats, and the distance of each from the sea-wall.



## CHAPTER XX.

### FUNCTIONS OF SUBMULTIPLE ANGLES.

#### 251. Trigonometrical ratios of $22\frac{1}{2}^\circ$ or $\frac{\pi}{8}$ .

From the identity

$$2 \sin^2 22\frac{1}{2}^\circ = 1 - \cos 45^\circ,$$

we have

$$4 \sin^2 22\frac{1}{2}^\circ = 2 - 2 \cos 45^\circ = 2 - \sqrt{2};$$

$$\therefore 2 \sin 22\frac{1}{2}^\circ = \sqrt{2 - \sqrt{2}} \dots\dots\dots(1).$$

In like manner from

$$2 \cos^2 22\frac{1}{2}^\circ = 1 + \cos 45^\circ,$$

we obtain

$$2 \cos 22\frac{1}{2}^\circ = \sqrt{2 + \sqrt{2}} \dots\dots\dots(2).$$

In each of these cases the positive sign must be taken before the radical, since  $22\frac{1}{2}^\circ$  is an acute angle.

$$\text{Again, } \tan 22\frac{1}{2}^\circ = \frac{1 - \cos 45^\circ}{\sin 45^\circ} = \operatorname{cosec} 45^\circ - \cot 45^\circ;$$

$$\therefore \tan 22\frac{1}{2}^\circ = \sqrt{2} - 1.$$

#### 252. We have seen that $2 \cos \frac{\pi}{8} = \sqrt{2 + \sqrt{2}}$ ;

but

$$4 \cos^2 \frac{\pi}{16} = 2 + 2 \cos \frac{\pi}{8};$$

$$\therefore 4 \cos^2 \frac{\pi}{16} = 2 + \sqrt{2 + \sqrt{2}};$$

$$\therefore 2 \cos \frac{\pi}{16} = \sqrt{2 + \sqrt{2 + \sqrt{2}}}.$$

Similarly,

$$2 \cos \frac{\pi}{32} = \sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2}}}};$$

and so on.

253. Suppose that  $\cos A = \frac{1}{2}$  and that it is required to find  $\sin \frac{A}{2}$ .

$$\sin \frac{A}{2} = \sqrt{\frac{1 - \cos A}{2}} = \sqrt{\frac{1}{2} \left(1 - \frac{1}{2}\right)} = \sqrt{\frac{1}{4}} = \pm \frac{1}{2}.$$

This case differs from those of the two previous articles in that the datum is less precise. All we know of the angle  $A$  is contained in the statement that its cosine is equal to  $\frac{1}{2}$ , and without some further knowledge respecting  $A$  we cannot remove the ambiguity of sign in the value found for  $\sin \frac{A}{2}$ .

We now proceed to a more general discussion.

254. Given  $\cos A$  to find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  and to explain the presence of the two values in each case.

From the identities

$$2 \sin^2 \frac{A}{2} = 1 - \cos A, \text{ and } 2 \cos^2 \frac{A}{2} = 1 + \cos A,$$

we have

$$\sin \frac{A}{2} = \pm \sqrt{\frac{1 - \cos A}{2}}, \text{ and } \cos \frac{A}{2} = \pm \sqrt{\frac{1 + \cos A}{2}}.$$

Thus corresponding to *one* value of  $\cos A$ , there are *two* values for  $\sin \frac{A}{2}$ , and *two* values for  $\cos \frac{A}{2}$ .

The presence of these two values may be explained as follows. If  $\cos A$  is given and nothing further is stated about the angle  $A$ , all we know is that  $A$  belongs to a certain group of *equi-cosinal angles*. Let  $a$  be the smallest positive angle belonging to this group, then  $A = 2n\pi \pm a$ . Thus in finding  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  we are really finding the values of

$$\sin \frac{1}{2}(2n\pi \pm a) \text{ and } \cos \frac{1}{2}(2n\pi \pm a).$$

$$\begin{aligned}
 \text{Now} \quad \sin \frac{1}{2}(2n\pi \pm a) &= \sin \left( n\pi \pm \frac{a}{2} \right) \\
 &= \sin n\pi \cos \frac{a}{2} \pm \cos n\pi \sin \frac{a}{2} \\
 &= \pm \sin \frac{a}{2},
 \end{aligned}$$

for  $\sin n\pi = 0$  and  $\cos n\pi = \pm 1$ .

$$\begin{aligned}
 \text{Again,} \quad \cos \frac{1}{2}(2n\pi \pm a) &= \cos n\pi \cos \frac{a}{2} \mp \sin n\pi \sin \frac{a}{2} \\
 &= \pm \cos \frac{a}{2}.
 \end{aligned}$$

Thus there are two values for  $\sin \frac{A}{2}$  and two values for  $\cos \frac{A}{2}$  when  $\cos A$  is given and nothing further is known respecting  $A$ .

**255. Geometrical Illustration.** Let  $a$  be the smallest positive angle which has the same cosine as  $A$ ; then


$$A = 2n\pi \pm a,$$

and we have to find the sine

and cosine of  $\frac{A}{2}$ , that is of

$$n\pi \pm \frac{a}{2}.$$

Each of the angles denoted by this formula is bounded by one of the lines  $OP_1, OP_2, OP_3, OP_4$ . Now



$$\sin XOP_2 = \sin \frac{a}{2}, \quad \sin XOP_3 = -\sin \frac{a}{2}, \quad \sin XOP_4 = -\sin \frac{a}{2},$$

$$\cos XOP_2 = -\cos \frac{a}{2}, \quad \cos XOP_3 = -\cos \frac{a}{2}, \quad \cos XOP_4 = \cos \frac{a}{2}.$$

Thus the values of  $\sin \frac{A}{2}$  are  $\pm \sin \frac{a}{2}$ , and the values of  $\cos \frac{A}{2}$  are  $\pm \cos \frac{a}{2}$ .

**256.** If  $\cos A$  is given, and  $A$  lies between certain known limits, the ambiguities of sign in the formulæ of Art. 254 may be removed.

*Example.* If  $\cos A = -\frac{7}{25}$ , and  $A$  lies between  $450^\circ$  and  $540^\circ$ , find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$ .

$$\sin \frac{A}{2} = \sqrt{\frac{1 - \cos A}{2}} = \sqrt{\frac{1}{2} \left(1 + \frac{7}{25}\right)} = \sqrt{\frac{16}{25}} = \pm \frac{4}{5};$$

$$\cos \frac{A}{2} = \sqrt{\frac{1 + \cos A}{2}} = \sqrt{\frac{1}{2} \left(1 - \frac{7}{25}\right)} = \sqrt{\frac{9}{25}} = \pm \frac{3}{5}.$$

Now  $\frac{A}{2}$  lies between  $225^\circ$  and  $270^\circ$ , so that  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  are both negative;

$$\therefore \sin \frac{A}{2} = -\frac{4}{5}, \text{ and } \cos \frac{A}{2} = -\frac{3}{5}.$$

**257.** To find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  in terms of  $\sin A$  and to explain the presence of four values in each case.

We have  $\sin^2 \frac{A}{2} + \cos^2 \frac{A}{2} = 1,$

and  $2 \sin \frac{A}{2} \cos \frac{A}{2} = \sin A.$

By addition,  $\left(\sin \frac{A}{2} + \cos \frac{A}{2}\right)^2 = 1 + \sin A;$

by subtraction,  $\left(\sin \frac{A}{2} - \cos \frac{A}{2}\right)^2 = 1 - \sin A.$

$$\therefore \sin \frac{A}{2} + \cos \frac{A}{2} = \pm \sqrt{1 + \sin A} \dots\dots\dots(1),$$

and  $\sin \frac{A}{2} - \cos \frac{A}{2} = \pm \sqrt{1 - \sin A} \dots\dots\dots(2).$

By addition and subtraction, we obtain  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$ ; and since there is a double sign before each radical, there are *four*

values for  $\sin \frac{A}{2}$ , and *four* values for  $\cos \frac{A}{2}$  corresponding to *one* value of  $\sin A$ .

The presence of these four values may be explained as follows.

If  $\sin A$  is given and nothing else is stated about the angle  $A$  all we know is that  $A$  belongs to a certain group of *equi-sin*al angles. Let  $a$  be the smallest positive angle belonging to this group, then  $A = n\pi + (-1)^n a$ . Thus in finding  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  we are really finding

$$\sin \frac{1}{2} \{n\pi + (-1)^n a\}, \text{ and } \cos \frac{1}{2} \{n\pi + (-1)^n a\}.$$

First suppose  $n$  even and equal to  $2m$ ; then

$$\begin{aligned} \sin \frac{1}{2} \{n\pi + (-1)^n a\} &= \sin \left( m\pi + \frac{a}{2} \right) \\ &= \sin m\pi \cos \frac{a}{2} + \cos m\pi \sin \frac{a}{2} \\ &= \pm \sin \frac{a}{2}, \end{aligned}$$

since  $\sin m\pi = 0$ , and  $\cos m\pi = \pm 1$ .

Next suppose  $n$  odd and equal to  $2m+1$ ; then

$$\begin{aligned} \sin \frac{1}{2} \{n\pi + (-1)^n a\} &= \sin \left( m\pi + \frac{\pi}{2} - \frac{a}{2} \right) \\ &= \sin m\pi \cos \left( \frac{\pi}{2} - \frac{a}{2} \right) + \cos m\pi \sin \left( \frac{\pi}{2} - \frac{a}{2} \right) \\ &= \pm \sin \left( \frac{\pi}{2} - \frac{a}{2} \right). \end{aligned}$$

Thus we have *four* values for  $\sin \frac{A}{2}$  when  $\sin A$  is given and nothing further is known respecting  $A$ .

In like manner it may be shewn that

$$\cos \frac{A}{2} \text{ has the } \textit{four} \text{ values } \pm \cos \frac{a}{2}, \quad \pm \cos \left( \frac{\pi}{2} - \frac{a}{2} \right).$$

**258. Geometrical Illustration.** Let  $a$  be the smallest positive angle which has the same sine as  $A$ ; then

$$A = n\pi + (-1)^n a,$$

and we have to find the sine and cosine of  $\frac{A}{2}$ , that is of

$$\frac{1}{2} \{n\pi + (-1)^n a\}.$$


If  $n$  is even and equal to  $2m$ , this expression becomes  $m\pi + \frac{a}{2}$ .

If  $n$  is odd and equal to  $2m+1$ , the expression becomes

$$m\pi + \left(\frac{\pi}{2} - \frac{a}{2}\right).$$

The angles denoted by the formula  $m\pi + \frac{a}{2}$  are bounded by one of the lines  $OP_1$  or  $OP_2$ ; and

those denoted by the formula  $m\pi + \left(\frac{\pi}{2} - \frac{a}{2}\right)$  are bounded by one of the lines  $OP_3$  or  $OP_4$ .



$$\text{Now} \quad \sin \angle OP_1 = \sin \frac{a}{2};$$

$$\sin \angle OP_2 = -\sin \angle OP_1 = -\sin \frac{a}{2};$$

$$\sin \angle OP_3 = \sin \left(\frac{\pi}{2} - \frac{a}{2}\right);$$

$$\sin \angle OP_4 = -\sin \angle OP_3 = -\sin \left(\frac{\pi}{2} - \frac{a}{2}\right).$$

Thus the values of  $\sin \frac{A}{2}$  are

$$\pm \sin \frac{a}{2} \quad \text{and} \quad \pm \sin \left(\frac{\pi}{2} - \frac{a}{2}\right).$$


Similarly the values of  $\cos \frac{A}{2}$  are  $\pm \cos \frac{a}{2}$  and  $\pm \cos \left(\frac{\pi}{2} - \frac{a}{2}\right)$ .



**259.** If in addition to the value of  $\sin A$  we know that  $A$  lies between certain limits, the ambiguities of sign in equations (1) and (2) of Art. 257 may be removed.

*Example 1.* Find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  in terms of  $\sin A$  when  $A$  lies between  $450^\circ$  and  $630^\circ$ .

In this case  $\frac{A}{2}$  lies between  $225^\circ$  and  $315^\circ$ .  
From the adjoining figure it is evident that between these limits  $\sin \frac{A}{2}$  is greater than  $\cos \frac{A}{2}$  and is negative.



$$\therefore \sin \frac{A}{2} + \cos \frac{A}{2} = -\sqrt{1 + \sin A},$$

and  $\sin \frac{A}{2} - \cos \frac{A}{2} = -\sqrt{1 - \sin A}.$

$$\therefore 2 \sin \frac{A}{2} = -\sqrt{1 + \sin A} - \sqrt{1 - \sin A},$$

and  $2 \cos \frac{A}{2} = -\sqrt{1 + \sin A} + \sqrt{1 - \sin A}.$

*Example 2.* Determine the limits between which  $A$  must lie in order that

$$2 \cos A = -\sqrt{1 + \sin 2A} - \sqrt{1 - \sin 2A}.$$

The given relation is obtained by combining


$$\sin A + \cos A = -\sqrt{1 + \sin 2A} \dots\dots\dots(1),$$

and  $\sin A - \cos A = +\sqrt{1 - \sin 2A} \dots\dots\dots(2).$

From (1), we see that of  $\sin A$  and  $\cos A$  the numerically greater is negative.

From (2), we see that the cosine is the greater.

Hence we have to choose limits between which  $\cos A$  is numerically greater than  $\sin A$  and is negative. From the figure we see that  $A$  lies between  $2n\pi + \frac{3\pi}{4}$  and  $2n\pi + \frac{5\pi}{4}.$



*Example 3.* Trace the changes of  $\cos \theta - \sin \theta$  in sign and magnitude as  $\theta$  increases from 0 to  $2\pi$ .

$$\begin{aligned}\cos \theta - \sin \theta &= \sqrt{2} \left( \frac{1}{\sqrt{2}} \cos \theta - \frac{1}{\sqrt{2}} \sin \theta \right) \\ &= \sqrt{2} \left( \cos \theta \cos \frac{\pi}{4} - \sin \theta \sin \frac{\pi}{4} \right) \\ &= \sqrt{2} \cos \left( \theta + \frac{\pi}{4} \right).\end{aligned}$$

As  $\theta$  increases from 0 to  $\frac{\pi}{4}$ , the expression is positive and decreases from 1 to 0.

As  $\theta$  increases from  $\frac{\pi}{4}$  to  $\frac{3\pi}{4}$ , the expression is negative and increases numerically from 0 to  $-\sqrt{2}$ .

As  $\theta$  increases from  $\frac{3\pi}{4}$  to  $\frac{5\pi}{4}$ , the expression is negative and decreases numerically from  $-\sqrt{2}$  to 0.

As  $\theta$  increases from  $\frac{5\pi}{4}$  to  $\frac{7\pi}{4}$ , the expression is positive and increases from 0 to  $\sqrt{2}$ .

As  $\theta$  increases from  $\frac{7\pi}{4}$  to  $2\pi$ , the expression is positive and decreases from  $\sqrt{2}$  to 1.

**260.** To find the sine and cosine of  $9^\circ$ .

Since  $\cos 9^\circ > \sin 9^\circ$  and is positive, we have

$$\sin 9^\circ + \cos 9^\circ = +\sqrt{1 + \sin 18^\circ},$$

and  $\sin 9^\circ - \cos 9^\circ = -\sqrt{1 - \sin 18^\circ}.$

$$\therefore \sin 9^\circ + \cos 9^\circ = +\sqrt{1 + \frac{\sqrt{5}-1}{4}} = +\frac{1}{2}\sqrt{3+\sqrt{5}},$$

and  $\sin 9^\circ - \cos 9^\circ = -\sqrt{1 - \frac{\sqrt{5}-1}{4}} = -\frac{1}{2}\sqrt{5-\sqrt{5}}.$

$$\therefore \sin 9^\circ = \frac{1}{4} \{ \sqrt{3+\sqrt{5}} - \sqrt{5-\sqrt{5}} \},$$

and  $\cos 9^\circ = \frac{1}{4} \{ \sqrt{3+\sqrt{5}} + \sqrt{5-\sqrt{5}} \}.$

## EXAMPLES. XX. a.

1. When  $A$  lies between  $-270^\circ$  and  $-360^\circ$ , prove that

$$\sin \frac{A}{2} = -\sqrt{\frac{1 - \cos A}{2}}.$$

2. If  $\cos A = \frac{119}{169}$ , find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  when  $A$  lies between  $270^\circ$  and  $360^\circ$ .

3. If  $\cos A = -\frac{161}{289}$ , find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  when  $A$  lies between  $540^\circ$  and  $630^\circ$ .

4. Find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  in terms of  $\sin A$  when  $A$  lies between  $270^\circ$  and  $450^\circ$ .

5. Find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  in terms of  $\sin A$  when  $\frac{A}{2}$  lies between  $225^\circ$  and  $315^\circ$ .

6. Find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  in terms of  $\sin A$  when  $A$  lies between  $-450^\circ$  and  $-630^\circ$ .

7. If  $\sin A = \frac{24}{25}$ , find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  when  $A$  lies between  $90^\circ$  and  $180^\circ$ .

8. If  $\sin A = -\frac{240}{289}$ , find  $\sin \frac{A}{2}$  and  $\cos \frac{A}{2}$  when  $A$  lies between  $270^\circ$  and  $360^\circ$ .

9. Determine the limits between which  $A$  must lie in order that

$$(1) \quad 2 \sin A = \sqrt{1 + \sin 2A} - \sqrt{1 - \sin 2A};$$

$$(2) \quad 2 \cos A = -\sqrt{1 + \sin 2A} + \sqrt{1 - \sin 2A};$$

$$(3) \quad 2 \sin A = -\sqrt{1 + \sin 2A} + \sqrt{1 - \sin 2A}.$$

10. If  $A = 240^\circ$ , is the following statement correct?

$$2 \sin \frac{A}{2} = \sqrt{1 + \sin A} - \sqrt{1 - \sin A}.$$

If not, how must it be modified?

11. Prove that

$$(1) \quad \tan 7\frac{1}{2}^\circ = \sqrt{6} - \sqrt{3} + \sqrt{2} - 2;$$

$$(2) \quad \cot 142\frac{1}{2}^\circ = \sqrt{2} + \sqrt{3} - 2 - \sqrt{6}.$$

12. Shew that  $\sin 9^\circ$  lies between  $\cdot 156$  and  $\cdot 157$ .

13. Prove that

$$(1) \quad 2 \sin 11^\circ 15' = \sqrt{2 - \sqrt{2 + \sqrt{2}}};$$

$$(2) \quad \tan 11^\circ 15' = \sqrt{4 + 2\sqrt{2}} - (\sqrt{2} + 1).$$

14. When  $\theta$  varies from 0 to  $2\pi$  trace the changes in sign and magnitude of

$$(1) \quad \cos \theta + \sin \theta;$$

$$(2) \quad \sin \theta - \sqrt{3} \cos \theta.$$

15. When  $\theta$  varies from 0 to  $\pi$ , trace the changes in sign and magnitude of

$$(1) \quad \frac{\tan \theta + \cot \theta}{\tan \theta - \cot \theta};$$

$$(2) \quad \frac{2 \sin \theta - \sin 2\theta}{2 \sin \theta + \sin 2\theta}.$$

261. To find  $\tan \frac{A}{2}$  when  $\tan A$  is given and to explain the presence of the two values.

Denote  $\tan A$  by  $t$ ; then

$$t = \tan A = \frac{2 \tan \frac{A}{2}}{1 - \tan^2 \frac{A}{2}};$$

$$\therefore t \tan^2 \frac{A}{2} + 2 \tan \frac{A}{2} - t = 0;$$

$$\therefore \tan \frac{A}{2} = \frac{-2 \pm \sqrt{4 + 4t^2}}{2t} = \frac{-1 \pm \sqrt{1 + t^2}}{t}.$$

The presence of these two values may be explained as follows.

If  $\alpha$  be the smallest positive angle which has the given tangent, then  $A = n\pi + \alpha$ , and we are really finding the value of

$$\tan \frac{1}{2}(n\pi + \alpha).$$

(1) Let  $n$  be even and equal to  $2m$ ; then

$$\tan \frac{1}{2}(n\pi + \alpha) = \tan \left( m\pi + \frac{\alpha}{2} \right) = \tan \frac{\alpha}{2}.$$

(2) Let  $n$  be odd and equal to  $2m+1$ ; then

$$\tan \frac{1}{2}(n\pi + \alpha) = \tan \left( m\pi + \frac{\pi}{2} + \frac{\alpha}{2} \right) = \tan \left( \frac{\pi}{2} + \frac{\alpha}{2} \right).$$

Thus  $\tan \frac{A}{2}$  has the two values  $\tan \frac{\alpha}{2}$  and  $\tan \left( \frac{\pi}{2} + \frac{\alpha}{2} \right)$ .

*Example 1.* If  $A = 170^\circ$ , prove that  $\tan \frac{A}{2} = \frac{-1 - \sqrt{1 + \tan^2 A}}{\tan A}$ .

Here  $\frac{A}{2}$  is an acute angle, so that  $\tan \frac{A}{2}$  must be positive. Hence

in the formula  $\frac{-1 \pm \sqrt{1 + \tan^2 A}}{\tan A}$  the numerator must have the same sign as the denominator. But when  $A = 170^\circ$ ,  $\tan A$  is negative, and therefore we must choose the sign which will make the numerator negative; thus

$$\tan \frac{A}{2} = \frac{-1 - \sqrt{1 + \tan^2 A}}{\tan A}.$$

*Example 2.* Given  $\cos A = .6$ , find  $\tan \frac{A}{2}$ , and explain the double answer.

$$\tan^2 \frac{A}{2} = \frac{1 - \cos A}{1 + \cos A} = \frac{.4}{1.6} = \frac{1}{4};$$

$$\therefore \tan \frac{A}{2} = \pm \frac{1}{2}.$$

Here all we know of the angle  $A$  is that it must be one of a group of equi-cosinal angles. Let  $\alpha$  be the smallest positive angle of this group; then  $A = 2n\pi \pm \alpha$ .

$$\therefore \tan \frac{A}{2} = \tan \left( n\pi \pm \frac{\alpha}{2} \right) = \tan \left( \pm \frac{\alpha}{2} \right) = \pm \tan \frac{\alpha}{2}.$$

Thus we have two values differing only in sign.

**262.** When any one of the functions of an acute angle  $A$  is given, we may in some cases conveniently obtain the functions of  $\frac{A}{2}$ , as in the following example.


*Example.* Given  $\cos A = \frac{b}{a}$ , to find the functions of  $\frac{A}{2}$ .

Make a right-angled triangle  $PQR$  in which the hypotenuse  $PQ = a$ , and base  $QR = b$ ; then

$$\cos PQR = \frac{QR}{PQ} = \frac{b}{a} = \cos A;$$

$$\therefore \angle PQR = A.$$

Produce  $RQ$  to  $S$  making  $QS = QP$ ;



$$\therefore \angle PSQ = \angle SPQ = \frac{1}{2} \angle PQR = \frac{A}{2}.$$

Now  $SR = a + b$ , and  $PR = \sqrt{a^2 - b^2}$ ,

$$\therefore PS^2 = (a + b)^2 + (a^2 - b^2) = 2a^2 + 2ab;$$

$$\therefore PS = \sqrt{2a(a + b)}.$$

The functions of  $\frac{A}{2}$  may now be written down in terms of the sides of the triangle  $PRS$ .

**263.** From Art. 125, we have

$$\cos A = 4 \cos^3 \frac{A}{3} - 3 \cos \frac{A}{3}.$$

Thus it appears that if  $\cos A$  be given we have a *cubic* equation to find  $\cos \frac{A}{3}$ ; so that  $\cos \frac{A}{3}$  has *three* values.

Similarly, from the equation

$$\sin A = 3 \sin \frac{A}{3} - 4 \sin^3 \frac{A}{3}$$

it appears that corresponding to *one* value of  $\sin A$  there are *three* values of  $\sin \frac{A}{3}$ .

It will be a useful exercise to prove these two statements analytically as in Arts. 254 and 257. In the next article we shall give a geometrical explanation for the case of the cosine.



**264.** Given  $\cos A$  to find  $\cos \frac{A}{3}$ , and to explain the presence of the three values.


Let  $a$  be the smallest positive angle with the given cosine; then  $A = 2n\pi \pm a$ , and we have to find all the values of

$$\cos \frac{1}{3}(2n\pi \pm a).$$

Consider the angles denoted by the formula

$$\frac{1}{3}(2n\pi \pm a),$$

and ascribe to  $n$  in succession the values 0, 1, 2, 3, .....



When  $n=0$ , the angles are  $\pm \frac{a}{3}$ , bounded by  $OP_1$  and  $OQ_1$ ;

when  $n=1$ , the angles are  $\frac{2\pi}{3} \pm \frac{a}{3}$ , bounded by  $OP_2$  and  $OQ_2$

when  $n=2$ , the angles are  $\frac{4\pi}{3} \pm \frac{a}{3}$ , bounded by  $OP_3$  and  $OQ_3$ .

By giving to  $n$  the values 3, 4, 5, ... we obtain a series of angles coterminal with those indicated in the figure.

Thus  $OP_1, OQ_1, OP_2, OQ_2, OP_3, OQ_3$  bound all the angles included in the formula  $\frac{1}{3}(2n\pi \pm a)$ .

Now  $\cos XOQ_1 = \cos XOP_1 = \cos \frac{a}{3};$

$$\cos XOP_3 = \cos XOQ_2 = \cos \left( \frac{2\pi}{3} - \frac{a}{3} \right);$$

$$\cos XOQ_3 = \cos XOP_2 = \cos \left( \frac{2\pi}{3} + \frac{a}{3} \right).$$

Thus the values of  $\cos \frac{A}{3}$  are  $\cos \frac{a}{3}, \cos \frac{2\pi+a}{3}, \cos \frac{2\pi-a}{3}.$

**EXAMPLES. XX. b.**

1. If  $A=320^\circ$ , prove that

$$\tan \frac{A}{2} = \frac{-1 + \sqrt{1 + \tan^2 A}}{\tan A}.$$

2. Shew that

$$\tan A = -\frac{1 + \sqrt{1 + \tan^2 2A}}{\tan 2A} \text{ when } A=110^\circ.$$

3. Find  $\tan A$  when  $\cos 2A = \frac{12}{13}$  and  $A$  lies between  $180^\circ$  and  $225^\circ$ .

4. Find  $\cot \frac{A}{2}$  when  $\cos A = -\frac{4}{5}$  and  $A$  lies between  $180^\circ$  and  $270^\circ$ .

5. If  $\cot 2\theta = \cot 2a$ , shew that  $\cot \theta$  has the two values  $\cot a$  and  $-\tan a$ .

6. Given that  $\sin \theta = \sin a$ , shew that the values of  $\sin \frac{\theta}{3}$  are

$$\sin \frac{a}{3}, \quad \sin \frac{\pi - a}{3}, \quad -\sin \frac{\pi + a}{3}.$$

7. If  $\tan \theta = \tan a$ , shew that the values of  $\tan \frac{\theta}{3}$  are

$$\tan \frac{a}{3}, \quad \tan \frac{\pi + a}{3}, \quad -\tan \frac{\pi - a}{3}.$$

8. Given that  $\cos 3\theta = \cos 3a$ , shew that the values of  $\sin \theta$  are

$$\pm \sin a, \quad -\sin \left( \frac{\pi}{3} \pm a \right), \quad \sin \left( \frac{2\pi}{3} \pm a \right).$$

9. Given that  $\sin 3\theta = \sin 3a$ , shew that the values of  $\cos \theta$  are

$$\pm \cos a, \quad \cos \left( \frac{\pi}{3} \pm a \right), \quad \cos \left( \frac{2\pi}{3} \pm a \right).$$


## CHAPTER XXI.

### LIMITS AND APPROXIMATIONS.

265. If  $\theta$  be the radian measure of an angle less than a right angle, to shew that  $\sin \theta$ ,  $\theta$ ,  $\tan \theta$  are in ascending order of magnitude.

Let the angle  $\theta$  be represented by  $\angle AOP$ .

With centre  $O$  and radius  $OA$  describe a circle. Draw  $PT$  at right angles to  $OP$  to meet  $OA$  produced in  $T$ , and join  $PA$ .



Let  $r$  be the radius of the circle.

$$\text{Area of } \triangle AOP = \frac{1}{2} AO \cdot OP \sin \angle AOP = \frac{1}{2} r^2 \sin \theta;$$

$$\text{area of sector } AOP = \frac{1}{2} r^2 \theta;$$

$$\text{area of } \triangle OPT = \frac{1}{2} OP \cdot PT = \frac{1}{2} r \cdot r \tan \theta = \frac{1}{2} r^2 \tan \theta.$$

But the areas of the triangle  $AOP$ , the sector  $AOP$ , and the triangle  $OPT$  are in ascending order of magnitude; that is,

$$\frac{1}{2} r^2 \sin \theta, \quad \frac{1}{2} r^2 \theta, \quad \frac{1}{2} r^2 \tan \theta$$

are in ascending order of magnitude;

$\therefore \sin \theta$ ,  $\theta$ ,  $\tan \theta$  are in ascending order of magnitude.

266. When  $\theta$  is indefinitely diminished, to prove that  $\frac{\sin \theta}{\theta}$  and  $\frac{\tan \theta}{\theta}$  each have unity for their limit.

In the last article, we have proved that  $\sin \theta$ ,  $\theta$ ,  $\tan \theta$  are in ascending order of magnitude. Divide each of these quantities by  $\sin \theta$ ; then

1,  $\frac{\theta}{\sin \theta}$ ,  $\frac{1}{\cos \theta}$  are in ascending order of magnitude;

that is,  $\frac{\theta}{\sin \theta}$  lies between 1 and  $\sec \theta$ .

But when  $\theta$  is indefinitely diminished, the limit of  $\sec \theta$  is 1; hence the limit of  $\frac{\theta}{\sin \theta}$  is 1; that is, the limit of  $\frac{\sin \theta}{\theta}$  is unity.

Again, by dividing each of the quantities  $\sin \theta$ ,  $\theta$ ,  $\tan \theta$  by  $\tan \theta$ , we find that  $\cos \theta$ ,  $\frac{\theta}{\tan \theta}$ , 1 are in ascending order of magnitude. Hence the limit of  $\frac{\tan \theta}{\theta}$  is unity.

These results are often written concisely in the forms

$$\lim_{\theta=0} \left( \frac{\sin \theta}{\theta} \right) = 1, \quad \lim_{\theta=0} \left( \frac{\tan \theta}{\theta} \right) = 1.$$

*Example.* Find the limit of  $n \sin \frac{\theta}{n}$  when  $n = \infty$ .

$$n \sin \frac{\theta}{n} = \theta \cdot \frac{n}{\theta} \cdot \sin \frac{\theta}{n} = \theta \left( \sin \frac{\theta}{n} \div \frac{\theta}{n} \right);$$

but since  $\frac{\theta}{n}$  is indefinitely small, the limit of  $\sin \frac{\theta}{n} \div \frac{\theta}{n}$  is unity;

$$\therefore \lim_{n=\infty} \left( n \sin \frac{\theta}{n} \right) = \theta.$$

Similarly

$$\lim_{n=\infty} \left( n \tan \frac{\theta}{n} \right) = \theta.$$

267. It is important to remember that the conclusions of the foregoing articles only hold when the angle is expressed in radian measure. If any other system of measurement is used, the results will require modification.

*Example.* Find the value of  $\text{Lt.}_{n=0} \left( \frac{\sin n^\circ}{n} \right)$ .

Let  $\theta$  be the number of radians in  $n^\circ$ ; then

$$\frac{n}{180} = \frac{\theta}{\pi}, \text{ and } n = \frac{180\theta}{\pi}; \text{ also } \sin n^\circ = \sin \theta;$$

$$\therefore \frac{\sin n^\circ}{n} = \frac{\pi \sin \theta}{180\theta} = \frac{\pi}{180} \cdot \frac{\sin \theta}{\theta}.$$

When  $n$  is indefinitely small,  $\theta$  is indefinitely small;

$$\therefore \text{Lt.}_{n=0} \left( \frac{\sin n^\circ}{n} \right) = \frac{\pi}{180} \cdot \text{Lt.}_{\theta=0} \left( \frac{\sin \theta}{\theta} \right);$$

$$\therefore \text{Lt.}_{n=0} \left( \frac{\sin n^\circ}{n} \right) = \frac{\pi}{180}.$$

268. When  $\theta$  is the radian measure of a very small angle, we have shewn that

$$\frac{\sin \theta}{\theta} = 1, \quad \cos \theta = 1, \quad \frac{\tan \theta}{\theta} = 1;$$

that is,  $\sin \theta = \theta, \quad \cos \theta = 1, \quad \tan \theta = \theta.$

Hence  $r \tan \theta = r\theta$ , and therefore in the figure of Art. 265, the tangent  $PT$  is equal to the arc  $PA$ , when  $\angle AOP$  is very small.

In Art. 270, it will be shewn that these results hold so long as  $\theta$  is so small that its square may be neglected. When this is the case, we have

$$\sin(a + \theta) = \sin a \cos \theta + \cos a \sin \theta$$

$$= \sin a + \theta \cos a;$$


$$\cos(a + \theta) = \cos a \cos \theta - \sin a \sin \theta$$

$$= \cos a - \theta \sin a.$$

*Example 1.* The inclination of a railway to the horizontal plane is  $52' 30''$ , find how many feet it rises in a mile.

Let  $OA$  be the horizontal plane, and  $OP$  a mile of the railway. Draw  $PN$  perpendicular to  $OA$ .

Let  $PN = x$  feet,  $\angle PON = \theta$ ;



then  $\frac{PN}{OP} = \sin \theta = \theta$  approximately.

But  $\theta = \text{radian measure of } 52' 30'' = \frac{52\frac{1}{2}}{60} \times \frac{\pi}{180} = \frac{7}{8} \times \frac{\pi}{180}$ ;

$$\therefore \frac{x}{1760 \times 3} = \frac{7}{8} \times \frac{22}{7} \times \frac{1}{180};$$

$$\therefore x = \frac{1760 \times 3 \times 22}{8 \times 180} = \frac{242}{3} = 80\frac{2}{3}.$$


Thus the rise is  $80\frac{2}{3}$  feet.

*Example 2.* A pole 6 ft. long stands on the top of a tower 54 ft. high: find the angle subtended by the pole at a point on the ground which is at a distance of 180 yds. from the foot of the tower.

Let  $A$  be the point on the ground,  $BC$  the tower,  $CD$  the pole.

Let  $\angle BAC = \alpha$ ,  $\angle CAD = \theta$ ;

then  $\tan \alpha = \frac{BC}{AB} = \frac{54}{540} = \frac{1}{10}$ ;



$$\tan (\alpha + \theta) = \frac{BD}{AB} = \frac{60}{540} = \frac{1}{9}.$$

But  $\tan (\alpha + \theta) = \frac{\tan \alpha + \tan \theta}{1 - \tan \alpha \tan \theta} = \frac{\tan \alpha + \theta}{1 - \theta \tan \alpha}$  approximately;

$$\therefore \frac{1}{9} = \frac{\frac{1}{10} + \theta}{1 - \frac{\theta}{10}} = \frac{1 + 10\theta}{10 - \theta};$$

whence  $\theta = \frac{1}{91}$ ; that is, the angle is  $\frac{1}{91}$  of a radian, and therefore contains  $\frac{1}{91} \times \frac{180}{\pi}$  degrees.

On reduction, we find that the angle is  $37' 46''$  nearly.



269. If  $\theta$  be the number of radians in an acute angle, to prove that  $\cos \theta > 1 - \frac{\theta^2}{2}$ , and  $\sin \theta > \theta - \frac{\theta^3}{4}$ .

Since  $\cos \theta = 1 - 2 \sin^2 \frac{\theta}{2}$ , and  $\sin \frac{\theta}{2} < \frac{\theta}{2}$ ;

$$\therefore \cos \theta > 1 - 2 \left( \frac{\theta}{2} \right)^2;$$

that is,  $\cos \theta > 1 - \frac{\theta^2}{2}$ .

Again,  $\sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} = 2 \tan \frac{\theta}{2} \cos^2 \frac{\theta}{2}$ ;

but  $\tan \frac{\theta}{2} > \frac{\theta}{2}$ ;

$$\therefore \sin \theta > 2 \frac{\theta}{2} \cos^2 \frac{\theta}{2};$$

$$\therefore \sin \theta > \theta \left( 1 - \sin^2 \frac{\theta}{2} \right).$$

But  $\sin \frac{\theta}{2} < \frac{\theta}{2}$ , and therefore

$$1 - \sin^2 \frac{\theta}{2} > 1 - \left( \frac{\theta}{2} \right)^2;$$

$$\therefore \sin \theta > \theta \left\{ 1 - \left( \frac{\theta}{2} \right)^2 \right\};$$

$$\therefore \sin \theta > \theta - \frac{\theta^3}{4}.$$

270. From the propositions established in this chapter, it follows that if  $\theta$  is an acute angle,

$$\cos \theta \text{ lies between } 1 \text{ and } 1 - \frac{\theta^2}{2},$$

and  $\sin \theta$  lies between  $\theta$  and  $\theta - \frac{\theta^3}{4}$ .

Thus  $\cos \theta = 1 - k\theta^2$  and  $\sin \theta = \theta - k'\theta^3$ , where  $k$  and  $k'$  are proper fractions less than  $\frac{1}{2}$  and  $\frac{1}{4}$  respectively.

Hence if  $\theta$  be so small that its square can be neglected,  
 $\cos \theta = 1, \quad \sin \theta = \theta.$

*Example.* Find the approximate value of  $\sin 10''$ .

The circular measure of  $10''$  is  $\frac{10\pi}{180 \times 60 \times 60}$  or  $\frac{\pi}{64800}$ ;

$$\therefore \sin 10'' < \frac{\pi}{64800} \text{ and } > \frac{\pi}{64800} - \frac{1}{4} \left( \frac{\pi}{64800} \right)^3.$$

But 
$$\frac{\pi}{64800} = \frac{3.1415926535...}{64800} = .000048481368.....;$$

$$\therefore \frac{\pi}{64800} < .00005 \text{ and } \left( \frac{\pi}{64800} \right)^3 < .000000000000125;$$

$$\therefore \sin 10'' < \frac{\pi}{64800} \text{ and } > \frac{\pi}{64800} - \frac{1}{4} (.000000000000125).$$

Hence to 12 places of decimals,

$$\sin 10'' = \frac{\pi}{64800} = .000048481368....$$

271. To shew that when  $n$  is an indefinitely large integer, the  
*limit of*  $\cos \frac{\theta}{2} \cos \frac{\theta}{4} \cos \frac{\theta}{8} \dots \cos \frac{\theta}{2^n} = \frac{\sin \theta}{\theta}.$

We have 
$$\begin{aligned} \sin \theta &= 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} \\ &= 2^2 \sin \frac{\theta}{4} \cos \frac{\theta}{4} \cos \frac{\theta}{2} \\ &= 2^3 \sin \frac{\theta}{8} \cos \frac{\theta}{8} \cos \frac{\theta}{4} \cos \frac{\theta}{2} \\ &\dots\dots\dots \\ &= 2^n \sin \frac{\theta}{2^n} \cos \frac{\theta}{2^n} \dots \cos \frac{\theta}{8} \cos \frac{\theta}{4} \cos \frac{\theta}{2}. \end{aligned}$$

$$\therefore \cos \frac{\theta}{2} \cos \frac{\theta}{4} \cos \frac{\theta}{8} \dots \cos \frac{\theta}{2^n} = \frac{\sin \theta}{2^n \sin \frac{\theta}{2^n}}.$$

But the limit of  $2^n \sin \frac{\theta}{2^n}$  is  $\theta$ , and thus the proposition is established. [See Art. 266.]

272. To shew that  $\frac{\sin \theta}{\theta}$  continually decreases from 1 to  $\frac{2}{\pi}$  as  $\theta$  continually increases from 0 to  $\frac{\pi}{2}$ .

We shall first shew that the fraction

$$\frac{\sin \theta}{\theta} - \frac{\sin (\theta + h)}{\theta + h} \text{ is positive,}$$

$h$  denoting the radian measure of a small positive angle.

$$\begin{aligned} \text{This fraction} &= \frac{(\theta + h) \sin \theta - \theta (\sin \theta \cos h + \cos \theta \sin h)}{\theta (\theta + h)} \\ &= \frac{\theta \sin \theta (1 - \cos h) + (h \sin \theta - \theta \cos \theta \sin h)}{\theta (\theta + h)}. \end{aligned}$$

Now  $\tan \theta > \theta$ , that is  $\sin \theta > \theta \cos \theta$ , and  $h > \sin h$ ;

$$\therefore h \sin \theta > \theta \cos \theta \sin h.$$

Also  $1 - \cos h$  is positive; hence the numerator is positive, and therefore the fraction is positive;

$$\therefore \frac{\sin (\theta + h)}{\theta + h} < \frac{\sin \theta}{\theta};$$

$\therefore \frac{\sin \theta}{\theta}$  continually decreases as  $\theta$  continually increases.

When  $\theta = 0$ ,  $\frac{\sin \theta}{\theta} = 1$ ; and when  $\theta = \frac{\pi}{2}$ ,  $\frac{\sin \theta}{\theta} = \frac{2}{\pi}$ .

Thus the proposition is established.

### EXAMPLES. XXI. a.

$$\left[ \text{In this Exercise take } \pi = \frac{22}{7}. \right]$$

1. A tower 44 feet high subtends an angle of  $35'$  at a point  $A$  on the ground: find the distance of  $A$  from the tower.

2. From the top of a wall 7 ft. 4 in. high the angle of depression of an object on the ground is  $24' 30''$ : find its distance from the wall.

3. Find the height of an object whose angle of elevation at a distance of 840 yards is  $1^{\circ} 30'$ .

4. Find the angle subtended by a pole 10 ft. 1 in. high at a distance of a mile.

5. Find the angle subtended by a circular target 4 feet in diameter at a distance of 1000 yards.

6. Taking the diameter of a penny as 1.25 inches, find at what distance it must be held from the eye so as just to hide the moon, supposing the diameter of the moon to be half a degree.

7. Find the distance at which a globe 11 inches in diameter subtends an angle of  $5'$ .

8. Two places on the same meridian are 11 miles apart: find the difference in their latitudes, taking the radius of the earth as 3960 miles.

9. A man 6 ft. high stands on a tower whose height is 120 ft.: shew that at a point 24 ft. from the tower the man subtends an angle of  $31.5'$  nearly.

10. A flagstaff standing on the top of a cliff 490 feet high subtends an angle of .04 radians at a point 980 feet from the base of the cliff: find the height of the flagstaff.

11. When  $n=0$ , find the limit of

$$(1) \frac{\sin n'}{n}; \quad (2) \frac{\sin n''}{n}.$$

12. When  $n=\infty$ , find the limit of  $\frac{1}{2}nr^2 \sin \frac{2\pi}{n}$ .

When  $\theta=0$ , find the limit of

$$13. \frac{1 - \cos \theta}{\theta \sin \theta}. \quad 14. \frac{m \sin m\theta - n \sin n\theta}{\tan m\theta + \tan n\theta}.$$

15. If  $\theta=.01$  of a radian, calculate  $\cos\left(\frac{\pi}{3} + \theta\right)$ .

16. Find the value of  $\sin 30^{\circ} 10' 30''$ .

17. Given  $\cos\left(\frac{\pi}{3} + \theta\right) = .49$ , find the sexagesimal value of  $\theta$ .


### Distance and Dip of the Visible Horizon.

273. Let  $A$  be a point above the earth's surface,  $BCD$  a section of the earth by a plane passing through its centre  $E$  and  $A$ .

Let  $AE$  cut the circumference in  $B$  and  $D$ .

From  $A$  draw  $AC$  to touch the circle  $BCD$  in  $C$ , and join  $EC$ .

Draw  $AF$  at right angles to  $AD$ ; then  $\angle FAC$  is called the **dip of the horizon** as seen from  $A$ .



Thus the *dip of the horizon* is the angle of depression of any point on the horizon visible from  $A$ .

274. *To find the distance of the horizon.*

In the figure of the last article, let

$$AB=h, \quad EB=ED=r, \quad AC=x;$$

then by Euc. III. 36,  $AC^2=AB \cdot AD$ ;

that is,  $x^2=h(2r+h)=2hr+h^2$ .

For ordinary altitudes  $h^2$  is very small in comparison with  $2hr$ ; hence approximately

$$x^2=2hr \quad \text{and} \quad x=\sqrt{2hr}.$$

In this formula, suppose the measurements are made in *miles*, and let  $a$  be the number of *feet* in  $AB$ ; then

$$a=1760 \times 3 \times h.$$

By taking  $r=3960$ , we have

$$x^2=\frac{2 \times 3960 \times a}{1760 \times 3}=\frac{3a}{2}.$$

Thus we have the following rule :

*Twice the square of the distance of the horizon measured in miles is equal to three times the height of the place of observation measured in feet.*

Hence a man whose eye is 6 feet from the ground can see to a distance of 3 miles on a horizontal plane.

*Example.* The top of a ship's mast is  $66\frac{2}{3}$  ft. above the sea-level, and from it the lamp of a lighthouse can just be seen. After the ship has sailed directly towards the lighthouse for half-an-hour the lamp can be seen from the deck, which is 24 ft. above the sea. Find the rate at which the ship is sailing.

Let  $L$  denote the lamp,  $D$  and  $E$  the two positions of the ship,  $B$  the top of the mast,  $C$  the point on the deck from which the lamp is seen; then  $LCB$  is a tangent to the earth's surface at  $A$ .

[In problems like this some of the lines must necessarily be greatly out of proportion.]

Let  $AB$  and  $AC$  be expressed in miles; then since  $DB = 66\frac{2}{3}$  feet and  $EC = 24$  feet, we have by the rule

$$AB^2 = \frac{3}{2} \times 66\frac{2}{3} = 100;$$

$$\therefore AB = 10 \text{ miles.}$$

$$AC^2 = \frac{3}{2} \times 24 = 36;$$

$$\therefore AC = 6 \text{ miles.}$$

But the angles subtended by  $AB$  and  $AC$  at  $O$  the centre of the earth are very small;

$$\therefore \text{arc } AD = AB, \text{ and arc } AC = AE. \quad [\text{Art. 268.}]$$

$$\therefore \text{arc } DE = AD - AE = AB - AC = 4 \text{ miles.}$$

Thus the ship sails 4 miles in half-an-hour, or 8 miles per hour.


**275.** Let  $\theta$  be the number of radians in the dip of the horizon; then with the figure of Art. 273, we have

$$\cos \theta = \frac{EC}{EA} = \frac{r}{h+r} = \left(1 + \frac{h}{r}\right)^{-1};$$

$$\therefore 1 - 2 \sin^2 \frac{\theta}{2} = 1 - \frac{h}{r} + \frac{h^2}{r^2} - \dots;$$

$$\therefore 2 \sin^2 \frac{\theta}{2} = \frac{h}{r} - \frac{h^2}{r^2} + \dots$$

Since  $\theta$  and  $\frac{h}{r}$  are small, we may replace  $\sin \frac{\theta}{2}$  by  $\frac{\theta}{2}$  and neglect the terms on the right after the first.





Thus 
$$\frac{\theta^2}{2} = \frac{h}{r}, \quad \text{or} \quad \theta = \sqrt{\frac{2h}{r}}.$$

Let  $N$  be the number of degrees in  $\theta$  radians ; then

$$N = \frac{180\theta}{\pi} = \frac{180}{\pi} \sqrt{\frac{2h}{r}}.$$

Now  $\sqrt{r} = 63$  nearly ; hence we have approximately

$$N = \frac{180 \times 7 \times \sqrt{2h}}{22 \times 63},$$

or 
$$N = \frac{10}{11} \sqrt{2h},$$

a formula connecting the dip of the horizon in degrees and the height of the place of observation in miles.

### EXAMPLES. XXI. b.

$$\left[ \text{Here } \pi = \frac{22}{7}, \text{ and radius of earth} = 3960 \text{ miles.} \right]$$

1. Find the greatest distance at which the lamp of a lighthouse can be seen, the light being 96 feet above the sea-level.

2. If the lamp of a lighthouse begins to be seen at a distance of 15 miles, find its height above the sea-level.

3. The tops of the masts of two ships are 32 ft. 8 in. and 42 ft. 8 in. above the sea-level : find the greatest distance at which one mast can be seen from the other.

4. Find the height of a ship's mast which is just visible at a distance of 20 miles from a point on the mast of another ship which is 54 ft. above the sea-level.

5. From the mast of a ship 73 ft. 6 in. high the lamp of a lighthouse is just visible at a distance of 28 miles : find the height of the lamp.

6. Find the sexagesimal measure of the dip of the horizon from a hill 2640 feet high.

7. Along a straight coast there are lighthouses at intervals of 24 miles: find at what height the lamp must be placed so that the light of one at least may be visible at a distance of  $3\frac{1}{2}$  miles from any point of the coast.

8. From the top of a mountain the dip of the horizon is  $1^{\circ} 81'$ : find its height in feet.

9. The distance of the horizon as seen from the top of a hill is 30.25 miles: find the height of the hill and the dip of the horizon.

10. If  $x$  miles be the distance of the visible horizon and  $N$  degrees the dip, shew that

$$N = \frac{x}{66} \sqrt{\frac{10}{11}}.$$

When  $\theta=0$ , find the limit of

$$11. \frac{\sin 4\theta \cot \theta}{\operatorname{vers} 2\theta \cot^2 2\theta}. \quad 12. \frac{1 - \cos \theta + \sin \theta}{1 - \cos \theta - \sin \theta}.$$

13. When  $\theta=a$ , find the limit of

$$(1) \frac{\sin \theta - \sin a}{\theta - a}; \quad (2) \frac{\cos \theta - \cos a}{\theta - a}.$$

14. Two sides of a triangle are 31 and 32, and they include a right angle: find the other angles.

15. A person walks directly towards a distant object  $P$ , and observes that at the three points  $A, B, C$ , the elevations of  $P$  are  $a, 2a, 3a$  respectively: shew that  $AB=3BC$  nearly.


16. Shew that  $\frac{\tan \theta}{\theta}$  continually increases from 1 to  $\infty$  as  $\theta$  continually increases from 0 to  $\frac{\pi}{2}$ .

## CHAPTER XXII.

### GEOMETRICAL PROOFS.

276. *To find the expansion of  $\tan (A+B)$  geometrically.*

Let  $\angle LOM = A$ , and  $\angle MON = B$ ; then  $\angle LON = A+B$ .



In  $ON$  take any point  $P$ , and draw  $PQ$  and  $PR$  perpendicular to  $OL$  and  $OM$  respectively. Also draw  $RS$  and  $RT$  perpendicular to  $OL$  and  $PQ$  respectively.

$$\begin{aligned}\tan (A+B) &= \frac{PQ}{OQ} = \frac{RS+PT}{OS-TR} \\ &= \frac{\frac{RS}{OS} + \frac{PT}{OS}}{1 - \frac{TR}{OS}} = \frac{\frac{RS}{OS} + \frac{PT}{OS}}{1 - \frac{TR}{TP} \cdot \frac{TP}{OS}}.\end{aligned}$$

Now  $\frac{RS}{OS} = \tan A$ , and  $\frac{TR}{TP} = \tan A$ ;


also the triangles  $ROS$  and  $TPR$  are similar, and therefore

$$\frac{TP}{OS} = \frac{PR}{OR} = \tan B.$$

$$\therefore \tan (A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}.$$

In like manner, with the help of the figure on page 95, we may obtain the expansion of  $\tan (A - B)$  geometrically.

**277.** *To prove geometrically the formulæ for transformation of sums into products.*



Let  $\angle EOF$  be denoted by  $A$ , and  $\angle EOG$  by  $B$ .

With centre  $O$  and any radius describe an arc of a circle meeting  $OG$  in  $H$  and  $OF$  in  $K$ .

Bisect  $\angle KOH$  by  $OL$ ; then  $OL$  bisects  $HK$  at right angles.

Draw  $KP$ ,  $HQ$ ,  $LR$  perpendicular to  $OE$ , and through  $L$  draw  $MLN$  parallel to  $OE$  meeting  $KP$  in  $M$  and  $HQ$  in  $N$ .

It is easy to prove that the triangles  $MKL$  and  $NHL$  are equal in all respects, so that  $KM = NH$ ,  $ML = LN$ ,  $PR = RQ$ .

Also  $\angle GOF = A - B$ , and therefore

$$\angle HOL = \angle KOL = \frac{A - B}{2};$$

$$\therefore \angle EOL = B + \frac{A - B}{2} = \frac{A + B}{2}.$$

$$\begin{aligned} \sin A + \sin B &= \frac{KP}{OK} + \frac{HQ}{OH} = \frac{KP + HQ}{OK} \\ &= \frac{(KM + LR) + (LR - NH)}{OK} = 2 \frac{LR}{OK}; \end{aligned}$$

$$\therefore \sin A + \sin B = 2 \frac{LR}{OL} \cdot \frac{OL}{OK} = 2 \sin ROL \cos KOL$$

$$= 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}.$$

$$\cos A + \cos B = \frac{OP}{OK} + \frac{OQ}{OH} = \frac{OP+OQ}{OK}$$

$$= \frac{(OR-PR)+(OR+RQ)}{OK} = 2 \frac{OR}{OK}$$

$$= 2 \frac{OR}{OL} \cdot \frac{OL}{OK} = 2 \cos ROL \cos KOL$$

$$= 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}.$$

$$\sin A - \sin B = \frac{KP}{OK} - \frac{HQ}{OH} = \frac{KP-HQ}{OK}$$

$$= \frac{(KM+LR)-(LR-NH)}{OK} = 2 \frac{KM}{OK}$$

$$= 2 \frac{KM}{KL} \cdot \frac{KL}{OK} = 2 \cos LKM \sin KOL$$

$$= 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2},$$

$$\text{since } \angle LKM = \text{comp}^t \text{ of } \angle KLM = \angle MLO = \angle LOE = \frac{A+B}{2}.$$

$$\cos B - \cos A = \frac{OQ}{OH} - \frac{OP}{OK} = \frac{OQ-OP}{OK}$$

$$= \frac{(OR+RQ)-(OR-PR)}{OK} = 2 \frac{PR}{OK} = 2 \frac{ML}{OK}$$

$$= 2 \frac{ML}{KL} \cdot \frac{KL}{OK} = 2 \sin LKM \sin KOL$$

$$= 2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}.$$

**278.** *Geometrical proof of the  $2A$  formulæ.*

Let  $BPD$  be a semicircle,  $BD$  the diameter,  $C$  the centre.


On the circumference, take any point  $P$ , and join  $PB$ ,  $PC$ ,  $PD$ .

Draw  $PN$  perpendicular to  $BD$ .

Let  $\angle PBD = A$ , then

$$\angle PCD = 2A.$$

And  $\angle NPD = \text{comp}^t$  of  $\angle PDN = \angle PBD = A$ .



$$\begin{aligned}\sin 2A &= \frac{PN}{CP} = \frac{2PN}{2CP} = \frac{2PN}{BD} = 2 \frac{PN}{BP} \cdot \frac{BP}{BD} \\ &= 2 \sin PBN \cos PBD \\ &= 2 \sin A \cos A.\end{aligned}$$

$$\begin{aligned}\cos 2A &= \frac{CN}{CP} = \frac{2CN}{BD} = \frac{CN + CN}{BD} \\ &= \frac{(BN - BC) + (CD - ND)}{BD} = \frac{BN - ND}{BD} \\ &= \frac{BN}{BP} \cdot \frac{BP}{BD} - \frac{ND}{PD} \cdot \frac{PD}{BD} \\ &= \cos A \cdot \cos A - \sin A \cdot \sin A \\ &= \cos^2 A - \sin^2 A.\end{aligned}$$

$$\begin{aligned}\cos 2A &= \frac{CN}{CP} = \frac{CD - DN}{CP} = 1 - \frac{DN}{CP} = 1 - \frac{2DN}{BD} \\ &= 1 - 2 \frac{DN}{DP} \cdot \frac{DP}{BD} = 1 - 2 \sin A \cdot \sin A \\ &= 1 - 2 \sin^2 A.\end{aligned}$$

$$\begin{aligned}\cos 2A &= \frac{CN}{CP} = \frac{BN - BC}{CP} = \frac{BN}{CP} - 1 = \frac{2BN}{BD} - 1 \\ &= 2 \frac{BN}{BP} \cdot \frac{BP}{BD} - 1 = 2 \cos A \cdot \cos A - 1 \\ &= 2 \cos^2 A - 1.\end{aligned}$$



$$\begin{aligned}
 \tan 2A &= \frac{PN}{CN} = \frac{2PN}{2CN} = \frac{2PN}{BN - ND} \\
 &= \frac{2 \frac{PN}{BN}}{1 - \frac{ND}{BN}} = \frac{2 \frac{PN}{BN}}{1 - \frac{ND}{PN} \cdot \frac{PN}{BN}} \\
 &= \frac{2 \tan A}{1 - \tan A \cdot \tan A} \\
 &= \frac{2 \tan A}{1 - \tan^2 A}.
 \end{aligned}$$

279. To find the value of  $\sin 18^\circ$  geometrically.

Let  $ABD$  be an isosceles triangle in which each angle at the base  $BD$  is double the vertical angle  $A$ ; then

$$A + 2A + 2A = 180^\circ,$$

and therefore  $A = 36^\circ$ .

Bisect  $\angle BAD$  by  $AE$ ; then  $AE$  bisects  $BD$  at right angles;

$$\therefore \angle BAE = 18^\circ.$$

Thus  $\sin 18^\circ = \frac{BE}{AB} = \frac{x}{a},$

where  $AB = a$ , and  $BE = x$ .

From the construction given in Euc. iv. 10,

$$AC = BD = 2BE = 2x,$$

and

$$AB \cdot BC = AC^2;$$


$$\therefore a(a - 2x) = (2x)^2;$$

$$\therefore 4x^2 + 2ax - a^2 = 0;$$

$$\therefore x = \frac{-2a \pm \sqrt{20a^2}}{8} = \frac{-1 \pm \sqrt{5}}{4} a.$$


The upper sign must be taken, since  $x$  is positive. Thus

$$\sin 18^\circ = \frac{\sqrt{5} - 1}{4}.$$



### Proofs by Projection.

**280. DEFINITION.** If from any two points  $A$  and  $B$ , lines  $AC$  and  $BD$  are drawn perpendicular to  $OX$ , then the intercept  $CD$  is called the **projection** of  $AB$  upon  $OX$ .



Through  $A$  draw  $AE$  parallel to  $OX$ ; then

$$CD = AE = AB \cos BAE;$$

that is,

$$CD = AB \cos a,$$

where  $a$  is the angle of inclination of the lines  $AB$  and  $OX$ .

**281.** *To shew that the projection of a straight line is equal to the projection of an equal and parallel straight line drawn from a fixed point.*

Let  $AB$  be any straight line,  $O$  a fixed point, which we shall call the origin,  $OP$  a straight line equal and parallel to  $AB$ .


Let  $CD$  and  $OM$  be the projections of  $AB$  and  $OP$  upon any straight line  $OX$  drawn through the origin.

Draw  $AE$  parallel to  $OX$ .

The two triangles  $AEB$  and  $OMP$  are identically equal;

$$\therefore OM = AE = CD;$$

that is, projection of  $OP$  = projection of  $AB$ .



**282.** In the figure of the last article, two straight lines  $OP$  and  $OQ$  can be drawn from  $O$  equal and parallel to  $AB$ ; it is therefore necessary to have some means of fixing the *direction* in which the line from  $O$  is to be drawn. Accordingly it is agreed to consider that

*the direction of a line is fixed by the order of the letters.*

Thus  $AB$  denotes a line drawn from  $A$  to  $B$ , and  $BA$  denotes a line drawn from  $B$  to  $A$ .

Hence  $OP$  denotes a line drawn from the origin parallel to  $AB$ , and  $OQ$  denotes a line drawn from the origin parallel to  $BA$ .

Similarly the direction of a projected line is fixed by the order of the letters.

Thus  $CD$  is drawn to the right from  $C$  to  $D$  and is positive, while  $DC$  is drawn to the left from  $D$  to  $C$  and is negative.

Hence in sign as well as in magnitude

$$OM = CD, \text{ and } ON = DC;$$

that is, projection of  $OP$  = projection of  $AB$ ,

and projection of  $OQ$  = projection of  $BA$ .

Thus *the projection of a straight line can be represented both in sign and magnitude by the projection of an equal and parallel straight line drawn from the origin.*

**283.** Whatever be the direction of  $AB$ , the line  $OP$  will fall within one of the four quadrants.


Also from the definitions given in Art. 75, we have

$$\frac{OM}{OP} = \cos XOP,$$

that is,

$$OM = OP \cos XOP,$$

whatever be the magnitude of the angle  $XOP$ . We shall always suppose, unless the contrary is stated, that the angles are measured in the positive direction.




**284.** Let  $O$  be the origin,  $P$  and  $Q$  any two points.

Join  $OP$ ,  $OQ$ ,  $PQ$ , and draw  $PM$  and  $QN$  perpendicular to  $OX$ .

We have

$$OM = ON + NM,$$

since the line  $NM$  is to be regarded as negative; that is,



the projection of  $OP$  = projection of  $OQ$  + projection of  $QP$ .



Hence, *the projection of one side of a triangle is equal to the sum of the projections of the other two sides taken in order.*  
Thus

projection of  $OQ$  = projection of  $OP$  + projection of  $PQ$ ;

projection of  $QP$  = projection of  $QO$  + projection of  $OP$ .

### General Proof of the Addition Formulæ.

285. In Fig. 1, let a line starting from  $OX$  revolve until it has traced the angle  $A$ , taking up the position  $OM$ , and then let it further revolve until it has traced the angle  $B$ , taking up the final position  $ON$ . Thus  $XON$  is the angle  $A+B$ .



In  $ON$  take any point  $P$ , and draw  $PQ$  perpendicular to  $OM$ ; also draw  $OR$  equal and parallel to  $QP$ .

*Projecting upon  $OX$* , we have

projection of  $OP$  = projection of  $OQ$  + projection of  $QP$   
 = projection of  $OQ$  + projection of  $OR$ .

$$\therefore OP \cos XOP = OQ \cos XOQ + OR \cos XOR \dots\dots\dots(1)$$

$$= OP \cos B \cos XOQ + OP \sin B \cos XOR;$$

$$\therefore \cos XOP = \cos B \cos XOQ + \sin B \cos XOR;$$

that is,  $\cos (A+B) = \cos B \cos A + \sin B \cos (90^\circ + A)$   
 $= \cos A \cos B - \sin A \sin B.$

*Projecting upon  $OY$* , we have only to write  $Y$  for  $X$  in (1);

$$\begin{aligned}
 \text{thus } OP \cos YOP &= OQ \cos YOQ + OR \cos YOR \\
 &= OP \cos B \cos YOQ + OP \sin B \cos YOR; \\
 \therefore \cos YOP &= \cos B \cos YOQ + \sin B \cos YOR;
 \end{aligned}$$

that is,

$$\begin{aligned}
 \cos (A + B - 90^\circ) &= \cos B \cos (A - 90^\circ) + \sin B \cos A; \\
 \therefore \sin (A + B) &= \sin A \cos B + \cos A \sin B.
 \end{aligned}$$

In Fig. 2, let a line starting from  $OX$  revolve until it has traced the angle  $A$ , taking up the position  $OM$ , and then let it revolve *back again* until it has traced the angle  $B$ , taking up the final position  $ON$ . Thus  $XON$  is the angle  $A - B$ .

In  $ON$  take any point  $P$ , and draw  $PQ$  perpendicular to  $MO$  produced; also draw  $OR$  equal and parallel to  $QP$ .

*Projecting upon  $OX$* , we have as in the previous case

$$\begin{aligned}
 OP \cos XOP &= OQ \cos XOQ + OR \cos XOR \\
 &= OP \cos (180^\circ - B) \cos XOQ \\
 &\quad + OP \sin (180^\circ - B) \cos XOR; \\
 \therefore \cos XOP &= -\cos B \cos XOQ + \sin B \cos XOR;
 \end{aligned}$$

that is,

$$\begin{aligned}
 \cos (A - B) &= -\cos B \cos (A - 180^\circ) + \sin B \cos (A - 90^\circ) \\
 &= -\cos B (-\cos A) + \sin B \sin A \\
 &= \cos A \cos B + \sin A \sin B.
 \end{aligned}$$

*Projecting upon  $OY$* , we have

$$\begin{aligned}
 OP \cos YOP &= OQ \cos YOQ + OR \cos YOR; \\
 &= OP \cos (180^\circ - B) \cos YOQ \\
 &\quad + OP \sin (180^\circ - B) \cos YOR; \\
 \therefore \cos YOP &= -\cos B \cos YOQ + \sin B \cos YOR;
 \end{aligned}$$

that is,

$$\begin{aligned}
 \cos (A - B - 90^\circ) &= -\cos B \cos (A - 270^\circ) + \sin B \cos (A - 180^\circ); \\
 \therefore \sin (A - B) &= -\cos B (-\sin A) + \sin B (-\cos A) \\
 &= \sin A \cos B - \cos A \sin B.
 \end{aligned}$$

236. The above method of proof is applicable to every case, and therefore the Addition Formulæ are universally established.

The universal truth of the Addition Formulæ may also be deduced from the special geometrical investigations of Arts. 110 and 111 by analysis, as in the next article.

287. When each of the angles  $A$ ,  $B$ ,  $A+B$  is less than  $90^\circ$ , we have shewn that

$$\cos(A+B) = \cos A \cos B - \sin A \sin B \dots\dots\dots(1).$$

$$\text{But } \cos(A+B) = \sin(\overline{A+B} + 90^\circ) = \sin(\overline{A+90^\circ} + B);$$

$$\text{also} \qquad \qquad \qquad \cos A = \sin(A+90^\circ),$$

$$\text{and} \qquad \qquad \qquad -\sin A = \cos(A+90^\circ). \quad [\text{Art. 98.}]$$

Hence by substitution in (1), we have

$$\sin(\overline{A+90^\circ} + B) = \sin(A+90^\circ) \cos B + \cos(A+90^\circ) \sin B.$$

In like manner, it may be proved that

$$\cos(\overline{A+90^\circ} + B) = \cos(A+90^\circ) \cos B - \sin(A+90^\circ) \sin B.$$

Thus the formulæ for the sine and cosine of  $A+B$  hold when  $A$  is increased by  $90^\circ$ . Similarly we may shew that they hold when  $B$  is increased by  $90^\circ$ .

By repeated applications of the same process it may be proved that the formulæ are true when either or both of the angles  $A$  and  $B$  is increased by any multiple of  $90^\circ$ .

$$\text{Again, } \cos(A+B) = \cos A \cos B - \sin A \sin B \dots\dots\dots(1).$$

$$\text{But } \cos(A+B) = -\sin(\overline{A+B} - 90^\circ) = -\sin(\overline{A-90^\circ} + B);$$

$$\text{also} \qquad \qquad \qquad \cos A = -\sin(A-90^\circ),$$

$$\text{and} \qquad \qquad \qquad \sin A = \cos(A-90^\circ). \quad [\text{Arts. 99 and 102.}]$$

Hence by substitution in (1), we have

$$\sin(\overline{A-90^\circ} + B) = \sin(A-90^\circ) \cos B + \cos(A-90^\circ) \sin B.$$

Similarly we may shew that

$$\cos(\overline{A-90^\circ} + B) = \cos(A-90^\circ) \cos B - \sin(A-90^\circ) \sin B.$$

Thus the formulæ for the sine and cosine of  $A+B$  hold when  $A$  is diminished by  $90^\circ$ . In like manner we may prove that they are true when  $B$  is diminished by  $90^\circ$ .



By repeated applications of the same process it may be shewn that the formulæ hold when either or both of the angles  $A$  and  $B$  is diminished by any multiple of  $90^\circ$ . Further, it will be seen that the formulæ are true if either of the angles  $A$  or  $B$  is increased by a multiple of  $90^\circ$  and the other is diminished by a multiple of  $90^\circ$ .

Thus  $\sin(P+Q) = \sin P \cos Q + \cos P \sin Q$ ,  
 and  $\cos(P+Q) = \cos P \cos Q - \sin P \sin Q$ ,  
 where  $P = A \pm m \cdot 90^\circ$ , and  $Q = B \pm n \cdot 90^\circ$ ,  
 $m$  and  $n$  being any positive integers, and  $A$  and  $B$  any acute angles.

Thus the Addition Formulæ are true for the algebraical sum of any two angles.

### MISCELLANEOUS EXAMPLES. H.

1. If the sides of a right-angled triangle are  $\cos 2a + \cos 2\beta + 2 \cos(a+\beta)$  and  $\sin 2a + \sin 2\beta + \sin 2(a+\beta)$ , shew that the hypotenuse is  $4 \cos^2 \frac{a-\beta}{2}$ .
2. If the in-centre and circum-centre be at equal distances from  $BC$ , prove that  $\cos B + \cos C = 1$ .
3. The shadow of a tower is observed to be half the known height of the tower, and some time afterwards to be equal to the height: how much will the sun have gone down in the interval? Given  $\log 2$ ,  
 $L \tan 63^\circ 26' = 10 \cdot 3009994$ , diff. for  $1' = 3159$ .
4. If  $(1 + \sin a)(1 + \sin \beta)(1 + \sin \gamma)$   
 $= (1 - \sin a)(1 - \sin \beta)(1 - \sin \gamma)$ ,  
 shew that each expression is equal to  $\pm \cos a \cos \beta \cos \gamma$ .
5. Two parallel chords of a circle lying on the same side of the centre subtend  $72^\circ$  and  $144^\circ$  at the centre: prove that the distance between them is one-half of the radius.  
 Also shew that the sum of the squares of the chords is equal to five times the square of the radius.

6. Two straight railways are inclined at an angle of  $60^\circ$ . From their point of intersection two trains  $A$  and  $B$  start at the same time, one along each line.  $A$  travels at the rate of 48 miles per hour, at what rate must  $B$  travel so that after one hour they shall be 43 miles apart?

7. If 
$$a = \cos^{-1} \frac{x}{a} + \cos^{-1} \frac{y}{b},$$

shew that 
$$\sin^2 a = \frac{x^2}{a^2} - \frac{2xy}{ab} \cos a + \frac{y^2}{b^2}.$$

8. If  $p, q, r$  denote the sides of the ex-central triangle, prove that

$$\frac{a^2}{p^2} + \frac{b^2}{q^2} + \frac{c^2}{r^2} + \frac{2abc}{pqr} = 1.$$

9. A tower is situated within the angle formed by two straight roads  $OA$  and  $OB$ , and subtends angles  $\alpha$  and  $\beta$  at the points  $A$  and  $B$  where the roads are nearest to it. If  $OA = a$ , and  $OB = b$ , shew that the height of the tower is

$$\sqrt{a^2 - b^2} \sin \alpha \sin \beta / \sqrt{\sin (\alpha + \beta) \sin (\alpha - \beta)}.$$

10. In a triangle, shew that

$$r^2 + r_1^2 + r_2^2 + r_3^2 = 16R^2 - a^2 - b^2 - c^2.$$

11. If  $AD$  be a median of the triangle  $ABC$ , shew that

$$(1) \quad \cot BAD = 2 \cot A + \cot B;$$

$$(2) \quad 2 \cot ADC = \cot B - \cot C.$$

12. If  $p, q, r$  are the distances of the orthocentre from the sides, prove that

$$4 \left( \frac{a}{p} + \frac{b}{q} + \frac{c}{r} \right) = \left( \frac{a}{p} + \frac{b}{q} - \frac{c}{r} \right) \left( \frac{b}{q} + \frac{c}{r} - \frac{a}{p} \right) \left( \frac{c}{r} + \frac{a}{p} - \frac{b}{q} \right).$$


## Graphical Representation of the Circular Functions.

288. DEFINITION. Let  $f(x)$  be a function of  $x$  which has a single value for all values of  $x$ , and let the values of  $x$  be represented by lines measured from  $O$  along  $OX$  or  $OX'$ , and the values of  $f(x)$  by lines drawn perpendicular to  $XX'$ . Then with the figure of the next article, if  $OM$  represent any value of  $x$ , and  $MP$  the corresponding value of  $f(x)$ , the curve traced out by the point  $P$  is called the **Graph** of  $f(x)$ .

### Graphs of $\sin \theta$ and $\cos \theta$ .

289. Suppose that the unit of length is chosen to represent a radian; then any angle of  $\theta$  radians will be represented by a line  $OM$  which contains  $\theta$  units of length.

Graph of  $\sin \theta$ .



Let  $MP$ , drawn perpendicular to  $OX$ , represent the value of  $\sin \theta$  corresponding to the value  $OM$  of  $\theta$ ; then the curve traced out by the point  $P$  represents the graph of  $\sin \theta$ .

As  $OM$  or  $\theta$  increases from 0 to  $\frac{\pi}{2}$ ,  $MP$  or  $\sin \theta$  increases from 0 to 1, which is its greatest value.

As  $OM$  increases from  $\frac{\pi}{2}$  to  $\pi$ ,  $MP$  decreases from 0 to 1.

As  $OM$  increases from  $\pi$  to  $\frac{3\pi}{2}$ ,  $MP$  increases numerically from 0 to  $-1$ .

As  $OM$  increases from  $\frac{3\pi}{2}$  to  $2\pi$ ,  $MP$  decreases numerically from  $-1$  to 0.

As  $OM$  increases from  $2\pi$  to  $4\pi$ , from  $4\pi$  to  $6\pi$ , from  $6\pi$  to  $8\pi$ , .....,  $MP$  passes through the same series of values as when  $OM$  increases from 0 to  $2\pi$ .

Since  $\sin(-\theta) = -\sin \theta$ , the values of  $MP$  lying to the left of  $O$  are equal in magnitude but are of opposite sign to values of  $MP$  lying at an equal distance to the right of  $O$ .

Thus the graph of  $\sin \theta$  is a *continuous* waving line extending to an infinite distance on each side of  $O$ .

The graph of  $\cos \theta$  is the same as that of  $\sin \theta$ , the origin being at the point marked  $\frac{\pi}{2}$  in the figure.

### Graphs of $\tan \theta$ and $\cot \theta$ .

290. As before, suppose that the unit of length is chosen to represent a radian; then any angle of  $\theta$  radians will be represented by a line  $OM$  which contains  $\theta$  units of length.

Let  $MP$ , drawn perpendicular to  $OX$ , represent the value of  $\tan \theta$  corresponding to the value  $OM$  of  $\theta$ ; then the curve traced out by the point  $P$  represents the graph of  $\tan \theta$ .


By tracing the changes in the value of  $\tan \theta$  as  $\theta$  varies from 0 to  $2\pi$ , from  $2\pi$  to  $4\pi$ , ....., it will be seen that the graph of  $\tan \theta$  consists of an infinite number of *discontinuous* equal branches as represented in the figure below. The part of each branch beneath  $XX'$  is convex towards  $XX'$ , and the part of each branch above  $XX'$  is also convex towards  $XX'$ ; hence at the point where any branch cuts  $XX'$  there is what is called a *point of inflexion*, where the direction of curvature changes. The proof of these statements is however beyond the range of the present work.

The various branches touch the dotted lines passing through the points marked

$$\pm \frac{\pi}{2}, \quad \pm \frac{3\pi}{2}, \quad \pm \frac{5}{2}, \quad \dots\dots,$$

at an infinite distance from  $XX'$ .

Graph of  $\tan \theta$ .



The student should draw the graph of  $\cot \theta$ , which is very similar to that of  $\tan \theta$ .


### Graphs of $\sec \theta$ and $\operatorname{cosec} \theta$ .

291. The graph of  $\sec \theta$  is represented in the figure below. It consists of an infinite number of equal festoons lying alternately above and below  $XX'$ , the vertex of each being at the unit of distance from  $XX'$ . The various festoons touch the dotted lines passing through the points marked

$$\pm \frac{\pi}{2}, \quad \pm \frac{3\pi}{2}, \quad \pm \frac{5\pi}{2}, \dots,$$

at an infinite distance from  $XX'$ .

Graph of  $\sec \theta$ .



The graph of  $\operatorname{cosec} \theta$  is the same as that of  $\sec \theta$ , the origin being at the point marked  $-\frac{\pi}{2}$  in the figure.

## CHAPTER XXIII.

### SUMMATION OF FINITE SERIES.

292. An expression in which the successive terms are formed by some regular law is called a **series**. If the series ends at some assigned term it is called a **finite series**; if the number of terms is unlimited it is called an **infinite series**.

A series may be denoted by an expression of the form

$$u_1 + u_2 + u_3 + \dots + u_{n-1} + u_n + u_{n+1} + \dots,$$

where  $u_{n+1}$ , the  $(n+1)^{\text{th}}$  term, is obtained from  $u_n$ , the  $n^{\text{th}}$  term, by replacing  $n$  by  $n+1$ .

Thus if  $u_n = \cos (a + n\beta)$ , then  $u_{n+1} = \cos \{a + (n+1)\beta\}$ ;  
and if  $u_n = \cot 2^{n-1}a$ , then  $u_{n+1} = \cot 2^n a$ .

293. If the  $r^{\text{th}}$  term of a series can be expressed as the difference of two quantities one of which is the same function of  $r$  that the other is of  $r+1$ , the sum of the series may be readily found.

For let the series be denoted by

$$u_1 + u_2 + u_3 + \dots + u_n,$$

and its sum by  $S$ , and suppose that any term

$$u_r = v_{r+1} - v_r;$$

then  $S = (v_2 - v_1) + (v_3 - v_2) + (v_4 - v_3) + \dots + (v_n - v_{n-1}) + (v_{n+1} - v_n)$   
 $= v_{n+1} - v_1.$

*Example.* Find the sum of the series

$$\operatorname{cosec} a + \operatorname{cosec} 2a + \operatorname{cosec} 4a + \dots + \operatorname{cosec} 2^{n-1}a.$$

$$\operatorname{cosec} a = \frac{1}{\sin a} = \frac{\sin \frac{a}{2}}{\sin \frac{a}{2} \sin a} = \frac{\sin \left( a - \frac{a}{2} \right)}{\sin \frac{a}{2} \sin a}.$$



Hence  $\operatorname{cosec} a = \cot \frac{a}{2} - \cot a.$

If we replace  $a$  by  $2a$ , we obtain

$$\operatorname{cosec} 2a = \cot a - \cot 2a.$$

Similarly,  $\operatorname{cosec} 4a = \cot 2a - \cot 4a,$

.....

$$\operatorname{cosec} 2^{n-1} a = \cot 2^{n-2} a - \cot 2^{n-1} a.$$

By addition,  $S = \cot \frac{a}{2} - \cot 2^{n-1} a.$

**294.** *To find the sum of the sines of a series of  $n$  angles which are in arithmetical progression.*

Let the sine-series be denoted by

$$\sin a + \sin (a + \beta) + \sin (a + 2\beta) + \dots + \sin \{a + (n-1)\beta\}.$$

We have the identities

$$2 \sin a \sin \frac{\beta}{2} = \cos \left( a - \frac{\beta}{2} \right) - \cos \left( a + \frac{\beta}{2} \right),$$

$$2 \sin (a + \beta) \sin \frac{\beta}{2} = \cos \left( a + \frac{\beta}{2} \right) - \cos \left( a + \frac{3\beta}{2} \right),$$

$$2 \sin (a + 2\beta) \sin \frac{\beta}{2} = \cos \left( a + \frac{3\beta}{2} \right) - \cos \left( a + \frac{5\beta}{2} \right),$$

.....

$$2 \sin \{a + (n-1)\beta\} \sin \frac{\beta}{2} = \cos \left( a + \frac{2n-3}{2} \beta \right) - \cos \left( a + \frac{2n-1}{2} \beta \right).$$

By addition,

$$2S \sin \frac{\beta}{2} = \cos \left( a - \frac{\beta}{2} \right) - \cos \left( a + \frac{2n-1}{2} \beta \right)$$

$$= 2 \sin \left( a + \frac{n-1}{2} \beta \right) \sin \frac{n\beta}{2};$$

$$\therefore S = \frac{\sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}} \sin \left( a + \frac{n-1}{2} \beta \right).$$

295. In like manner we may shew that the sum of the cosine-series

$$\begin{aligned} \cos \alpha + \cos (\alpha + \beta) + \cos (\alpha + 2\beta) + \dots + \cos \{\alpha + (n-1)\beta\} \\ = \frac{\sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}} \cos \left( \alpha + \frac{n-1}{2} \beta \right). \end{aligned}$$

296. The formulæ of the two last articles may be expressed verbally as follows.

*The sum of the sines of a series of n angles in A.P.*

$$= \frac{\sin \frac{n \text{ diff.}}{2}}{\sin \frac{\text{diff.}}{2}} \sin \frac{\text{first angle} + \text{last angle}}{2}.$$

*The sum of the cosines of a series of n angles in A.P.*

$$= \frac{\sin \frac{n \text{ diff.}}{2}}{\sin \frac{\text{diff.}}{2}} \cos \frac{\text{first angle} + \text{last angle}}{2}.$$

*Example.* Find the sum of the series

$$\cos \alpha + \cos 3\alpha + \cos 5\alpha + \dots + \cos (2n-1)\alpha.$$

Here the common difference of the angles is  $2\alpha$ ;

$$\begin{aligned} \therefore S &= \frac{\sin n\alpha}{\sin \alpha} \cos \frac{\alpha + (2n-1)\alpha}{2} \\ &= \frac{\sin n\alpha \cos n\alpha}{\sin \alpha} = \frac{\sin 2n\alpha}{2 \sin \alpha}. \end{aligned}$$

297. If  $\sin \frac{n\beta}{2} = 0$ , each of the expressions found in Arts. 294 and 295 for the sum vanishes. In this case

$$\frac{n\beta}{2} = k\pi, \text{ or } \beta = \frac{2k\pi}{n}, \text{ where } k \text{ is any integer.}$$

Hence the sum of the sines and the sum of the cosines of  $n$  angles in arithmetical progression are each equal to zero, when the common difference of the angles is an even multiple of  $\frac{\pi}{n}$ .

**298.** Some series may be brought under the rule of Art. 296 by a simple transformation.

*Example 1.* Find the sum of  $n$  terms of the series

$$\cos \alpha - \cos (\alpha + \beta) + \cos (\alpha + 2\beta) - \cos (\alpha + 3\beta) + \dots$$

This series is equal to

$\cos \alpha + \cos (\alpha + \beta + \pi) + \cos (\alpha + 2\beta + 2\pi) + \cos (\alpha + 3\beta + 3\pi) + \dots$ ,  
a series in which the common difference of the angles is  $\beta + \pi$ , and the last angle is  $\alpha + (n-1)(\beta + \pi)$ .

$$\therefore S = \frac{\sin \frac{n(\beta + \pi)}{2}}{\sin \frac{\beta + \pi}{2}} \cos \left\{ \alpha + \frac{(n-1)(\beta + \pi)}{2} \right\}.$$

*Example 2.* Find the sum of  $n$  terms of the series

$$\sin \alpha + \cos (\alpha + \beta) - \sin (\alpha + 2\beta) - \cos (\alpha + 3\beta) + \sin (\alpha + 4\beta) + \dots$$

This series is equal to

$$\sin \alpha + \sin \left( \alpha + \beta + \frac{\pi}{2} \right) + \sin (\alpha + 2\beta + \pi) + \sin \left( \alpha + 3\beta + \frac{3\pi}{2} \right) + \dots,$$

a series in which the common difference of the angles is  $\beta + \frac{\pi}{2}$ .

$$\therefore S = \frac{\sin \frac{n(2\beta + \pi)}{4}}{\sin \frac{2\beta + \pi}{4}} \sin \left\{ \alpha + \frac{(n-1)(2\beta + \pi)}{4} \right\}.$$

### EXAMPLES. XXIII. a.

Sum each of the following series to  $n$  terms :

1.  $\sin \alpha + \sin 3\alpha + \sin 5\alpha + \dots$

2.  $\cos \alpha + \cos (\alpha - \beta) + \cos (\alpha - 2\beta) + \dots$

3.  $\sin \alpha + \sin \left( \alpha - \frac{\pi}{n} \right) + \sin \left( \alpha - \frac{2\pi}{n} \right) + \dots$

4.  $\cos \frac{\pi}{k} + \cos \frac{2\pi}{k} + \cos \frac{3\pi}{k} + \dots$

Find the sum of each of the following series :

$$5. \cos \frac{\pi}{19} + \cos \frac{3\pi}{19} + \cos \frac{5\pi}{19} + \dots + \cos \frac{17\pi}{19}.$$

$$6. \cos \frac{2\pi}{21} + \cos \frac{4\pi}{21} + \cos \frac{6\pi}{21} + \dots + \cos \frac{20\pi}{21}.$$

$$7. \sin \frac{\pi}{n} + \sin \frac{2\pi}{n} + \sin \frac{3\pi}{n} + \dots \text{ to } n-1 \text{ terms.}$$

$$8. \cos \frac{\pi}{n} + \cos \frac{3\pi}{n} + \cos \frac{5\pi}{n} + \dots \text{ to } 2n-1 \text{ terms.}$$

$$9. \sin na + \sin (n-1)a + \sin (n-2)a + \dots \text{ to } 2n \text{ terms.}$$

Sum each of the following series to  $n$  terms :

$$10. \sin \theta - \sin 2\theta + \sin 3\theta - \sin 4\theta + \dots.$$

$$11. \cos a - \cos (a - \beta) + \cos (a - 2\beta) - \cos (a - 3\beta) + \dots.$$

$$12. \cos a - \sin (a - \beta) - \cos (a - 2\beta) + \sin (a - 3\beta) + \dots.$$

$$13. \sin 2\theta \sin \theta + \sin 3\theta \sin 2\theta + \sin 4\theta \sin 3\theta + \dots.$$

$$14. \sin a \cos 3a + \sin 3a \cos 5a + \sin 5a \cos 7a + \dots.$$

$$15. \sec a \sec 2a + \sec 2a \sec 3a + \sec 3a \sec 4a + \dots.$$

$$16. \operatorname{cosec} \theta \operatorname{cosec} 3\theta + \operatorname{cosec} 3\theta \operatorname{cosec} 5\theta \\ + \operatorname{cosec} 5\theta \operatorname{cosec} 7\theta + \dots.$$

$$17. \tan \frac{a}{2} \sec a + \tan \frac{a}{2^2} \sec \frac{a}{2} + \tan \frac{a}{2^3} \sec \frac{a}{2^2} + \dots.$$

$$18. \cos 2a \operatorname{cosec} 3a + \cos 6a \operatorname{cosec} 9a + \cos 18a \operatorname{cosec} 27a + \dots.$$

$$19. \sin a \sec 3a + \sin 3a \sec 9a + \sin 9a \sec 27a + \dots.$$

20. The circumference of a semicircle of radius  $a$  is divided into  $n$  equal arcs. Shew that the sum of the distances of the several points of section from either extremity of the diameter is

$$a \left( \cot \frac{\pi}{4n} - 1 \right).$$

21. From the angular points of a regular polygon, perpendiculars are drawn to  $XX'$  and  $YY'$  the horizontal and vertical diameter of the circumscribing circle: shew that the algebraical sums of each of the two sets of perpendiculars are equal to zero.

299. By means of the identities

$$\begin{aligned} 2 \sin^2 a &= 1 - \cos 2a, & 2 \cos^2 a &= 1 + \cos 2a, \\ 4 \sin^3 a &= 3 \sin a - \sin 3a, & 4 \cos^3 a &= 3 \cos a + \cos 3a, \end{aligned}$$

we can find the sum of the squares and cubes of the sines and cosines of a series of angles in arithmetical progression.

*Example 1.* Find the sum of  $n$  terms of the series

$$\sin^2 a + \sin^2(a + \beta) + \sin^2(a + 2\beta) + \dots$$

$$2S = \{1 - \cos 2a\} + \{1 - \cos(2a + 2\beta)\} + \{1 - \cos(2a + 4\beta)\} + \dots$$

$$= n - \{\cos 2a + \cos(2a + 2\beta) + \cos(2a + 4\beta) + \dots\};$$

$$= n - \frac{\sin n\beta}{\sin \beta} \cos \frac{2a + \{2a + (n-1)2\beta\}}{2};$$

$$\therefore S = \frac{n}{2} - \frac{\sin n\beta}{2 \sin \beta} \cos \{2a + (n-1)\beta\}.$$

*Example 2.* Find the sum of the series

$$\cos^3 a + \cos^3 3a + \cos^3 5a + \dots + \cos^3(2n-1)a.$$

$$4S = (3 \cos a + \cos 3a) + (3 \cos 3a + \cos 9a) + (3 \cos 5a + \cos 15a) + \dots$$

$$= 3(\cos a + \cos 3a + \cos 5a + \dots) + (\cos 3a + \cos 9a + \cos 15a + \dots)$$

$$= \frac{3 \sin na}{\sin a} \cos \left\{ \frac{a + (2n-1)a}{2} \right\} + \frac{\sin 3na}{\sin 3a} \cos \left\{ \frac{3a + (2n-1)3a}{2} \right\};$$

$$\therefore S = \frac{3 \sin na \cos na}{4 \sin a} + \frac{\sin 3na \cos 3na}{4 \sin 3a}.$$

300. The following further examples illustrate the principle of Art. 293.

*Example 1.* Find the sum of the series

$$\tan^{-1} \frac{x}{1+1 \cdot 2 \cdot x^2} + \tan^{-1} \frac{x}{1+2 \cdot 3 \cdot x^2} + \dots + \tan^{-1} \frac{x}{1+n(n+1)x^2}.$$

As in Art. 249, we have

$$\tan^{-1} \frac{x}{1+r(r+1)x^2} = \tan^{-1}(r+1)x - \tan^{-1}rx;$$

$$\therefore S = \tan^{-1}(n+1)x - \tan^{-1}x.$$

*Example 2.* Find the sum of  $n$  terms of the series

$$\tan a + \frac{1}{2} \tan \frac{a}{2} + \frac{1}{2^2} \tan \frac{a}{2^2} + \frac{1}{2^3} \tan \frac{a}{2^3} + \dots$$

We have  $\tan a = \cot a - 2 \cot 2a$ .

Replacing  $a$  by  $\frac{a}{2}$  and dividing by 2, we obtain

$$\frac{1}{2} \tan \frac{a}{2} = \frac{1}{2} \cot \frac{a}{2} - \cot a.$$

Similarly,  $\frac{1}{2^2} \tan \frac{a}{2^2} = \frac{1}{2^2} \cot \frac{a}{2^2} - \frac{1}{2} \cot \frac{a}{2};$

.....

$$\frac{1}{2^{n-1}} \tan \frac{a}{2^{n-1}} = \frac{1}{2^{n-1}} \cot \frac{a}{2^{n-1}} - \frac{1}{2^{n-2}} \cot \frac{a}{2^{n-2}}.$$

By addition,  $S = \frac{1}{2^{n-1}} \cot \frac{a}{2^{n-1}} - 2 \cot 2a.$

### EXAMPLES. XXIII. b.

Sum each of the following series to  $n$  terms :

1.  $\cos^2 \theta + \cos^2 3\theta + \cos^2 5\theta + \dots$

2.  $\sin^2 a + \sin^2 \left(a + \frac{\pi}{n}\right) + \sin^2 \left(a + \frac{2\pi}{n}\right) + \dots$

3.  $\cos^2 a + \cos^2 \left(a - \frac{\pi}{n}\right) + \cos^2 \left(a - \frac{2\pi}{n}\right) + \dots$

4.  $\sin^3 \theta + \sin^3 2\theta + \sin^3 3\theta + \dots$

5.  $\sin^3 a + \sin^3 \left(a + \frac{2\pi}{n}\right) + \sin^3 \left(a + \frac{4\pi}{n}\right) + \dots$

6.  $\cos^3 a + \cos^3 \left(a - \frac{2\pi}{n}\right) + \cos^3 \left(a - \frac{4\pi}{n}\right) + \dots$

7.  $\tan \theta + 2 \tan 2\theta + 2^2 \tan 2^2\theta + \dots$

8.  $\frac{1}{\cos a + \cos 3a} + \frac{1}{\cos a + \cos 5a} + \frac{1}{\cos a + \cos 7a} + \dots$



$$9. \quad \sin^2 \theta \sin 2\theta + \frac{1}{2} \sin^2 2\theta \sin 4\theta + \frac{1}{4} \sin^2 4\theta \sin 8\theta + \dots$$

$$10. \quad 2 \cos \theta \sin^2 \frac{\theta}{2} + 2^2 \cos \frac{\theta}{2} \sin^2 \frac{\theta}{2^2} + 2^3 \cos \frac{\theta}{2^2} \sin^2 \frac{\theta}{2^3} + \dots$$

$$11. \quad \tan^{-1} \frac{x}{1 \cdot 2 + x^2} + \tan^{-1} \frac{x}{2 \cdot 3 + x^2} + \tan^{-1} \frac{x}{3 \cdot 4 + x^2} + \dots$$

$$12. \quad \tan^{-1} \frac{1}{1+1+1^2} + \tan^{-1} \frac{1}{1+2+2^2} + \tan^{-1} \frac{1}{1+3+3^2} + \dots$$

$$13. \quad \tan^{-1} \frac{2}{2+1^2+1^4} + \tan^{-1} \frac{4}{2+2^2+2^4} + \tan^{-1} \frac{6}{2+3^2+3^4} + \dots$$

$$14. \quad \tan^{-1} \frac{2}{1-1^2+1^4} + \tan^{-1} \frac{4}{1-2^2+2^4} + \tan^{-1} \frac{6}{1-3^2+3^4} + \dots$$

15. From any point on the circumference of a circle of radius  $r$ , chords are drawn to the angular points of the regular inscribed polygon of  $n$  sides: shew that the sum of the squares of the chords is  $2nr^2$ .

16. From a point  $P$  within a regular polygon of  $2n$  sides, perpendiculars  $PA_1, PA_2, PA_3, \dots, PA_{2n}$  are drawn to the sides: shew that

$$PA_1 + PA_3 + \dots + PA_{2n-1} = PA_2 + PA_4 + \dots + PA_{2n} = nr,$$

where  $r$  is the radius of the inscribed circle.

17. If  $A_1 A_2 A_3 \dots A_{2n+1}$  is a regular polygon and  $P$  a point on the circumscribed circle lying on the arc  $A_1 A_{2n+1}$ , shew that

$$PA_1 + PA_2 + \dots + PA_{2n+1} = PA_2 + PA_4 + \dots + PA_{2n}.$$

18. From any point on the circumference of a circle, perpendiculars are drawn to the sides of the regular circumscribing polygon of  $n$  sides: shew that

$$(1) \quad \text{the sum of the squares of the perpendiculars is } \frac{3nr^2}{2};$$

$$(2) \quad \text{the sum of the cubes of the perpendiculars is } \frac{5nr^2}{2}.$$

## CHAPTER XXIV.

### MISCELLANEOUS TRANSFORMATIONS AND IDENTITIES.

#### Symmetrical Expressions.

301. An expression is said to be *symmetrical* with respect to certain of the letters it contains, if the value of the expression remains unaltered when any pair of these letters are interchanged. Thus

$$\begin{aligned} \cos a + \cos \beta + \cos \gamma, \quad \sin a \sin \beta \sin \gamma, \\ \tan (a - \theta) + \tan (\beta - \theta) + \tan (\gamma - \theta), \end{aligned}$$

are expressions which are symmetrical with respect to the letters  $a, \beta, \gamma$ .

302. A symmetrical expression involving the *sum* of a number of quantities may be concisely denoted by writing down one of the *terms* and prefixing the symbol  $\Sigma$ . Thus  $\Sigma \cos a$  stands for the sum of all the terms of which  $\cos a$  is the type,  $\Sigma \sin a \sin \beta$  stands for the sum of all the terms of which  $\sin a \sin \beta$  is the type; and so on.

For instance, if the expression is symmetrical with respect to the three letters  $a, \beta, \gamma$ ,

$$\begin{aligned} \Sigma \cos \beta \cos \gamma &= \cos \beta \cos \gamma + \cos \gamma \cos a + \cos a \cos \beta; \\ \Sigma \sin (a - \theta) &= \sin (a - \theta) + \sin (\beta - \theta) + \sin (\gamma - \theta). \end{aligned}$$

303. A symmetrical expression involving the *product* of a number of quantities may be denoted by writing down one of the *factors* and prefixing the symbol  $\Pi$ . Thus  $\Pi \sin a$  stands for the product of all the factors of which  $\sin a$  is the type.

For instance, if the expression is symmetrical with respect to the three letters  $a, \beta, \gamma$ ,

$$\begin{aligned} \Pi \tan (a + \theta) &= \tan (a + \theta) \tan (\beta + \theta) \tan (\gamma + \theta); \\ \Pi (\cos \beta + \cos \gamma) &= (\cos \beta + \cos \gamma) (\cos \gamma + \cos a) (\cos a + \cos \beta). \end{aligned}$$

304. With the notation just explained, certain theorems in Chap. XII. involving the three angles  $A, B, C$ , which are connected by the relation  $A+B+C=180^\circ$ , may be written more concisely. For instance

$$\Sigma \sin 2A = 4\Pi \sin A;$$

$$\Sigma \sin A = 4\Pi \cos \frac{A}{2};$$

$$\Sigma \tan A = \Pi \tan A;$$

$$\Sigma \tan \frac{B}{2} \tan \frac{C}{2} = 1.$$

*Example 1.* Find the ratios of  $a : b : c$  from the equations

$$a \cos \theta + b \sin \theta = c \quad \text{and} \quad a \cos \phi + b \sin \phi = c.$$

From the given equations, we have

$$a \cos \theta + b \sin \theta - c = 0,$$

and

$$a \cos \phi + b \sin \phi - c = 0;$$

whence by cross multiplication

$$\frac{a}{\sin \phi - \sin \theta} = \frac{b}{\cos \theta - \cos \phi} = \frac{c}{\sin \phi \cos \theta - \cos \phi \sin \theta};$$

$$\therefore \frac{a}{2 \cos \frac{\phi + \theta}{2} \sin \frac{\phi - \theta}{2}} = \frac{b}{2 \sin \frac{\phi + \theta}{2} \sin \frac{\phi - \theta}{2}} = \frac{c}{\sin (\phi - \theta)}.$$

Dividing each denominator by  $2 \sin \frac{\phi - \theta}{2}$ , we have

$$\frac{a}{\cos \frac{\theta + \phi}{2}} = \frac{b}{\sin \frac{\theta + \phi}{2}} = \frac{c}{\cos \frac{\theta - \phi}{2}}.$$

**NOTE.** This result is important in Analytical Geometry.

It should be remarked that  $\cos (\theta - \phi)$  is a symmetrical function of  $\theta$  and  $\phi$ , for  $\cos (\theta - \phi) = \cos (\phi - \theta)$ ; hence the values obtained for  $a : b : c$  involve  $\theta$  and  $\phi$  symmetrically.

*Example 2.* If  $\alpha$  and  $\beta$  are two different values of  $\theta$  which satisfy the equation  $a \cos \theta + b \sin \theta = c$ , find the values of

$$4 \cos^2 \frac{\alpha}{2} \cos^2 \frac{\beta}{2}, \quad \sin \alpha + \sin \beta, \quad \sin \alpha \sin \beta.$$

From the given equation, by transposing and squaring,

$$(a \cos \theta - c)^2 = b^2 \sin^2 \theta = b^2 (1 - \cos^2 \theta);$$

$$\therefore (a^2 + b^2) \cos^2 \theta - 2ac \cos \theta + c^2 - b^2 = 0.$$

The roots of this quadratic in  $\cos \theta$  are  $\cos \alpha$  and  $\cos \beta$ ;

$$\therefore \cos \alpha + \cos \beta = \frac{2ac}{a^2 + b^2} \dots\dots\dots(1),$$

and

$$\cos \alpha \cos \beta = \frac{c^2 - b^2}{a^2 + b^2} \dots\dots\dots(2).$$

$$\text{And} \quad 4 \cos^2 \frac{\alpha}{2} \cos^2 \frac{\beta}{2} = (1 + \cos \alpha)(1 + \cos \beta)$$

$$= 1 + \frac{2ac}{a^2 + b^2} + \frac{c^2 - b^2}{a^2 + b^2}$$

$$= \frac{(a + c)^2}{a^2 + b^2}.$$

From the data, we see that  $\frac{\pi}{2} - \alpha$  and  $\frac{\pi}{2} - \beta$  are values of  $\theta$  which satisfy the equation  $a \sin \theta + b \cos \theta = c$ .

By writing  $a$  for  $b$  and  $b$  for  $a$ , equation (1) becomes

$$\cos \left( \frac{\pi}{2} - \alpha \right) + \cos \left( \frac{\pi}{2} - \beta \right) = \frac{2bc}{b^2 + a^2},$$

or

$$\sin \alpha + \sin \beta = \frac{2bc}{a^2 + b^2}.$$

Similarly, from equation (2) we have

$$\sin \alpha \sin \beta = \frac{c^2 - a^2}{a^2 + b^2}.$$

These last two results may also be derived from the equation

$$(b \sin \theta - c)^2 = a^2 \cos^2 \theta = a^2 (1 - \sin^2 \theta).$$

*Example 3.* If  $\alpha$  and  $\beta$  are two different values of  $\theta$  which satisfy the equation  $a \cos \theta + b \sin \theta = c$ , prove that  $\tan \frac{\alpha + \beta}{2} = \frac{b}{a}$ . Also if the values of  $\alpha$  and  $\beta$  are equal, shew that  $a^2 + b^2 = c^2$ .

$$\text{By substituting} \quad \cos \theta = \frac{1 - \tan^2 \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}} \quad \text{and} \quad \sin \theta = \frac{2 \tan \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}}$$

in the given equation  $a \cos \theta + b \sin \theta = c$ , we have

$$a \left( 1 - \tan^2 \frac{\theta}{2} \right) + 2b \tan \frac{\theta}{2} = c \left( 1 + \tan^2 \frac{\theta}{2} \right);$$

that is,  $(c+a) \tan^2 \frac{\theta}{2} - 2b \tan \frac{\theta}{2} + (c-a) = 0 \dots\dots\dots(1).$

The roots of this equation are  $\tan \frac{\alpha}{2}$  and  $\tan \frac{\beta}{2}$ ;

$$\therefore \tan \frac{\alpha}{2} + \tan \frac{\beta}{2} = \frac{2b}{c+a}, \quad \text{and} \quad \tan \frac{\alpha}{2} \tan \frac{\beta}{2} = \frac{c-a}{c+a};$$

$$\therefore \tan \frac{\alpha+\beta}{2} = \frac{2b}{c+a} \bigg/ \left( 1 - \frac{c-a}{c+a} \right) = \frac{b}{a}.$$

If the roots of equation (1) are equal, we have

$$b^2 = (c+a)(c-a);$$

whence  $a^2 + b^2 = c^2.$

NOTE. The substitution here employed is frequently used in Analytical Geometry.

*Example 4.* If  $\cos \theta + \cos \phi = a$  and  $\sin \theta + \sin \phi = b$ , find the values of  $\cos (\theta + \phi)$  and  $\sin 2\theta + \sin 2\phi$ .

From the given equations, we have

$$\frac{\sin \theta + \sin \phi}{\cos \theta + \cos \phi} = \frac{b}{a};$$

$$\therefore \tan \frac{\theta + \phi}{2} = \frac{b}{a}.$$

For shortness write  $t$  instead of  $\tan \frac{\theta + \phi}{2}$ ; then

$$\cos (\theta + \phi) = \frac{1 - t^2}{1 + t^2} = \frac{a^2 - b^2}{a^2 + b^2},$$

and

$$\sin (\theta + \phi) = \frac{2t}{1 + t^2} = \frac{2ab}{a^2 + b^2}.$$

Multiplying the two given equations together, we have

$$\sin 2\theta + \sin 2\phi + 2 \sin (\theta + \phi) = 2ab;$$

$$\therefore \sin 2\theta + \sin 2\phi = 2ab \left( 1 - \frac{2}{a^2 + b^2} \right).$$

*Example 5.* Resolve into factors the expression

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + 2 \cos \alpha \cos \beta \cos \gamma - 1,$$

and shew that it vanishes if any one of the four angles  $\alpha \pm \beta \pm \gamma$  is an odd multiple of two right angles.

$$\begin{aligned} \text{The expression} &= \cos^2 \alpha + (\cos^2 \beta + \cos^2 \gamma - 1) + 2 \cos \alpha \cos \beta \cos \gamma \\ &= \cos^2 \alpha + (\cos^2 \beta - \sin^2 \gamma) + 2 \cos \alpha \cos \beta \cos \gamma \\ &= \cos^2 \alpha + \cos (\beta + \gamma) \cos (\beta - \gamma) + \cos \alpha \{ \cos (\beta + \gamma) + \cos (\beta - \gamma) \} \\ &= \{ \cos \alpha + \cos (\beta + \gamma) \} \{ \cos \alpha + \cos (\beta - \gamma) \} \\ &= 4 \cos \frac{\alpha + \beta + \gamma}{2} \cos \frac{\alpha - \beta - \gamma}{2} \cos \frac{\alpha + \beta - \gamma}{2} \cos \frac{\alpha - \beta + \gamma}{2}. \end{aligned}$$

The expression vanishes if one of the quantities  $\cos \frac{\alpha \pm \beta \pm \gamma}{2} = 0$ ;

that is, if one of the four angles  $\frac{\alpha \pm \beta \pm \gamma}{2} = (2n+1) \frac{\pi}{2}$ ;

that is, if  $\alpha \pm \beta \pm \gamma = (2n+1) \pi$ , where  $n$  is any integer.

*Example 6.* If  $\tan \theta = \frac{\sin \alpha \sin \beta}{\cos \alpha + \cos \beta}$ ,

prove that one value of  $\tan \frac{\theta}{2}$  is  $\tan \frac{\alpha}{2} \tan \frac{\beta}{2}$ .

From the given equation, we have

$$\begin{aligned} \sec^2 \theta &= 1 + \frac{\sin^2 \alpha \sin^2 \beta}{(\cos \alpha + \cos \beta)^2} = \frac{(\cos \alpha + \cos \beta)^2 + (1 - \cos^2 \alpha)(1 - \cos^2 \beta)}{(\cos \alpha + \cos \beta)^2} \\ &= \frac{1 + 2 \cos \alpha \cos \beta + \cos^2 \alpha \cos^2 \beta}{(\cos \alpha + \cos \beta)^2}. \end{aligned}$$

Taking the positive root,  $\sec \theta = \frac{1 + \cos \alpha \cos \beta}{\cos \alpha + \cos \beta}$ ;

$$\therefore \cos \theta = \frac{\cos \alpha + \cos \beta}{1 + \cos \alpha \cos \beta}.$$

$$\therefore \frac{1 - \cos \theta}{1 + \cos \theta} = \frac{1 - \cos \alpha - \cos \beta + \cos \alpha \cos \beta}{1 + \cos \alpha + \cos \beta + \cos \alpha \cos \beta} = \frac{(1 - \cos \alpha)(1 - \cos \beta)}{(1 + \cos \alpha)(1 + \cos \beta)};$$

$$\therefore \tan^2 \frac{\theta}{2} = \tan^2 \frac{\alpha}{2} \tan^2 \frac{\beta}{2};$$

and therefore one value of  $\tan \frac{\theta}{2}$  is  $\tan \frac{\alpha}{2} \tan \frac{\beta}{2}$ .



*Example 7.* In any triangle, shew that

$$\Sigma a^3 \cos A = abc (1 + 4\Pi \cos A).$$

Let 
$$k = \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C};$$

so that 
$$a = k \sin A, \quad b = k \sin B, \quad c = k \sin C.$$

By substituting these values in the given identity, and dividing by  $k^3$ , we have to prove that

$$\Sigma \sin^3 A \cos A = \sin A \sin B \sin C (1 + 4\Pi \cos A).$$

Now 
$$\begin{aligned} 8\Sigma \sin^3 A \cos A &= 4\Sigma \sin^2 A \sin 2A \\ &= 2\Sigma (1 - \cos 2A) \sin 2A \\ &= 2\Sigma \sin 2A - \Sigma \sin 4A; \end{aligned}$$

and it has been shewn in Example 1, Art. 135, that

$$\Sigma \sin 2A = 4\Pi \sin A;$$

and it is easy to prove that

$$\begin{aligned} \Sigma \sin 4A &= -4\Pi \sin 2A = -32\Pi \sin A \cdot \Pi \cos A; \\ \therefore 8\Sigma \sin^3 A \cos A &= 8\Pi \sin A + 32\Pi \sin A \cdot \Pi \cos A; \\ \therefore \Sigma \sin^3 A \cos A &= \Pi \sin A (1 + 4\Pi \cos A). \end{aligned}$$

### EXAMPLES. XXIV. a.

1. If  $\theta = a$ , and  $\theta = \beta$  satisfy the equation

$$\frac{1}{a} \cos \theta + \frac{1}{b} \sin \theta = \frac{1}{c},$$

prove that 
$$a \cos \frac{a+\beta}{2} = b \sin \frac{a+\beta}{2} = c \cos \frac{a-\beta}{2}.$$

Solve the simultaneous equations :

$$2. \quad \frac{x}{a} \cos a + \frac{y}{b} \sin a = 1, \quad \frac{x}{a} \cos \beta + \frac{y}{b} \sin \beta = 1.$$

$$3. \quad \frac{x}{a} \cos a + \frac{y}{b} \sin a = 1, \quad \frac{x}{a} \sin a - \frac{y}{b} \cos a = 1.$$

If  $a$  and  $\beta$  are two different solutions of  $a \cos \theta + b \sin \theta = c$ , prove that

$$4. \quad \cos(a + \beta) = \frac{a^2 - b^2}{a^2 + b^2}. \quad 5. \quad \cos^2 \frac{a - \beta}{2} = \frac{c^2}{a^2 + b^2}.$$

$$6. \quad \sin 2a + \sin 2\beta = \frac{4ab(2c^2 - a^2 - b^2)}{(a^2 + b^2)^2}.$$

$$7. \quad \sin^2 a + \sin^2 \beta = \frac{2a^2(a^2 + b^2) - 2c^2(a^2 - b^2)}{(a^2 + b^2)^2}.$$

8. If  $a \cos a + b \sin a = a \cos \beta + b \sin \beta = c$ , prove that

$$\sin(a + \beta) = \frac{2ab}{a^2 + b^2}, \quad \text{and} \quad \cot a + \cot \beta = \frac{2ab}{c^2 - a^2}.$$

If  $\cos \theta + \cos \phi = a$  and  $\sin \theta + \sin \phi = b$ , prove that

$$9. \quad \cos \theta \cos \phi = \frac{(a^2 + b^2)^2 - 4b^2}{4(a^2 + b^2)}.$$

$$10. \quad \cos 2\theta + \cos 2\phi = \frac{(a^2 - b^2)(a^2 + b^2 - 2)}{a^2 + b^2}.$$

$$11. \quad \tan \theta + \tan \phi = \frac{8ab}{(a^2 + b^2)^2 - 4b^2}.$$

$$12. \quad \tan \frac{\theta}{2} + \tan \frac{\phi}{2} = \frac{4b}{a^2 + b^2 + 2a}.$$

13. Express

$$1 - \cos^2 a - \cos^2 \beta - \cos^2 \gamma + 2 \cos a \cos \beta \cos \gamma$$

as the product of four sines, and shew that it vanishes if any one of the four angles  $a \pm \beta \pm \gamma$  is zero or an even multiple of  $\pi$ .

14. Express

$$\sin^2 a + \sin^2 \beta - \sin^2 \gamma + 2 \sin a \sin \beta \cos \gamma$$

as the product of two sines and two cosines.

15. Express

$$\sin^2 a + \sin^2 \beta + \sin^2 \gamma - 2 \sin a \sin \beta \sin \gamma - 1$$

as the product of four cosines.

16. If  $\cos \theta = \frac{\cos a - \cos \beta}{1 - \cos a \cos \beta},$

prove that one value of  $\tan \frac{\theta}{2}$  is  $\tan \frac{a}{2} \cot \frac{\beta}{2}.$

17. If  $\tan^2 \theta \cos^2 \frac{a+\beta}{2} = \sin a \sin \beta,$

prove that one value of  $\tan^2 \frac{\theta}{2}$  is  $\tan \frac{a}{2} \tan \frac{\beta}{2}.$

18. If  $\tan \theta (\cos a + \sin \beta) = \sin a \cos \beta,$

prove that one value of  $\tan \frac{\theta}{2}$  is  $\tan \frac{a}{2} \tan \left( \frac{\pi}{4} - \frac{\beta}{2} \right).$

In any triangle, shew that

19.  $\Sigma a^3 \sin B \sin C = 2abc (1 + \cos A \cos B \cos C).$

20.  $\Sigma a \cos^3 A = \frac{abc}{4R^2} (1 - 4 \cos A \cos B \cos C).$

21.  $\Sigma a^3 \cos (B - C) = 3abc.$

22. If  $a$  and  $\beta$  are roots of the equation  $a \cos \theta + b \sin \theta = c,$  form the equations whose roots are

(1)  $\sin a$  and  $\sin \beta$ ; (2)  $\cos 2a$  and  $\cos 2\beta.$

### Alternating Expressions.

305. An expression is said to be *alternating* with respect to certain of the letters it contains, if the sign of the expression but not its numerical value is altered when any pair of these letters are interchanged.

Thus  $\cos a - \cos \beta, \sin (a - \beta), \tan (a - \beta),$

$\cos^2 a \sin (\beta - \gamma) + \cos^2 \beta \sin (\gamma - a) + \cos^2 \gamma \sin (a - \beta)$

are alternating expressions.

306. Alternating expressions may be abridged by means of the symbols  $\Sigma$  and  $\Pi$ . Thus

$\Sigma \sin^2 a \sin (\beta - \gamma) = \sin^2 a \sin (\beta - \gamma) + \sin^2 \beta \sin (\gamma - a)$   
 $+ \sin^2 \gamma \sin (a - \beta);$

$\Pi \tan (\beta - \gamma) = \tan (\beta - \gamma) \tan (\gamma - a) \tan (a - \beta).$

We shall confine our attention chiefly to alternating expressions involving the three letters  $\alpha$ ,  $\beta$ ,  $\gamma$ , and we shall adopt the *cyclical arrangement*  $\beta - \gamma$ ,  $\gamma - \alpha$ ,  $\alpha - \beta$  in which  $\beta$  follows  $\alpha$ ,  $\gamma$  follows  $\beta$ , and  $\alpha$  follows  $\gamma$ .

*Example 1.* Prove that  $\Sigma \cos (\alpha + \theta) \sin (\beta - \gamma) = 0$ .

$$\begin{aligned}\Sigma \cos (\alpha + \theta) \sin (\beta - \gamma) &= \Sigma (\cos \alpha \cos \theta - \sin \alpha \sin \theta) \sin (\beta - \gamma) \\ &= \cos \theta \Sigma \cos \alpha \sin (\beta - \gamma) - \sin \theta \Sigma \sin \alpha \sin (\beta - \gamma) \\ &= 0,\end{aligned}$$

since  $\Sigma \cos \alpha \sin (\beta - \gamma) = 0$  and  $\Sigma \sin \alpha \sin (\beta - \gamma) = 0$ .

*Example 2.* Shew that  $\Sigma \sin 2 (\beta - \gamma) = -4\Pi \sin (\beta - \gamma)$ .

$$\begin{aligned}\sin 2 (\beta - \gamma) + \sin 2 (\gamma - \alpha) + \sin 2 (\alpha - \beta) \\ &= 2 \sin (\beta - \alpha) \cos (\alpha + \beta - 2\gamma) + 2 \sin (\alpha - \beta) \cos (\alpha - \beta) \\ &= 2 \sin (\alpha - \beta) \{ \cos (\alpha - \beta) - \cos (\alpha + \beta - 2\gamma) \} \\ &= 4 \sin (\alpha - \beta) \sin (\alpha - \gamma) \sin (\beta - \gamma) \\ &= -4\Pi \sin (\beta - \gamma).\end{aligned}$$

*Example 3.* Prove that

$$(1) \quad \Sigma \tan (\beta - \gamma) = \Pi \tan (\beta - \gamma);$$

$$(2) \quad \Sigma \tan \beta \tan \gamma \tan (\beta - \gamma) = -\Pi \tan (\beta - \gamma).$$

(1) From Art. 118, if  $A + B + C = 0$ , we see that

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

Hence by writing  $A = \beta - \gamma$ ,  $B = \gamma - \alpha$ ,  $C = \alpha - \beta$ , we have

$$\Sigma \tan (\beta - \gamma) = \Pi \tan (\beta - \gamma).$$

(2) From the formulæ for  $\tan (\beta - \gamma)$ ,  $\tan (\gamma - \alpha)$ ,  $\tan (\alpha - \beta)$ , we have

$$\Sigma (1 + \tan \beta \tan \gamma) \tan (\beta - \gamma) = \Sigma (\tan \beta - \tan \gamma) = 0;$$

whence by transposition

$$\begin{aligned}\Sigma \tan \beta \tan \gamma \tan (\beta - \gamma) &= -\Sigma \tan (\beta - \gamma) \\ &= -\Pi \tan (\beta - \gamma).\end{aligned}$$

*Example 4.* Shew that

$$\Sigma \cos 3\alpha \sin (\beta - \gamma) = 4 \cos (\alpha + \beta + \gamma) \Pi \sin (\beta - \gamma).$$

Since  $2 \cos 3\alpha \sin (\beta - \gamma) = \sin (3\alpha + \beta - \gamma) - \sin (3\alpha - \beta + \gamma)$ ,  
we have

$$2\Sigma \cos 3\alpha \sin (\beta - \gamma) = \sin (3\alpha + \beta - \gamma) - \sin (3\alpha - \beta + \gamma) + \sin (3\beta + \gamma - \alpha) \\ - \sin (3\beta - \gamma + \alpha) + \sin (3\gamma + \alpha - \beta) - \sin (3\gamma - \alpha + \beta).$$

Combining the second and third terms, the fourth and fifth terms, the sixth and first terms, and dividing by 2, we have

$$\Sigma \cos 3\alpha \sin (\beta - \gamma) \\ = \cos (\alpha + \beta + \gamma) \{ \sin 2 (\beta - \alpha) + \sin 2 (\gamma - \beta) + \sin 2 (\alpha - \gamma) \} \\ = 4 \cos (\alpha + \beta + \gamma) \Pi \sin (\beta - \gamma). \quad [\text{See Example 2.}]$$

307. The following example is given as a specimen of a concise solution.

*Example.* If  $(y+z) \tan \alpha + (z+x) \tan \beta + (x+y) \tan \gamma = 0$ ,  
and  $x \tan \beta \tan \gamma + y \tan \gamma \tan \alpha + z \tan \alpha \tan \beta = x + y + z$ ,  
prove that  $x \sin 2\alpha + y \sin 2\beta + z \sin 2\gamma = 0$ .

From the given equations, we have

$$x (1 - \tan \beta \tan \gamma) + y (1 - \tan \gamma \tan \alpha) + z (1 - \tan \alpha \tan \beta) = 0, \\ \text{and} \quad x (\tan \beta + \tan \gamma) + y (\tan \gamma + \tan \alpha) + z (\tan \alpha + \tan \beta) = 0.$$

If we find the values of  $x : y : z$  by cross multiplication, the denominator of  $x$

$$= (1 - \tan \gamma \tan \alpha) (\tan \alpha + \tan \beta) - (1 - \tan \alpha \tan \beta) (\tan \gamma + \tan \alpha) \\ = (\tan \beta - \tan \gamma) + \tan^2 \alpha (\tan \beta - \tan \gamma) \\ = (1 + \tan^2 \alpha) (\tan \beta - \tan \gamma) \\ = \sec^2 \alpha (\tan \beta - \tan \gamma) \\ = \frac{\sec \alpha \sin (\beta - \gamma)}{\cos \alpha \cos \beta \cos \gamma}.$$

$$\text{Hence} \quad \frac{x}{\sec \alpha \sin (\beta - \gamma)} = \frac{y}{\sec \beta \sin (\gamma - \alpha)} = \frac{z}{\sec \gamma \sin (\alpha - \beta)} = k \text{ say.}$$

$$\therefore x \sin 2\alpha + y \sin 2\beta + z \sin 2\gamma = k \Sigma \sin 2\alpha \sec \alpha \sin (\beta - \gamma) \\ = 2k \Sigma \sin \alpha \sin (\beta - \gamma) \\ = 0.$$

### Allied formulæ in Algebra and Trigonometry.

308. From well-known algebraical identities we can deduce some interesting trigonometrical identities.

*Example 1.* In the identity

$$(x-a)(b-c) + (x-b)(c-a) + (x-c)(a-b) = 0,$$

put  $x = \cos 2\theta$ ,  $a = \cos 2\alpha$ ,  $b = \cos 2\beta$ ,  $c = \cos 2\gamma$ ;

then  $x - a = \cos 2\theta - \cos 2\alpha = 2 \sin (\alpha + \theta) \sin (\alpha - \theta)$ ,

and  $b - c = \cos 2\beta - \cos 2\gamma = -2 \sin (\beta + \gamma) \sin (\beta - \gamma)$ ;

$$\therefore \Sigma \sin (\alpha + \theta) \sin (\alpha - \theta) \sin (\beta + \gamma) \sin (\beta - \gamma) = 0.$$

*Example 2.* In the identity

$$\Sigma a^2 (b-c) = -\Pi (b-c),$$

put  $a = \sin^2 \alpha$ ,  $b = \sin^2 \beta$ ,  $c = \sin^2 \gamma$ ;

then  $b - c = \sin^2 \beta - \sin^2 \gamma = \sin (\beta + \gamma) \sin (\beta - \gamma)$ ;

$$\therefore \Sigma \sin^4 \alpha \sin (\beta + \gamma) \sin (\beta - \gamma) = -\Pi \sin (\beta + \gamma) \cdot \Pi \sin (\beta - \gamma).$$

*Example 3.* In the identity

$$\Sigma a^3 (b-c) = -(a+b+c) \Pi (b-c),$$

put  $a = \cos \alpha$ ,  $b = \cos \beta$ ,  $c = \cos \gamma$ ;

$$\therefore \Sigma \cos^3 \alpha (\cos \beta - \cos \gamma) = -(\cos \alpha + \cos \beta + \cos \gamma) \Pi (\cos \beta - \cos \gamma).$$

But  $\Sigma \cos \alpha (\cos \beta - \cos \gamma) = 0$ ;

$$\begin{aligned} \therefore \Sigma (4 \cos^3 \alpha - 3 \cos \alpha) (\cos \beta - \cos \gamma) \\ = -4 (\cos \alpha + \cos \beta + \cos \gamma) \Pi (\cos \beta - \cos \gamma); \end{aligned}$$

that is,

$$\Sigma \cos^3 \alpha (\cos \beta - \cos \gamma) = -4 (\cos \alpha + \cos \beta + \cos \gamma) \Pi (\cos \beta - \cos \gamma).$$

*Example 4.* If  $a+b+c=0$ , then  $a^3+b^3+c^3=3abc$ .

Here  $a, b, c$  may be any three quantities whose sum is zero; this condition is satisfied if we put  $a = \cos (\alpha + \theta) \sin (\beta - \gamma)$ , and  $b$  and  $c$  equal to corresponding quantities.

$$\text{Thus } \Sigma \cos^3 (\alpha + \theta) \sin^3 (\beta - \gamma) = 3 \Pi \cos (\alpha + \theta) \sin (\beta - \gamma).$$



309. An algebraical identity may sometimes be established by the aid of Trigonometry.

*Example.* If  $x + y + z = xyz$ , prove that

$$x(1-y^2)(1-z^2) + y(1-z^2)(1-x^2) + z(1-x^2)(1-y^2) = 4xyz.$$

By putting  $x = \tan \alpha$ ,  $y = \tan \beta$ ,  $z = \tan \gamma$ , we have

$$\tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma;$$

whence 
$$\tan \alpha = -\frac{\tan \beta + \tan \gamma}{1 - \tan \beta \tan \gamma} = -\tan(\beta + \gamma);$$

$$\therefore \alpha = n\pi - (\beta + \gamma), \text{ where } n \text{ is an integer};$$

$$\therefore \alpha + \beta + \gamma = n\pi;$$

$$\therefore 2\alpha + 2\beta + 2\gamma = 2n\pi.$$

From this relation it is easy to shew that

$$\tan 2\alpha + \tan 2\beta + \tan 2\gamma = \tan 2\alpha \tan 2\beta \tan 2\gamma;$$

$$\therefore \frac{2x}{1-x^2} + \frac{2y}{1-y^2} + \frac{2z}{1-z^2} = \frac{8xyz}{(1-x^2)(1-y^2)(1-z^2)};$$

$$\therefore x(1-y^2)(1-z^2) + y(1-z^2)(1-x^2) + z(1-x^2)(1-y^2) = 4xyz.$$

### EXAMPLES. XXIV. b.

Prove the following identities :

1.  $\Sigma \sin(\alpha - \theta) \sin(\beta - \gamma) = 0.$
2.  $\Sigma \cos \beta \cos \gamma \sin(\beta - \gamma) = \Sigma \sin \beta \sin \gamma \sin(\beta - \gamma).$
3.  $\Sigma \sin(\beta - \gamma) \cos(\beta + \gamma + \theta) = 0.$
4.  $\Sigma \cos 2(\beta - \gamma) = 4\Pi \cos(\beta - \gamma) - 1.$
5.  $\Sigma \sin \beta \sin \gamma \sin(\beta - \gamma) = -\Pi \sin(\beta - \gamma).$
6.  $\Sigma \cot(\alpha - \beta) \cot(\alpha - \gamma) + 1 = 0.$
7.  $\Sigma \sin 3\alpha \sin(\beta - \gamma) = 4 \sin(\alpha + \beta + \gamma) \Pi \sin(\beta - \gamma).$
8.  $\Sigma \cos^3 \alpha \sin(\beta - \gamma) = \cos(\alpha + \beta + \gamma) \Pi \sin(\beta - \gamma).$
9.  $\Sigma \cos(\theta + \alpha) \cos(\beta + \gamma) \sin(\theta - \alpha) \sin(\beta - \gamma) = 0.$
10.  $\Sigma \sin^2 \beta \sin^2 \gamma \sin(\beta + \gamma) \sin(\beta - \gamma)$   

$$= -\Pi \sin(\beta + \gamma) \cdot \Pi \sin(\beta - \gamma).$$

Prove the following identities :

$$11. \quad \Sigma \cos 2\beta \cos 2\gamma \sin (\beta + \gamma) \sin (\beta - \gamma) \\ = -4\Pi \sin (\beta + \gamma) \cdot \Pi \sin (\beta - \gamma).$$

$$12. \quad \Sigma \cos 4a \sin (\beta + \gamma) \sin (\beta - \gamma) \\ = -8\Pi \sin (\beta + \gamma) \cdot \Pi \sin (\beta - \gamma).$$

$$13. \quad \Sigma \sin 3a (\sin \beta - \sin \gamma) \\ = 4 (\sin a + \sin \beta + \sin \gamma) \Pi (\sin \beta - \sin \gamma).$$

$$14. \quad \Sigma \sin^3 (\beta + \gamma) \sin^3 (\beta - \gamma) = 3\Pi \sin (\beta + \gamma) \cdot \Pi \sin (\beta - \gamma).$$

$$15. \quad \Sigma \cos^3 (\beta + \gamma + \theta) \sin^3 (\beta - \gamma) \\ = 3\Pi \cos (\beta + \gamma + \theta) \cdot \Pi \sin (\beta - \gamma).$$

$$16. \quad \text{If } x + y + z = xyz, \text{ prove that}$$

$$\Sigma \frac{3x - x^3}{1 - 3x^2} = \Pi \frac{3x - x^3}{1 - 3x^2}.$$

$$17. \quad \text{If } yz + zx + xy = 1, \text{ prove that}$$

$$\Sigma x (1 - y^2) (1 - z^2) = 4xyz.$$

310. From a trigonometrical identity many others may be derived by various substitutions.

For instance, if  $A, B, C$  are *any* angles, positive or negative, connected by the relation  $A + B + C = \pi$ , we know that

$$\sin A + \sin B + \sin C = 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}.$$

$$\text{Let} \quad A = \pi - 2a, \quad B = \pi - 2\beta, \quad C = \pi - 2\gamma;$$

$$\text{then} \quad \sin A = \sin 2a, \text{ and } \cos \frac{A}{2} = \sin a.$$

$$\text{Also} \quad 2(a + \beta + \gamma) = 3\pi - (A + B + C) = 2\pi;$$

$$\therefore a + \beta + \gamma = \pi,$$

$$\text{and} \quad \sin 2a + \sin 2\beta + \sin 2\gamma = 4 \sin a \sin \beta \sin \gamma.$$

$$\text{Again, let} \quad A = \frac{\pi}{2} - \frac{a}{2}, \quad B = \frac{\pi}{2} - \frac{\beta}{2}, \quad C = \frac{\pi}{2} - \frac{\gamma}{2};$$

$$\text{then} \quad \sin A = \cos \frac{a}{2}, \text{ and } \cos \frac{A}{2} = \cos \frac{\pi - a}{4}.$$

Also  $\alpha + \beta + \gamma = 3\pi - 2(A + B + C) = 3\pi - 2\pi;$

$$\therefore \alpha + \beta + \gamma = \pi,$$

and  $\cos \frac{\alpha}{2} + \cos \frac{\beta}{2} + \cos \frac{\gamma}{2} = 4 \cos \frac{\pi - \alpha}{4} \cos \frac{\pi - \beta}{4} \cos \frac{\pi - \gamma}{4}.$

*Example.* If  $A + B + C = \pi$ , shew that

$$\cos \frac{A}{2} + \cos \frac{B}{2} - \cos \frac{C}{2} = 4 \cos \frac{\pi + A}{4} \cos \frac{\pi + B}{4} \cos \frac{\pi - C}{4}.$$

Put  $\frac{\pi + A}{4} = \frac{\alpha}{2}, \quad \frac{\pi + B}{4} = \frac{\beta}{2}, \quad \frac{C - \pi}{4} = \frac{\gamma}{2};$

then  $\cos \frac{A}{2} = \cos \left( \alpha - \frac{\pi}{2} \right) = \sin \alpha$ , and  $\cos \frac{C}{2} = \cos \left( \gamma + \frac{\pi}{2} \right) = -\sin \gamma$ ,

so that the above identity becomes

$$\sin \alpha + \sin \beta + \sin \gamma = 4 \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2},$$

which is clearly true since

$$\alpha + \beta + \gamma = \frac{\pi}{2} + \frac{A + B + C}{2} = \frac{\pi}{2} + \frac{\pi}{2} = \pi.$$

**311.** When  $A + B + C = n\pi$ ,

$$\tan(A + B) = \tan(n\pi - C) = -\tan C;$$

whence we obtain  $\Sigma \tan A = \Pi \tan A$ .

When  $n=0$ , the given condition is satisfied in the case of any three angles whose sum is 0; as for instance if

$$A = \beta + \gamma - 2\alpha, \quad B = \gamma + \alpha - 2\beta, \quad C = \alpha + \beta - 2\gamma.$$

Hence  $\Sigma \tan(\beta + \gamma - 2\alpha) = \Pi \tan(\beta + \gamma - 2\alpha).$

*Example.* If  $\alpha + \beta + \gamma = 0$ , shew that

$$\Sigma \cot(\gamma + \alpha - \beta) \cot(\alpha + \beta - \gamma) = 1.$$

Put  $\beta + \gamma - \alpha = A, \quad \gamma + \alpha - \beta = B, \quad \alpha + \beta - \gamma = C;$

then, by addition,

$$A + B + C = \alpha + \beta + \gamma = 0;$$

$$\therefore \cot(A + B) = -\cot C;$$

whence

$$\Sigma \cot A \cot B = 1,$$

that is,

$$\Sigma \cot(\gamma + \alpha - \beta) \cot(\alpha + \beta - \gamma) = 1.$$

**312.** The following example is a further illustration of the manner in which an identity may be established by appropriate substitutions in some simpler identity.

*Example.* Prove that

$$2\Pi \cos(\beta + \gamma) + \Pi \cos 2a = \Sigma \cos 2a \cos^2(\beta + \gamma).$$

In Example 5, Art. 133, we have proved that

$$4 \cos a \cos \beta \cos \gamma = \Sigma \cos(\beta + \gamma - a) + \cos(a + \beta + \gamma).$$

In this identity first replace  $a, \beta, \gamma$  by  $\beta + \gamma, \gamma + a, a + \beta$  respectively, and secondly replace  $a, \beta, \gamma$  by  $2a, 2\beta, 2\gamma$  respectively.

$$\text{Thus} \quad 8\Pi \cos(\beta + \gamma) = 2\Sigma \cos 2a + 2 \cos 2(a + \beta + \gamma),$$

$$\text{and} \quad 4\Pi \cos 2a = \Sigma \cos 2(\beta + \gamma - a) + \cos 2(a + \beta + \gamma);$$

whence by addition

$$\begin{aligned} 8\Pi \cos(\beta + \gamma) + 4\Pi \cos 2a &= 2\Sigma \cos 2a + \Sigma \cos 2(\beta + \gamma - a) + 3 \cos 2(a + \beta + \gamma) \\ &= 2\Sigma \cos 2a + \Sigma \{\cos 2(\beta + \gamma - a) + \cos 2(a + \beta + \gamma)\} \\ &= 2\Sigma \cos 2a + 2\Sigma \cos 2(\beta + \gamma) \cos 2a \\ &= 2\Sigma \cos 2a \{1 + \cos 2(\beta + \gamma)\} \\ &= 4\Sigma \cos 2a \cos^2(\beta + \gamma); \end{aligned}$$

$$\therefore 2\Pi \cos(\beta + \gamma) + \Pi \cos 2a = \Sigma \cos 2a \cos^2(\beta + \gamma).$$

**313.** Suppose that  $A'B'C'$  is the pedal triangle of  $ABC$ , and let the sides and angles of the pedal triangle be denoted by  $a', b', c'$ , and  $A', B', C'$ , and its circum-radius by  $R'$ . Then from Arts. 224 and 225, we have

$$a' = a \cos A, \quad b' = b \cos B, \quad c' = c \cos C, \quad R' = \frac{R}{2},$$

$$A' = 180^\circ - 2A, \quad B' = 180^\circ - 2B, \quad C' = 180^\circ - 2C.$$

By means of these relations, we may from any identity proved for the triangle  $ABC$  derive another, as in the following case.

In the triangle  $ABC$ , we know that

$$\Sigma a \cos A = 4R \sin A \sin B \sin C;$$

hence in the pedal triangle  $A'B'C'$ ,

$$\Sigma a' \cos A' = 4R' \sin A' \sin B' \sin C';$$

$$\therefore \Sigma a \cos A \cos(180^\circ - 2A) = 2R\Pi \sin(180^\circ - 2A);$$

$$\text{that is,} \quad -\Sigma a \cos A \cos 2A = 2R \sin 2A \sin 2B \sin 2C.$$

*Example.* In any triangle  $ABC$ , shew that

$$\frac{a^2 \cos^2 A - b^2 \cos^2 B - c^2 \cos^2 C}{2bc \cos B \cos C} = \cos 2A.$$

In the pedal triangle  $A'B'C'$ , we have

$$\frac{b'^2 + c'^2 - a'^2}{2b'c'} = \cos A';$$

hence, by substituting the equivalents of  $a'$ ,  $b'$ ,  $c'$ ,  $A'$ , we have

$$\frac{b^2 \cos^2 B + c^2 \cos^2 C - a^2 \cos^2 A}{2bc \cos B \cos C} = \cos (180^\circ - 2A) = -\cos 2A;$$

whence the required identity follows at once.

314. If  $A_1B_1C_1$  be the ex-central triangle of  $ABC$ , we may, as in the preceding article, from any identity proved for the triangle  $ABC$  derive another by means of the relations

$$a_1 = a \operatorname{cosec} \frac{A}{2}, \quad b_1 = b \operatorname{cosec} \frac{B}{2}, \quad c_1 = c \operatorname{cosec} \frac{C}{2}, \quad R_1 = 2R,$$

$$A_1 = 90^\circ - \frac{A}{2}, \quad B_1 = 90^\circ - \frac{B}{2}, \quad C_1 = 90^\circ - \frac{C}{2}.$$

315. The following Exercise consists of miscellaneous questions on the subject of this Chapter.

### EXAMPLES. XXIV. c.

1. Shew that

$$\Sigma \cot (2a + \beta - 3\gamma) \cot (2\beta + \gamma - 3a) = 1.$$

2. Shew that

$$(1) \quad 2\Pi \sin (\beta + \gamma) + \Pi \sin 2a = \Sigma \sin 2a \sin^2 (\beta + \gamma);$$

$$(2) \quad \Pi \sin (\beta + \gamma - a) + 2\Pi \sin a = \Sigma \sin^2 a \sin (\beta + \gamma - a).$$

3. In any triangle, prove that

$$(1) \quad a^2 \cos^2 A - b^2 \cos^2 B = Rc \cos C \sin 2(B - A);$$

$$(2) \quad a^2 \operatorname{cosec}^2 \frac{A}{2} - b^2 \operatorname{cosec}^2 \frac{B}{2} = 4Rc \operatorname{cosec} \frac{C}{2} \sin \frac{B - A}{2};$$

$$(3) \quad \Sigma (b \cos B + c \cos C) \cot A = -2R \Sigma \cos 2A.$$

4. If  $\sin 2\theta = 2 \sin a \sin \gamma$ ,  
 and  $\cos 2\theta = \cos 2a \cos 2\beta = \cos 2\gamma \cos 2\delta$ ,  
 prove that one value of  $\tan \theta$  is  $\tan \beta \tan \delta$ .

5. If  $\tan \frac{\theta}{2} \tan \frac{\phi}{2} = \tan \frac{\gamma}{2}$ ,  
 and  $\sec a \cos \theta = \sec \beta \cos \phi = \cos \gamma$ ,  
 prove that  $\sin^2 \gamma = (\sec a - 1)(\sec \beta - 1)$ .

6. If  $\frac{\cos \theta - \cos a}{\cos \theta - \cos \beta} = \frac{\sin^2 a \cos \beta}{\sin^2 \beta \cos a}$ ,  
 prove that one value of  $\tan \frac{\theta}{2}$  is  $\tan \frac{a}{2} \tan \frac{\beta}{2}$ .

7. If  $\sin \theta = \cot a \tan \gamma$  and  $\tan \theta = \cos a \tan \beta$ ,  
 prove that one value of  $\cos \theta$  is  $\cos \beta \sec \gamma$ .

8. If  $a$  and  $\beta$  are two different values of  $\theta$  which satisfy  
 $bc \cos \theta \cos \phi + ac \sin \theta \sin \phi = ab$ ,  
 prove that  
 $(b^2 + c^2 - a^2) \cos a \cos \beta + (c^2 + a^2 - b^2) \sin a \sin \beta = a^2 + b^2 - c^2$ .

9. If  $\beta$  and  $\gamma$  are two different values of  $\theta$  which satisfy  
 $\sin a \cos \theta + \cos a \sin \theta = \cos a \sin a$ ,  
 prove that  $\frac{\cos \beta \cos \gamma}{\cos^2 a} + \frac{\sin \beta \sin \gamma}{\sin^2 a} = 1$ .

10. If  $\beta$  and  $\gamma$  are two different values of  $\theta$  which satisfy  
 $k^2 \cos a \cos \theta + k(\sin a + \sin \theta) + 1 = 0$ ,  
 prove that  $k^2 \cos \beta \cos \gamma + k(\sin \beta + \sin \gamma) + 1 = 0$ .

11. If  $\beta$  and  $\gamma$  are two different values of  $\theta$  which satisfy  
 $\frac{\cos \theta \cos \phi}{\cos^2 a} + \frac{\sin \theta \sin \phi}{\sin^2 a} + 1 = 0$ ,  
 prove that  $\frac{\cos \beta \cos \gamma}{\cos^2 a} + \frac{\sin \beta \sin \gamma}{\sin^2 a} + 1 = 0$ .



## CHAPTER XXV.

### MISCELLANEOUS THEOREMS AND EXAMPLES.

#### Inequalities. Maxima and Minima.

316. THE methods of proving trigonometrical inequalities are in many cases identical with those by which algebraical inequalities are established.

*Example 1.* Shew that  $a^2 \tan^2 \theta + b^2 \cot^2 \theta > 2ab$ .

We have  $a^2 \tan^2 \theta + b^2 \cot^2 \theta = (a \tan \theta - b \cot \theta)^2 + 2ab$ ;

$$\therefore a^2 \tan^2 \theta + b^2 \cot^2 \theta > 2ab,$$

unless  $a \tan \theta - b \cot \theta = 0$ , or  $a \tan^2 \theta = b$ .

In this case the inequality becomes an equality.

This proposition may be otherwise expressed by saying that the *minimum value* of  $a^2 \tan^2 \theta + b^2 \cot^2 \theta$  is  $2ab$ .

*Example 2.* Shew that

$$1 + \sin^2 \alpha + \sin^2 \beta > \sin \alpha + \sin \beta + \sin \alpha \sin \beta.$$

Since  $(1 - \sin \alpha)^2$  is positive,

$$1 + \sin^2 \alpha > 2 \sin \alpha ;$$

similarly  $1 + \sin^2 \beta > 2 \sin \beta$ ,

and  $\sin^2 \alpha + \sin^2 \beta > 2 \sin \alpha \sin \beta$ .

Adding and dividing by 2, we have

$$1 + \sin^2 \alpha + \sin^2 \beta > \sin \alpha + \sin \beta + \sin \alpha \sin \beta.$$

*Example 3.* When is  $12 \sin \theta - 9 \sin^2 \theta$  a maximum?

The expression  $= 4 - (2 - 3 \sin \theta)^2$ , and is therefore a maximum when  $2 - 3 \sin \theta = 0$ , so that its maximum value is 4.

317. To find the numerically greatest values of

$$a \cos \theta + b \sin \theta.$$

Let  $a = r \cos a$  and  $b = r \sin a$ ,

so that  $r^2 = a^2 + b^2$  and  $\tan a = \frac{b}{a}$ ;

then  $a \cos \theta + b \sin \theta = r (\cos \theta \cos a + \sin \theta \sin a)$   
 $= r \cos (\theta - a).$

Thus the expression is numerically greatest when

$$\cos (\theta - a) = \pm 1;$$

that is, the greatest positive value  $= r = \sqrt{a^2 + b^2}$ ,

and the numerically greatest negative value  $= -r = -\sqrt{a^2 + b^2}$ .

Hence, if  $c^2 > a^2 + b^2$ ,

the maximum value of  $a \cos \theta + b \sin \theta + c$  is  $c + \sqrt{a^2 + b^2}$ ,

and the minimum value is  $c - \sqrt{a^2 + b^2}$ .

318. The expression  $a \cos (a + \theta) + b \cos (\beta + \theta)$

$$= (a \cos a + b \cos \beta) \cos \theta - (a \sin a + b \sin \beta) \sin \theta;$$

and therefore its numerically greatest values are equal to the positive and negative square roots of

$$(a \cos a + b \cos \beta)^2 + (a \sin a + b \sin \beta)^2;$$

that is, are equal to

$$\pm \sqrt{a^2 + b^2 + 2ab \cos (a - \beta)}.$$

In like manner, we may find the maximum and minimum values of the sum of any number of expressions of the form  $a \cos (a + \theta)$  or  $a \sin (a + \theta)$ .

319. If  $a$  and  $\beta$  are two angles, each lying between 0 and  $\frac{\pi}{2}$ , whose sum is given, to find the maximum value of  $\cos a \cos \beta$  and of  $\cos a + \cos \beta$ .

Suppose that  $a + \beta = \sigma$ ;

then  $2 \cos a \cos \beta = \cos (a + \beta) + \cos (a - \beta)$   
 $= \cos \sigma + \cos (a - \beta),$

and is therefore a maximum when  $a - \beta = 0$ , or  $a = \beta = \frac{\sigma}{2}$ .

Thus the maximum value of  $\cos a \cos \beta$  is  $\cos^2 \frac{\sigma}{2}$ .

$$\begin{aligned}\text{Again,} \quad \cos a + \cos \beta &= 2 \cos \frac{a+\beta}{2} \cos \frac{a-\beta}{2} \\ &= 2 \cos \frac{\sigma}{2} \cos \frac{a-\beta}{2},\end{aligned}$$

and is therefore a maximum when  $a = \beta = \frac{\sigma}{2}$ .

Thus the maximum value of  $\cos a + \cos \beta$  is  $2 \cos \frac{\sigma}{2}$ .

Similar theorems hold in case of the sine.

*Example 1.* If  $A, B, C$  are the angles of a triangle, find the maximum value of

$$\sin A + \sin B + \sin C \text{ and of } \sin A \sin B \sin C.$$

Let us suppose that  $C$  remains constant, while  $A$  and  $B$  vary.

$$\begin{aligned}\sin A + \sin B + \sin C &= 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2} + \sin C \\ &= 2 \cos \frac{C}{2} \cos \frac{A-B}{2} + \sin C.\end{aligned}$$

This expression is a maximum when  $A = B$ .

Hence, so long as any two of the angles  $A, B, C$  are unequal, the expression  $\sin A + \sin B + \sin C$  is not a maximum; that is, the expression is a maximum when  $A = B = C = 60^\circ$ .

$$\text{Thus the maximum value} = 3 \sin 60^\circ = \frac{3\sqrt{3}}{2}.$$

Again,

$$\begin{aligned}2 \sin A \sin B \sin C &= \{\cos(A-B) - \cos(A+B)\} \sin C \\ &= \{\cos(A-B) + \cos C\} \sin C.\end{aligned}$$

This expression is a maximum when  $A = B$ .

Hence, by reasoning as before,  $\sin A \sin B \sin C$  has its maximum value when  $A = B = C = 60^\circ$ .

$$\text{Thus the maximum value} = \sin^3 60^\circ = \frac{3\sqrt{3}}{8}.$$

*Example 2.* If  $\alpha$  and  $\beta$  are two angles, each lying between 0 and  $\frac{\pi}{2}$ , whose sum is constant, find the minimum value of  $\sec \alpha + \sec \beta$ .

$$\begin{aligned} \text{We have } \sec \alpha + \sec \beta &= \frac{1}{\cos \alpha} + \frac{1}{\cos \beta} = \frac{\cos \alpha + \cos \beta}{\cos \alpha \cos \beta} \\ &= \frac{4 \cos \frac{\alpha+\beta}{2} \cos \frac{\alpha-\beta}{2}}{\cos (\alpha+\beta) + \cos (\alpha-\beta)} = \frac{2 \cos \frac{\alpha+\beta}{2} \cos \frac{\alpha-\beta}{2}}{\cos^2 \frac{\alpha-\beta}{2} - \sin^2 \frac{\alpha+\beta}{2}} \\ &= \cos \frac{\alpha+\beta}{2} \left( \frac{1}{\cos \frac{\alpha-\beta}{2} + \sin \frac{\alpha+\beta}{2}} + \frac{1}{\cos \frac{\alpha-\beta}{2} - \sin \frac{\alpha+\beta}{2}} \right). \end{aligned}$$

Since  $\alpha + \beta$  is constant, this expression is least when the denominators are greatest; that is, when  $\alpha = \beta = \frac{\alpha+\beta}{2}$ .

Thus the minimum value is  $2 \sec \frac{\alpha+\beta}{2}$ .

**320.** If  $\alpha, \beta, \gamma, \delta, \dots$  are  $n$  angles, each lying between 0 and  $\frac{\pi}{2}$ , whose sum is constant, to find the maximum value of

$$\cos \alpha \cos \beta \cos \gamma \cos \delta \dots$$

$$\text{Let } \alpha + \beta + \gamma + \delta + \dots = \sigma.$$

Suppose that any two of the angles, say  $\alpha$  and  $\beta$ , are unequal; then if in the given product we replace the two unequal factors  $\cos \alpha$  and  $\cos \beta$  by the two equal factors  $\cos \frac{\alpha+\beta}{2}$  and  $\cos \frac{\alpha+\beta}{2}$ , the value of the product is increased while the sum of the angles remains unaltered. Hence so long as any two of the angles  $\alpha, \beta, \gamma, \delta, \dots$  are unequal the product is not a maximum; that is, the product is a maximum when all the angles are equal. In this case each angle  $= \frac{\sigma}{n}$ .

Thus the maximum value is  $\cos^n \frac{\sigma}{n}$ .

In like manner we may shew that

the maximum value of  $\cos \alpha + \cos \beta + \cos \gamma + \dots = n \cos \frac{\sigma}{n}$ .

321. The methods of solution used in the following examples are worthy of notice.

*Example 1.* Shew that  $\tan 3a \cot a$  cannot lie between 3 and  $\frac{1}{3}$ .

We have  $\tan 3a \cot a = \frac{\tan 3a}{\tan a} = \frac{3 - \tan^2 a}{1 - 3 \tan^2 a} = n$  say;

$$\therefore \tan^2 a = \frac{n-3}{3n-1} = \frac{3-n}{1-3n}.$$

These two fractional values of  $\tan^2 a$  must be positive, and therefore  $n$  must be greater than 3 or less than  $\frac{1}{3}$ .

*Example 2.* If  $a$  and  $b$  are positive quantities, of which  $a$  is the greater, find the minimum value of  $a \sec \theta - b \tan \theta$ .

Denote the expression by  $x$ , and put  $\tan \theta = t$ ;  
then

$$x = a \sqrt{1+t^2} - bt;$$

$$\therefore b^2 t^2 + 2bxt + x^2 = a^2 (1+t^2);$$

$$\therefore t^2 (b^2 - a^2) + 2bxt + x^2 - a^2 = 0.$$

In order that the values of  $t$  found from this equation may be real,

$$b^2 x^2 > (b^2 - a^2) (x^2 - a^2);$$

$$\therefore 0 > a^2 (a^2 - b^2 - x^2);$$

$$\therefore x^2 > a^2 - b^2.$$

Thus the minimum value is  $\sqrt{a^2 - b^2}$ .

*Example 3.* If  $a, b, c, k$  are constant quantities and  $\alpha, \beta, \gamma$  variable quantities subject to the relation  $a \tan \alpha + b \tan \beta + c \tan \gamma = k$ , find the minimum value of  $\tan^2 \alpha + \tan^2 \beta + \tan^2 \gamma$ .

By multiplying out and re-arranging the terms, we have

$$\begin{aligned} (a^2 + b^2 + c^2) (\tan^2 \alpha + \tan^2 \beta + \tan^2 \gamma) - (a \tan \alpha + b \tan \beta + c \tan \gamma)^2 \\ = (b \tan \gamma - c \tan \beta)^2 + (c \tan \alpha - a \tan \gamma)^2 + (a \tan \beta - b \tan \alpha)^2. \end{aligned}$$

But the minimum value of the right side of this equation is zero; hence the minimum value of

$$(a^2 + b^2 + c^2) (\tan^2 \alpha + \tan^2 \beta + \tan^2 \gamma) - k^2 = 0;$$

that is, the minimum value of

$$\tan^2 \alpha + \tan^2 \beta + \tan^2 \gamma = \frac{k^2}{a^2 + b^2 + c^2}.$$

**EXAMPLES. XXV. a.**

When  $\theta$  is variable find the minimum value of the following expressions :

1.  $p \cot \theta + q \tan \theta.$
2.  $4 \sin^2 \theta + \operatorname{cosec}^2 \theta.$
3.  $8 \sec^2 \theta + 18 \cos^2 \theta.$
4.  $3 - 2 \cos \theta + \cos^2 \theta.$

Prove the following inequalities :

5.  $\tan^2 \alpha + \tan^2 \beta + \tan^2 \gamma > \tan \beta \tan \gamma$   
 $\quad \quad \quad + \tan \gamma \tan \alpha + \tan \alpha \tan \beta.$
6.  $\sin^2 \alpha + \sin^2 \beta > 2 (\sin \alpha + \sin \beta - 1).$

When  $\theta$  is variable, find the maximum value of

7.  $\sin \theta + \cos \theta.$
8.  $\cos \theta + \sqrt{3} \sin \theta.$
9.  $a \cos (a + \theta) + b \sin \theta.$
10.  $p \cos \theta + q \sin (a + \theta).$

If  $\sigma = \alpha + \beta$ , where  $\alpha$  and  $\beta$  are two angles each lying between 0 and  $\frac{\pi}{2}$ , and  $\sigma$  is constant, find the maximum or minimum value of

11.  $\sin \alpha + \sin \beta.$
12.  $\sin \alpha \sin \beta.$
13.  $\tan \alpha + \tan \beta.$
14.  $\operatorname{cosec} \alpha + \operatorname{cosec} \beta.$

If  $A, B, C$  are the angles of a triangle, find the maximum or minimum value of

15.  $\cos A \cos B \cos C.$
16.  $\cot A + \cot B + \cot C.$
17.  $\sin^2 \frac{A}{2} + \sin^2 \frac{B}{2} + \sin^2 \frac{C}{2}.$
18.  $\sec A + \sec B + \sec C.$
19.  $\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2}.$   $\left[ \text{Use } \Sigma \tan \frac{B}{2} \tan \frac{C}{2} = 1. \right]$
20.  $\cot^2 A + \cot^2 B + \cot^2 C.$   $\left[ \text{Use } \Sigma \cot B \cot C = 1. \right]$

21. If  $b^2 < 4ac$ , find the maximum and minimum values of  
 $a \sin^2 \theta + b \sin \theta \cos \theta + c \cos^2 \theta.$



22. If  $\alpha, \beta, \gamma$  lie between 0 and  $\frac{\pi}{2}$ , shew that

$$\sin \alpha + \sin \beta + \sin \gamma > \sin (\alpha + \beta + \gamma).$$

23. If  $a$  and  $b$  are two positive quantities of which  $a$  is the greater, shew that  $a \operatorname{cosec} \theta > b \cot \theta + \sqrt{a^2 - b^2}$ .

24. Shew that  $\frac{\sec^2 \theta - \tan \theta}{\sec^2 \theta + \tan \theta}$  lies between 3 and  $\frac{1}{3}$ .

25. Find the maximum value of  $\frac{\tan^2 \theta - \cot^2 \theta + 1}{\tan^2 \theta + \cot^2 \theta - 1}$ .

26. If  $\alpha, b, c, k$  are constant positive quantities, and  $\alpha, \beta, \gamma$  variable quantities subject to the relation

$$a \cos \alpha + b \cos \beta + c \cos \gamma = k,$$

find the minimum value of

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma \text{ and of } a \cos^2 \alpha + b \cos^2 \beta + c \cos^2 \gamma.$$

### Elimination.

322. No general rules can be given for the elimination of some assigned quantity or quantities from two or more trigonometrical equations. The form of the equations will often suggest special methods, and in addition to the usual algebraical artifices we shall always have at our disposal the identical relations subsisting between the trigonometrical functions. Thus suppose it is required to eliminate  $\theta$  from the equations

$$x \cos \theta = a, \quad y \cot \theta = b.$$

Here  $\sec \theta = \frac{x}{a}$ , and  $\tan \theta = \frac{y}{b}$ ;

but for all values of  $\theta$ , we have

$$\sec^2 \theta - \tan^2 \theta = 1.$$

$\therefore$  by substitution,

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1.$$

From this example we see that since  $\theta$  satisfies *two* equations (either of which is sufficient to determine  $\theta$ ) there is a relation, independent of  $\theta$ , which subsists between the coefficients and

constants of the equations. To determine this relation we *eliminate*  $\theta$ , and the result is called the *eliminant* of the given equations.

**323.** The following examples will illustrate some useful methods of elimination.

*Example 1.* Eliminate  $\theta$  between the equations

$$l \cos \theta + m \sin \theta + n = 0 \text{ and } p \cos \theta + q \sin \theta + r = 0.$$

From the given equations, we have by cross multiplication

$$\frac{\cos \theta}{mr - nq} = \frac{\sin \theta}{np - lr} = \frac{1}{lq - mp};$$

$$\therefore \cos \theta = \frac{mr - nq}{lq - mp}, \text{ and } \sin \theta = \frac{np - lr}{lq - mp};$$

whence by squaring, adding, and clearing of fractions, we obtain

$$(mr - nq)^2 + (np - lr)^2 = (lq - mp)^2.$$

The particular instance in which  $q = l$  and  $p = -m$  is of frequent occurrence in Analytical Geometry. In this case the eliminant may be written down at once; for we have

$$l \cos \theta + m \sin \theta = -n,$$

and

$$l \sin \theta - m \cos \theta = -r;$$

whence by squaring and adding, we obtain

$$l^2 + m^2 = n^2 + r^2.$$

*Example 2.* Eliminate  $\theta$  between the equations

$$\frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = c^2 \text{ and } l \tan \theta = m.$$

From the second equation, we have

$$\frac{\sin \theta}{m} = \frac{\cos \theta}{l} = \frac{\sqrt{\sin^2 \theta + \cos^2 \theta}}{\sqrt{m^2 + l^2}} = \frac{1}{\sqrt{m^2 + l^2}};$$

$$\therefore \sin \theta = \frac{m}{\sqrt{m^2 + l^2}}, \text{ and } \cos \theta = \frac{l}{\sqrt{m^2 + l^2}}.$$

By substituting in the first equation, we obtain

$$\frac{ax}{l} - \frac{by}{m} = \frac{c^2}{\sqrt{m^2 + l^2}}.$$

*Example 3.* Eliminate  $\theta$  between the equations

$$x = \cot \theta + \tan \theta \text{ and } y = \sec \theta - \cos \theta.$$

From the given equations, we have

$$\begin{aligned} x &= \frac{1}{\tan \theta} + \tan \theta = \frac{1 + \tan^2 \theta}{\tan \theta} \\ &= \frac{\sec^2 \theta}{\tan \theta}, \end{aligned}$$

and

$$\begin{aligned} y &= \sec \theta - \frac{1}{\sec \theta} = \frac{\sec^2 \theta - 1}{\sec \theta} \\ &= \frac{\tan^2 \theta}{\sec \theta}. \end{aligned}$$

From these values of  $x$  and  $y$  we obtain

$$x^2 y = \sec^3 \theta \text{ and } xy^2 = \tan^3 \theta.$$

But

$$\sec^2 \theta - \tan^2 \theta = 1;$$

$$\therefore (x^2 y)^{\frac{2}{3}} - (xy^2)^{\frac{2}{3}} = 1;$$

that is,

$$x^{\frac{4}{3}} y^{\frac{2}{3}} - x^{\frac{2}{3}} y^{\frac{4}{3}} = 1.$$

*Example 4.* Eliminate  $\theta$  from the equations

$$\frac{x}{a} = \cos \theta + \cos 2\theta \text{ and } \frac{y}{b} = \sin \theta + \sin 2\theta.$$

From the given equations, we have

$$\frac{x}{a} = 2 \cos \frac{3\theta}{2} \cos \frac{\theta}{2},$$

and

$$\frac{y}{b} = 2 \sin \frac{3\theta}{2} \cos \frac{\theta}{2};$$

whence by squaring and adding, we obtain

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 4 \cos^2 \frac{\theta}{2}.$$

But

$$\frac{x}{a} = 2 \cos \frac{\theta}{2} \left( 4 \cos^3 \frac{\theta}{2} - 3 \cos \frac{\theta}{2} \right)$$

$$= 2 \cos^2 \frac{\theta}{2} \left( 4 \cos^2 \frac{\theta}{2} - 3 \right);$$

$$\therefore \frac{2x}{a} = \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} \right) \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} - 3 \right).$$

**324.** The following examples are instances of the elimination of two quantities.

*Example 1.* Eliminate  $\theta$  and  $\phi$  from the equations

$$a \sin^2 \theta + b \cos^2 \theta = m, \quad b \sin^2 \phi + a \cos^2 \phi = n, \quad a \tan \theta = b \tan \phi.$$

From the first equation, we have

$$a \sin^2 \theta + b \cos^2 \theta = m (\sin^2 \theta + \cos^2 \theta);$$

$$\therefore (a - m) \sin^2 \theta = (m - b) \cos^2 \theta;$$

$$\therefore \tan^2 \theta = \frac{m - b}{a - m}.$$

From the second equation, we have

$$b \sin^2 \phi + a \cos^2 \phi = n (\sin^2 \phi + \cos^2 \phi);$$

$$\therefore \tan^2 \phi = \frac{n - a}{b - n}.$$

From the third equation,

$$a^2 \tan^2 \theta = b^2 \tan^2 \phi;$$

$$\therefore \frac{a^2 (m - b)}{a - m} = \frac{b^2 (n - a)}{b - n};$$

$$\therefore a^2 (bm - b^2 - mn + bn) = b^2 (an - a^2 - mn + am);$$

$$\therefore mab(a - b) + nab(a - b) = mn(a^2 - b^2);$$

$$\therefore mab + nab = mn(a + b);$$

$$\therefore \frac{1}{n} + \frac{1}{m} = \frac{1}{a} + \frac{1}{b}.$$

*Example 2.* Eliminate  $\theta$  and  $\phi$  from the equations

$$x \cos \theta + y \sin \theta = x \cos \phi + y \sin \phi = 2a, \quad 2 \sin \frac{\theta}{2} \sin \frac{\phi}{2} = 1.$$

From the data, we see that  $\theta$  and  $\phi$  are the roots of the equation

$$x \cos \alpha + y \sin \alpha = 2a;$$

$$\therefore (x \cos \alpha - 2a)^2 = y^2 \sin^2 \alpha = y^2 (1 - \cos^2 \alpha);$$

$$\therefore (x^2 + y^2) \cos^2 \alpha - 4ax \cos \alpha + 4a^2 - y^2 = 0,$$

which is a quadratic in  $\cos \alpha$  with roots  $\cos \theta$  and  $\cos \phi$ .

But  $1 = 4 \sin^2 \frac{\theta}{2} \sin^2 \frac{\phi}{2} = (1 - \cos \theta)(1 - \cos \phi);$

whence

$$\cos \theta + \cos \phi = \cos \theta \cos \phi ;$$

$$\therefore \frac{4ax}{x^2 + y^2} = \frac{4a^2 - y^2}{x^2 + y^2} ;$$

$$\therefore y^2 = 4a(a - x).$$

**325.** The method exhibited in the following example is one frequently used in Analytical Geometry.

*Example.* If  $a, b, c$  are unequal, find the relations that hold between the coefficients, when

$$a \cos \theta + b \sin \theta = c,$$

and

$$a \cos^2 \theta + 2a \cos \theta \sin \theta + b \sin^2 \theta = c.$$

The required relation will be obtained by eliminating  $\theta$  from the given equations. This is most conveniently done by making each equation homogeneous in  $\sin \theta$  and  $\cos \theta$ .

From the first equation, we have

$$a \cos \theta + b \sin \theta = c \sqrt{\cos^2 \theta + \sin^2 \theta} ;$$

whence, by squaring and transposing,

$$(a^2 - c^2) \cos^2 \theta + 2ab \cos \theta \sin \theta + (b^2 - c^2) \sin^2 \theta = 0 \dots\dots(1).$$

From the second equation, we have

$$a \cos^2 \theta + 2a \cos \theta \sin \theta + b \sin^2 \theta = c (\cos^2 \theta + \sin^2 \theta) ;$$

$$\therefore (a - c) \cos^2 \theta + 2a \cos \theta \sin \theta + (b - c) \sin^2 \theta = 0 \dots\dots(2).$$

From (1) and (2) we have by cross-multiplication,

$$\begin{aligned} \frac{\cos^2 \theta}{2ab(b-c) - 2a(b^2 - c^2)} &= \frac{\cos \theta \sin \theta}{(b^2 - c^2)(a-c) - (a^2 - c^2)(b-c)} \\ &= \frac{\sin^2 \theta}{2a(a^2 - c^2) - 2ab(a-c)} ; \end{aligned}$$

$$\text{or} \quad \frac{\cos^2 \theta}{-2ac(b-c)} = \frac{\cos \theta \sin \theta}{(b-c)(a-c)(b-a)} = \frac{\sin^2 \theta}{2a(a-c)(a+c-b)} ;$$

$$\therefore -4a^2c(b-c)(a-c)(a+c-b) = (b-c)^2(a-c)^2(b-a)^2.$$

By supposition, the quantities  $a, b, c$  are unequal; hence dividing by  $(b-c)(a-c)$ , we obtain

$$4a^2c(a+c-b) + (b-c)(a-c)(a-b)^2 = 0.$$

**EXAMPLES. XXV. b.**

Eliminate  $\theta$  between the equations:

1.  $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1, \quad \frac{x}{a} \sin \theta - \frac{y}{b} \cos \theta = 1.$
2.  $a \sec \theta - x \tan \theta = y, \quad b \sec \theta + y \tan \theta = x.$
3.  $\cos \theta + \sin \theta = a, \quad \cos 2\theta = b,$
4.  $x = \sin \theta + \cos \theta, \quad y = \tan \theta + \cot \theta.$
5.  $a = \cot \theta + \cos \theta, \quad b = \cot \theta - \cos \theta.$

Find the eliminant in each of the following cases:

6.  $x = \cot \theta + \tan \theta, \quad y = \operatorname{cosec} \theta - \sin \theta.$
7.  $\operatorname{cosec} \theta - \sin \theta = a^3, \quad \sec \theta - \cos \theta = b^3.$
8.  $4x = 3a \cos \theta + a \cos 3\theta, \quad 4y = 3a \sin \theta - a \sin 3\theta.$
9.  $x = \tan^2 \theta (a \tan \theta - x), \quad y = \sec^2 \theta (y - a \sec \theta).$
10.  $x = a \cos \theta (2 \cos 2\theta - 1), \quad y = b \sin \theta (4 \cos^2 \theta - 1).$
11. If  $\cos(\theta - \alpha) = a$ , and  $\sin(\theta - \beta) = b$ ,  
shew that  $a^2 - 2ab \sin(\alpha - \beta) + b^2 = \cos^2(\alpha - \beta).$

Find the relation that must hold between  $x$  and  $y$  if

12.  $x + y = 3 - \cos 4\theta, \quad x - y = 4 \sin 2\theta.$
13.  $x = \sin \theta + \cos \theta \sin 2\theta, \quad y = \cos \theta + \sin \theta \sin 2\theta.$
14. If  $\sin \theta + \cos \theta = a$ , and  $\sin 2\theta + \cos 2\theta = b$ ,  
shew that  $(a^2 - b - 1)^2 = a^2 (2 - a^2).$
15. If  $\cos \theta - \sin \theta = b$ , and  $\cos 3\theta + \sin 3\theta = a$ ,  
shew that  $a = 3b - 2b^3.$
16. Eliminate  $\theta$  from the equations:  
 $a \cos \theta - b \sin \theta = c, \quad 2ab \cos 2\theta + (a^2 - b^2) \sin 2\theta = 2c^2.$
17. If  $x = a \cos \theta + b \cos 2\theta$ , and  $y = a \sin \theta + b \sin 2\theta$ ,  
shew that  $a^2 \{(x+b)^2 + y^2\} = (x^2 + y^2 - b^2)^2.$



18. If  $\frac{\tan(\theta + a)}{\tan(\theta - a)} = \frac{a + b}{a - b}$ , and  $a \cos 2a + b \cos 2\theta = c$ ,

shew that  $a^2 + c^2 - 2ac \cos 2a = b^2$ .

19. If  $x = a(\sin 3\theta - \sin \theta)$ , and  $y = a(\cos \theta - \cos 3\theta)$ ,

shew that  $(x^2 + y^2)(2a^2 - x^2 - y^2)^2 = 4a^4 x^2$ .

Eliminate  $\theta$  from the equations:

20.  $x \cos \theta - y \sin \theta = a \cos 2\theta$ ,  $x \sin \theta + y \cos \theta = 2a \sin 2\theta$ .

21.  $x \sin \theta - y \cos \theta = \sqrt{x^2 + y^2}$ ,  $\frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2} = \frac{1}{x^2 + y^2}$ .

22.  $\frac{x \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$ ,  $x \sin \theta - y \cos \theta = \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$ .

23. If  $\cos(a - 3\theta) = m \cos^3 \theta$ , and  $\sin(a - 3\theta) = m \sin^3 \theta$ ,

shew that  $m^2 + m \cos a = 2$ .

Eliminate  $\theta$  and  $\phi$  from the equations:

24.  $\tan \theta + \tan \phi = x$ ,  $\cot \theta + \cot \phi = y$ ,  $\theta + \phi = a$ .

25.  $\sin \theta + \sin \phi = a$ ,  $\cos \theta + \cos \phi = b$ ,  $\theta - \phi = a$ .

26.  $a \sin^2 \theta + b \cos^2 \theta = a \cos^2 \phi + b \sin^2 \phi = 1$ ,  $a \tan \theta = b \tan \phi$ .

27. If  $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = \frac{x}{a} \cos \phi + \frac{y}{b} \sin \phi = 1$ , and  $\theta - \phi = a$ ,

shew that  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = \sec^2 \frac{a}{2}$ .

28. If  $\tan \theta + \tan \phi = a$ ,  $\cot \theta + \cot \phi = b$ ,  $\theta - \phi = a$ ,

shew that  $ab(ab - 4) = (a + b)^2 \tan^2 a$ .

Eliminate  $\theta$  and  $\phi$  between the equations:

29.  $a \cos^2 \theta + b \sin^2 \theta = m \cos^2 \phi$ ,  $a \sin^2 \theta + b \cos^2 \theta = n \sin^2 \phi$ ,

$m \tan^2 \theta - n \tan^2 \phi = 0$ .

30.  $x \cos \theta + y \sin \theta = 2a \sqrt{3}$ ,  $x \cos(\theta + \phi) + y \sin(\theta + \phi) = 4a$ ,

$x \cos(\theta - \phi) + y \sin(\theta - \phi) = 2a$ .

31.  $c \sin \theta = a \sin(\theta + \phi)$ ,  $a \sin \phi = b \sin \theta$ ,  $\cos \theta - \cos \phi = 2m$ .

### Application of Trigonometry to the Theory of Equations.

**326.** In the Theory of Equations it is shewn that the solution of *any* cubic equation may be made to depend on the solution of a cubic equation of the form  $x^3 + ax + b = 0$ . In certain cases the solution is very conveniently obtained by Trigonometry.

**327.** Consider the equation

$$x^3 - qx - r = 0 \dots\dots\dots(1),$$

in which each of the letters  $q$  and  $r$  represents a positive quantity.

From the identity  $\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$ ,

we have 
$$\cos^3 \theta - \frac{3}{4} \cos \theta - \frac{\cos 3\theta}{4} = 0 \dots\dots\dots(2).$$

Let  $x = y \cos \theta$ , where  $y$  is a positive quantity; then from (1),

$$\cos^3 \theta - \frac{q}{y^2} \cos \theta - \frac{r}{y^3} = 0 \dots\dots\dots(3).$$

If the equations (2) and (3) are identical, we have  $\frac{q}{y^2} = \frac{3}{4}$ , so that

$y = + \sqrt{\frac{4q}{3}}$ , since  $y$  is positive; and

$$\frac{\cos 3\theta}{4} = \frac{r}{y^3} = \sqrt{\frac{27r^2}{64q^3}};$$

whence 
$$\cos 3\theta = \sqrt{\frac{27r^2}{4q^3}}.$$

Hence the values of  $\theta$  are real if  $27r^2 < 4q^3$ ;

that is, if 
$$\left(\frac{r}{2}\right)^2 < \left(\frac{q}{3}\right)^3.$$

Let  $a$  be the smallest angle whose cosine is equal to  $\sqrt{\frac{27r^2}{4q^3}}$ ; then  $\cos 3\theta = \cos a$ ; whence  $3\theta = 2n\pi \pm a$ .

Thus the values of  $\cos \theta$  are

$$\cos \frac{a}{3}, \quad \cos \frac{2\pi + a}{3}, \quad \cos \frac{2\pi - a}{3}. \quad [\text{See Art. 264.}]$$

But 
$$x = y \cos \theta = \sqrt{\frac{4q}{3}} \cos \theta,$$

and therefore the roots of  $x^3 - qx - r = 0$  are

$$\sqrt{\frac{4q}{3}} \cos a, \quad \sqrt{\frac{4q}{3}} \cos \frac{2\pi + a}{3}, \quad \sqrt{\frac{4q}{3}} \cos \frac{2\pi - a}{3}.$$

328. Following the method explained in the preceding article, we may use the identity

$$\sin^3 \theta - \frac{3}{4} \sin \theta + \frac{\sin 3\theta}{4} = 0$$

to obtain the solution of the equation

$$x^3 - qx + r = 0,$$

each of the quantities represented by  $q$  and  $r$  being positive.

*Example.* Solve the equation  $x^3 - 12x + 8 = 0$ .

We have 
$$\sin^3 \theta - \frac{3}{4} \sin \theta + \frac{\sin 3\theta}{4} = 0.$$

In the given equation put  $x = y \sin \theta$ , where  $y$  is positive; then

$$\sin^3 \theta - \frac{12}{y^3} \sin \theta + \frac{8}{y^3} = 0.$$

$$\therefore \frac{3}{4} = \frac{12}{y^3}; \text{ whence } y = 4;$$

and 
$$\frac{\sin 3\theta}{4} = \frac{8}{y^3} = \frac{1}{8}; \text{ whence } \sin 3\theta = \frac{1}{2}.$$

Suppose that  $\theta$  is estimated in sexagesimal measure; then

$$3\theta = n \cdot 180^\circ + (-1)^n 30^\circ.$$

By ascribing to  $n$  the values 0, 1, 2, 3, 4 we obtain

$$\theta = 10^\circ, \quad \theta = 50^\circ, \quad \theta = 130^\circ, \quad \theta = 170^\circ, \quad \theta = 250^\circ;$$

and by further ascribing to  $n$  the values 5, 6, 7, ... it will easily be seen that the values of  $\sin \theta$  are equal to some one of the three quantities

$$\sin 10^\circ, \quad \sin 50^\circ, \quad -\sin 70^\circ.$$

But  $x = y \sin \theta = 4 \sin \theta$ , and therefore the roots are

$$4 \sin 10^\circ, \quad 4 \sin 50^\circ, \quad -4 \sin 70^\circ.$$

### Application of the Theory of Equations to Trigonometry.

329. In the Theory of Equations it is shewn that the equation whose roots are  $a_1, a_2, a_3, \dots, a_n$  is

$$(x - a_1)(x - a_2)(x - a_3) \dots (x - a_n) = 0,$$

or  $x^n - S_1 x^{n-1} + S_2 x^{n-2} - S_3 x^{n-3} + \dots + (-1)^n S_n = 0,$

where  $S_1$  = sum of the roots;

$S_2$  = sum of the products of the roots taken two at a time;

$S_3$  = sum of the products of the roots taken three at a time;

.....

$S_n$  = product of the roots.

[See Hall and Knight's *Higher Algebra*, Art. 538 and Art. 539.]

*Example 1.* If  $\alpha, \beta, \gamma$  are the values of  $\theta$  which satisfy the equation

$$a \tan^3 \theta + (2a - x) \tan \theta + y = 0 \dots \dots \dots (1),$$

shew that (i) if  $\tan \alpha + \tan \beta = h$ , then  $ah^3 + (2a - x)h = y$ ;

(ii) if  $\tan \alpha \tan \beta = k$ , then  $y^2 + (2a - x)ak^2 = a^2k^3$ .

(i) From the theory of equations, we have from (1),

$$\tan \alpha + \tan \beta + \tan \gamma = 0;$$

$$\therefore h + \tan \gamma = 0, \text{ or } \tan \gamma = -h.$$

But

$$a \tan^3 \gamma + (2a - x) \tan \gamma + y = 0;$$

$$\therefore ah^3 + (2a - x)h - y = 0.$$

(ii) From the theory of equations, we have from (1),

$$\tan \alpha \tan \beta \tan \gamma = -\frac{y}{a};$$

$$\therefore k \tan \gamma = -\frac{y}{a}, \text{ or } \tan \gamma = -\frac{y}{ak}.$$

Substituting in  $a \tan^3 \gamma + (2a - x) \tan \gamma + y = 0$ , we have

$$-\frac{ay^3}{a^3k^3} - (2a - x)\frac{y}{ak} + y = 0;$$

$$\therefore y^2 + (2a - x)ak^2 - a^2k^3 = 0.$$

*Example 2.* Shew that

$$\cos^2 \alpha + \cos^2 \left( \frac{2\pi}{3} + \alpha \right) + \cos^2 \left( \frac{2\pi}{3} - \alpha \right) = \frac{3}{2}.$$

Suppose that  $\cos 3\theta = k$ ;

then  $4 \cos^3 \theta - 3 \cos \theta = \cos 3\theta = k$ ;

$$\therefore \cos^3 \theta - \frac{3}{4} \cos \theta - \frac{k}{4} = 0.$$

The roots of this cubic in  $\cos \theta$  are

$$\cos \alpha, \cos \left( \frac{2\pi}{3} + \alpha \right), \text{ and } \cos \left( \frac{2\pi}{3} - \alpha \right),$$

where  $\alpha$  is any angle which satisfies the equation  $\cos 3\alpha = k$ . For shortness, denote the roots by  $a, b, c$ ; then

$$\begin{aligned} a^2 + b^2 + c^2 &= (a + b + c)^2 - 2(bc + ca + ab) \\ &= 0 - 2 \left( -\frac{3}{4} \right); \end{aligned}$$

$$\therefore \cos^2 \alpha + \cos^2 \left( \frac{2\pi}{3} + \alpha \right) + \cos^2 \left( \frac{2\pi}{3} - \alpha \right) = \frac{3}{2}.$$

**330.** If  $5\theta = 2n\pi$ , where  $n$  is any integer, we have

$$3\theta = 2n\pi - 2\theta;$$

$$\therefore \sin 3\theta = -\sin 2\theta.$$

The values of  $\sin \theta$  found from this equation are

$$0, \sin \frac{2\pi}{5}, \sin \frac{4\pi}{5}, \sin \frac{6\pi}{5}, \sin \frac{8\pi}{5},$$

being obtained by giving to  $n$  the values 0, 1, 2, 3, 4. It will easily be seen that no new values of  $\sin \theta$  are obtained by ascribing to  $n$  the values 5, 6, 7, ....

But  $\sin \frac{6\pi}{5} = -\sin \frac{4\pi}{5} = -\sin \frac{\pi}{5},$

and  $\sin \frac{8\pi}{5} = -\sin \frac{2\pi}{5};$

hence rejecting the zero solution, the values of  $\sin \theta$  found from the equation  $\sin 3\theta = -\sin 2\theta$  are

$$\pm \sin \frac{\pi}{5}, \text{ and } \pm \sin \frac{2\pi}{5}.$$

If we put  $\sin \theta = x$ , the equation  $\sin 3\theta = -\sin 2\theta$  becomes

$$3x - 4x^3 = -2x\sqrt{1-x^2}.$$

Dividing by  $x$ , and thus removing the solution  $x=0$ , we have

$$(3-4x^2)^2 = 4(1-x^2),$$

or

$$16x^4 - 20x^2 + 5 = 0.$$

This is a quadratic in  $x^2$ , and as we have just seen the values of  $x^2$  are

$$\sin^2 \frac{\pi}{5} \text{ and } \sin^2 \frac{2\pi}{5}.$$

From the theory of quadratic equations, we have

$$\sin^2 \frac{\pi}{5} + \sin^2 \frac{2\pi}{5} = \frac{20}{16} = \frac{5}{4};$$

$$\sin^2 \frac{\pi}{5} \sin^2 \frac{2\pi}{5} = \frac{5}{16}.$$

*Example.* Shew that

$$\sin \frac{2\pi}{7} + \sin \frac{4\pi}{7} + \sin \frac{8\pi}{7} = \frac{1}{2}\sqrt{7}.$$

If  $7\theta = 2n\pi$ , where  $n$  is any integer, we have

$$\sin 4\theta = -\sin 3\theta.$$

The values of  $\sin \theta$  found from this equation are

$$0, \pm \sin \frac{2\pi}{7}, \pm \sin \frac{4\pi}{7}, \pm \sin \frac{8\pi}{7},$$

since

$$\sin \frac{6\pi}{7} = -\sin \frac{8\pi}{7}.$$

If  $\sin \theta = x$ , the equation  $\sin 4\theta = -\sin 3\theta$  becomes

$$4x(1-2x^2)\sqrt{1-x^2} = 4x^3 - 3x;$$

whence rejecting the solution  $x=0$ , we obtain

$$16(1-4x^2+4x^4)(1-x^2) = 16x^4 - 24x^2 + 9,$$

or

$$64x^6 - 112x^4 + 56x^2 - 7 = 0 \dots\dots\dots(1).$$

The values of  $x^2$  found from this equation are

$$\sin^2 \frac{2\pi}{7}, \sin^2 \frac{4\pi}{7}, \sin^2 \frac{8\pi}{7};$$



hence 
$$\sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7} = \frac{112}{64} = \frac{7}{4}.$$

But 
$$\sin \frac{2\pi}{7} \sin \frac{4\pi}{7} + \sin \frac{2\pi}{7} \sin \frac{8\pi}{7} + \sin \frac{4\pi}{7} \sin \frac{8\pi}{7}$$

$$= \frac{1}{2} \left\{ \left( \cos \frac{2\pi}{7} - \cos \frac{6\pi}{7} \right) + \left( \cos \frac{6\pi}{7} - \cos \frac{10\pi}{7} \right) + \left( \cos \frac{4\pi}{7} - \cos \frac{12\pi}{7} \right) \right\}$$

$$= 0.$$

$$\therefore \left( \sin \frac{2\pi}{7} + \sin \frac{4\pi}{7} + \sin \frac{8\pi}{7} \right)^2 = \sin^2 \frac{2\pi}{7} + \sin^2 \frac{4\pi}{7} + \sin^2 \frac{8\pi}{7} = \frac{7}{4};$$

$$\therefore \sin \frac{2\pi}{7} + \sin \frac{4\pi}{7} + \sin \frac{8\pi}{7} = \frac{1}{2} \sqrt{7}.$$

331. If  $7\theta = 2n\pi$ , where  $n$  is any integer, we have

$$4\theta = 2n\pi - 3\theta;$$

$$\therefore \cos 4\theta = \cos 3\theta.$$

By giving to  $n$  the values 0, 1, 2, 3, the values of  $\cos \theta$  obtained from this equation are

$$1, \cos \frac{2\pi}{7}, \cos \frac{4\pi}{7}, \cos \frac{6\pi}{7}.$$

It will easily be seen that no new values of  $\cos \theta$  are found by ascribing to  $n$  the values 4, 5, 6, 7, .....; for

$$\cos \frac{8\pi}{7} = \cos \frac{6\pi}{7}, \cos \frac{10\pi}{7} = \cos \frac{4\pi}{7}, \dots\dots\dots$$

Now 
$$\cos 4\theta = 8 \cos^4 \theta - 8 \cos^2 \theta + 1,$$

and therefore if  $x = \cos \theta$ , the equation  $\cos 4\theta = \cos 3\theta$  becomes

$$8x^4 - 8x^2 + 1 = 4x^3 - 3x,$$

or 
$$8x^4 - 4x^3 - 8x^2 + 3x + 1 = 0.$$

Removing the factor  $x - 1$ , which corresponds to the root  $\cos \theta = 1$ , we obtain

$$8x^3 + 4x^2 - 4x - 1 = 0,$$

the roots of which equation are

$$\cos \frac{2\pi}{7}, \cos \frac{4\pi}{7}, \cos \frac{6\pi}{7}.$$

*Example 1.* Find the values of

$$\tan^2 \frac{\pi}{7} + \tan^2 \frac{2\pi}{7} + \tan^2 \frac{3\pi}{7} \text{ and } \tan \frac{\pi}{7} \tan \frac{2\pi}{7} \tan \frac{3\pi}{7}.$$

If  $7\theta = n\pi$ , where  $n$  is any integer, we have

$$\tan 4\theta = -\tan 3\theta.$$

By writing  $\tan \theta = t$ , this equation becomes

$$\frac{4t - 4t^3}{1 - 6t^2 + t^4} = -\frac{3t - t^3}{1 - 3t^2},$$

or

$$t^6 - 21t^4 + 35t^2 - 7 = 0.$$

The roots of this cubic in  $t^2$  are

$$\tan^2 \frac{\pi}{7}, \quad \tan^2 \frac{2\pi}{7}, \quad \tan^2 \frac{3\pi}{7}.$$

$$\therefore \tan^2 \frac{\pi}{7} + \tan^2 \frac{2\pi}{7} + \tan^2 \frac{3\pi}{7} = 21,$$

and

$$\tan \frac{\pi}{7} \tan \frac{2\pi}{7} \tan \frac{3\pi}{7} = \sqrt{7},$$

the positive value of the square root being taken, since each of the factors on the left is positive.

*Example 2.* Shew that

$$\cos^4 \frac{\pi}{7} + \cos^4 \frac{2\pi}{7} + \cos^4 \frac{3\pi}{7} = \frac{13}{16};$$

and

$$\sec^4 \frac{\pi}{7} + \sec^4 \frac{2\pi}{7} + \sec^4 \frac{3\pi}{7} = 416.$$

Let  $y$  denote any one of the quantities

$$\cos^2 \frac{\pi}{7}, \quad \cos^2 \frac{2\pi}{7}, \quad \cos^2 \frac{3\pi}{7};$$

then  $2y = 1 + x$ , where  $x$  denotes one of the quantities

$$\cos \frac{2\pi}{7}, \quad \cos \frac{4\pi}{7}, \quad \cos \frac{6\pi}{7}.$$

From Art. 331, the equation whose roots are

$$\cos \frac{2\pi}{7}, \quad \cos \frac{4\pi}{7}, \quad \cos \frac{6\pi}{7}$$

is

$$8x^3 + 4x^2 - 4x - 1 = 0;$$

whence by substituting  $x=2y-1$ , it follows that

$$\cos^2 \frac{\pi}{7}, \cos^2 \frac{2\pi}{7}, \cos^2 \frac{3\pi}{7}$$

are the roots of the equation

$$8(2y-1)^3 + 4(2y-1)^2 - 4(2y-1) - 1 = 0,$$

or 
$$64y^3 - 80y^2 + 24y - 1 = 0.$$

$$\therefore \cos^2 \frac{\pi}{7} + \cos^2 \frac{2\pi}{7} + \cos^2 \frac{3\pi}{7} = \frac{80}{64} = \frac{5}{4};$$

and 
$$\Sigma \cos^2 \frac{\pi}{7} \cos^2 \frac{2\pi}{7} = \frac{24}{64} = \frac{3}{8}.$$

By squaring the first of these equations and subtracting twice the second equation, we have

$$\cos^4 \frac{\pi}{7} + \cos^4 \frac{2\pi}{7} + \cos^4 \frac{3\pi}{7} = \frac{13}{16}.$$

By putting  $z = \frac{1}{y}$ , we see that

$$\sec^2 \frac{\pi}{7}, \sec^2 \frac{2\pi}{7}, \sec^2 \frac{3\pi}{7}$$

are the roots of the equation

$$z^3 - 24z^2 + 80z - 64 = 0;$$

$$\therefore \sec^4 \frac{\pi}{7} + \sec^4 \frac{2\pi}{7} + \sec^4 \frac{3\pi}{7} = (24)^2 - (2 \times 80) = 416.$$

**332.** To find  $\cos 5\theta$  and  $\sin 5\theta$ , we may proceed as follows:

$$\begin{aligned} \cos 5\theta + \cos \theta &= 2 \cos 3\theta \cos 2\theta \\ &= (4 \cos^3 \theta - 3 \cos \theta) (4 \cos^2 \theta - 2); \end{aligned}$$

$$\therefore \cos 5\theta = 16 \cos^5 \theta - 20 \cos^3 \theta + 5 \cos \theta.$$

$$\begin{aligned} \sin 5\theta + \sin \theta &= 2 \sin 3\theta \cos 2\theta \\ &= (3 \sin \theta - 4 \sin^3 \theta) (2 - 4 \sin^2 \theta); \end{aligned}$$

$$\therefore \sin 5\theta = 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta.$$

It is easy to prove that

$$\cos 6\theta = 32 \cos^6 \theta - 48 \cos^4 \theta + 18 \cos^2 \theta - 1,$$

and 
$$\sin 6\theta = \cos \theta (32 \sin^5 \theta - 32 \sin^3 \theta + 6 \sin \theta).$$

## EXAMPLES. XXV. c.

Solve the following equations :

1.  $x^3 - 3x - 1 = 0.$

2.  $x^3 - 3x + 1 = 0.$

3.  $x^3 - 3x - \sqrt{3} = 0.$

4.  $8x^3 - 6x + \sqrt{2} = 0.$

5.  $8a^3x^3 - 6ax + 2 \sin 3A = 0.$

6.  $x^3 - 3a^2x - 2a^3 \cos 3A = 0.$

7. If  $\sin a$  and  $\sin \beta$  are the roots of the equation

$$a \sin^2 \theta + b \sin \theta + c = 0,$$

shew that (1) if  $\sin a + 2 \sin \beta = 1$ , then  $a^2 + 2b^2 + 3ab + ac = 0$ ,

(2) if  $c \sin a = a \sin \beta$ , then  $a + c = \pm b$ .

8. If  $\tan a$  and  $\tan \beta$  are the roots of the equation

$$a \tan^2 \theta - b \tan \theta + c = 0, \text{ and if } a \tan a + b \tan \beta = 2b,$$

shew that

$$b^2(2a - b) + c(a - b)^2 = 0.$$

9. If  $\tan a$ ,  $\tan \beta$ ,  $\tan \gamma$  are the roots of the equation

$$a \tan^3 \theta + (2a - x) \tan \theta + y = 0,$$

and if  $a(\tan^2 a + \tan^2 \beta) = 2x - 5a$ , shew that  $x \pm y = 3a$ .

10. If  $\cos a$ ,  $\cos \beta$ ,  $\cos \gamma$  are the roots of the equation

$$\cos^3 \theta + a \cos^2 \theta + b \cos \theta + c = 0,$$

and if  $\cos a(\cos \beta + \cos \gamma) = 2b$ , prove that  $abc + 2b^3 + c^2 = 0$ .

Prove the following identities :

$$11. \sec a + \sec \left( \frac{2\pi}{3} + a \right) + \sec \left( \frac{2\pi}{3} - a \right) = -3 \sec 3a.$$

$$12. \sin^2 a + \sin^2 \left( \frac{2\pi}{3} + a \right) + \sin^2 \left( \frac{4\pi}{3} + a \right) = \frac{3}{2}.$$

$$13. \operatorname{cosec} a + \operatorname{cosec} \left( \frac{2\pi}{3} + a \right) + \operatorname{cosec} \left( \frac{4\pi}{3} + a \right) = 3 \operatorname{cosec} 3a.$$

$$14. \operatorname{cosec}^2 \frac{\pi}{5} + \operatorname{cosec}^2 \frac{2\pi}{5} = 4.$$

$$15. \cos \frac{2\pi}{5} + \cos \frac{4\pi}{5} = -\frac{1}{2}, \text{ and } \cos \frac{2\pi}{5} \cos \frac{4\pi}{5} = -\frac{1}{4}.$$

16. Form the equation whose roots are

$$(1) \cos \frac{\pi}{7}, \cos \frac{3\pi}{7}, \cos \frac{5\pi}{7};$$

$$(2) \sin^2 \frac{\pi}{14}, \sin^2 \frac{3\pi}{14}, \sin^2 \frac{5\pi}{14}.$$

17. Form the equation whose roots are

$$\sin^2 \frac{\pi}{7}, \sin^2 \frac{2\pi}{7}, \sin^2 \frac{3\pi}{7};$$

and shew that  $\sum_{n=1}^{n=3} \sin^4 \frac{n\pi}{7} = \frac{21}{16}$  and  $\sum_{n=1}^{n=3} \operatorname{cosec}^4 \frac{n\pi}{7} = 32$ .

18. Form the equation whose roots are

$$(1) \cos \frac{2\pi}{9}, \cos \frac{4\pi}{9}, \cos \frac{6\pi}{9}, \cos \frac{8\pi}{9};$$

$$(2) \cos \frac{\pi}{9}, \cos \frac{3\pi}{9}, \cos \frac{5\pi}{9}, \cos \frac{7\pi}{9}.$$

19. Form the equation whose roots are

$$\cos^2 \frac{\pi}{9}, \cos^2 \frac{2\pi}{9}, \cos^2 \frac{3\pi}{9}, \cos^2 \frac{4\pi}{9},$$

and shew that  $\sum_{n=1}^{n=4} \cos^4 \frac{n\pi}{9} = \frac{19}{16}$ , and  $\sum_{n=1}^{n=4} \sec^4 \frac{n\pi}{9} = 1120$ .

20. Form the equation whose roots are

$$\tan^2 \frac{\pi}{9}, \tan^2 \frac{2\pi}{9}, \tan^2 \frac{3\pi}{9}, \tan^2 \frac{4\pi}{9},$$

and shew that  $\cot^2 \frac{\pi}{9} + \cot^2 \frac{2\pi}{9} + \cot^2 \frac{4\pi}{9} = 9$ .

21. Form the equation whose roots are

$$\cos \frac{\pi}{11}, \cos \frac{2\pi}{11}, \cos \frac{3\pi}{11}, \cos \frac{4\pi}{11}, \cos \frac{5\pi}{11};$$

and shew that  $\sum_{n=1}^{n=5} \sec^2 \frac{n\pi}{11} = 60$ , and  $\prod_{n=1}^{n=5} \sec \frac{n\pi}{11} = 32$ .

## MISCELLANEOUS EXAMPLES. I.

1. If  $a \tan \alpha + b \tan \beta = (a+b) \tan \frac{\alpha+\beta}{2}$ ,  
 prove that  $a \cos \beta = b \cos \alpha$ .

2. If  $\frac{\sin^4 \alpha}{a} + \frac{\cos^4 \alpha}{b} = \frac{1}{a+b}$ ,  
 prove that  $\frac{\sin^8 \alpha}{a^3} + \frac{\cos^8 \alpha}{b^3} = \frac{1}{(a+b)^3}$ .

3. Shew that

$$2 \tan^{-1} \left\{ \tan \frac{\alpha}{2} \tan \left( \frac{\pi}{4} - \frac{\beta}{2} \right) \right\} = \tan^{-1} \left( \frac{\sin \alpha \cos \beta}{\cos \alpha + \sin \beta} \right).$$

4. If the equation

$$\frac{\sin^{2n+2} \theta}{\sin^{2n} \alpha} + \frac{\cos^{2n+2} \theta}{\cos^{2n} \alpha} = 1$$

is true when  $n=1$ , prove that it will be true when  $n$  is any positive integer.

5. If  $a \cos \theta + b \sin \theta = c$  and  $a \cos^2 \theta + b \sin^2 \theta = c$ ,  
 prove that  $4a^2b^2 + (b-c)(a-c)(a-b)^2 = 0$ .

6. Prove the following identities :

$$(i) \quad \Sigma \sin (\beta - \gamma) \cos (\alpha - \beta) \cos (\alpha - \gamma) = -\Pi \sin (\beta - \gamma);$$

$$(ii) \quad \Sigma \sin \alpha \sin (\beta - \gamma) \cos (\beta + \gamma - \alpha) = 0;$$

$$(iii) \quad \Sigma \sin \alpha \sin (\beta - \gamma) \sin (\beta + \gamma - \alpha) = 2\Pi \sin (\beta - \gamma).$$

7. If  $P$  be a point within a triangle  $ABC$ , such that

$$\angle PAB = \angle PBC = \angle PCA = \omega,$$

prove that (1)  $\cot \omega = \cot A + \cot B + \cot C$ ;

$$(2) \quad \operatorname{cosec}^2 \omega = \operatorname{cosec}^2 A + \operatorname{cosec}^2 B + \operatorname{cosec}^2 C.$$

8. A hill of inclination 1 in 169 faces West. Shew that a railway on it which runs S.E. has an inclination of 1 in 239.

9. Two vertical walls of equal height  $a$  are inclined to one another at an angle  $\alpha$ . At noon the breadth of their shadows are  $b$  and  $c$  : shew that the altitude  $\theta$  of the sun is given by the equation

$$a^2 \sin^2 \gamma \cot^2 \theta = b^2 + c^2 + 2bc \cos \gamma.$$



# ANSWERS.

## I. PAGE 4.

- |                    |                   |                    |                 |
|--------------------|-------------------|--------------------|-----------------|
| 1. .75.            | 2. .125.          | 3. .375.           | 4. .0241.       |
| 5. .089.           | 6. .0204045.      | 7. 76° 91' 66.7".  | 8. 21° 12' 50". |
| 9. 56° 24' 25".    | 10. 48° 75' 25".  | 11. 12° 23' 40.7". |                 |
| 12. 158° 6' 94.4". | 13. 22' 50".      | 14. 6' 36.7".      |                 |
| 15. 51° 11' 15".   | 16. 35° 9' 22.5". | 17. 36° 0' 40.6".  |                 |
| 18. 55' 5.8".      | 19. 2° 43' 6.4".  | 20. 7° 17' 26.1".  |                 |
| 21. 3' 22.5".      | 22. 20' 0.4".     | 23. 45°, 27°.      | 24. 72°.        |

## II. PAGE 11.

- |   |  |
|---|--|
| 1. $\frac{15}{17}, \frac{17}{8}, \frac{8}{15}, \frac{17}{15}.$  | 2. $\frac{12}{5}, \frac{13}{12}, \frac{5}{13}, \frac{12}{13}.$ |
| 3. $25, \frac{4}{5}, \frac{4}{5}, \frac{3}{4}, \frac{5}{3}.$  | 4. $7, \frac{7}{25}, \frac{24}{7}, \frac{25}{24}.$             |
| 5. $\frac{37}{35}, \frac{37}{12}, \frac{35}{12}, \frac{35}{37}.$  | 6. 12 inches, $\frac{4}{5}, \frac{3}{5}, \frac{4}{3}.$         |
| 7. $25, \frac{24}{25}, \frac{7}{25}.$   | 8. 40 ft., $\frac{40}{41}, \frac{9}{40}.$                      |
| 9. 20 ft., $\text{sine} = \frac{20}{29}, \text{cosine} = \frac{21}{29}, \text{tangent} = \frac{20}{21}.$  |  |
| 10. $\text{sine} = \frac{1}{\sqrt{5}}, \text{cosine} = \frac{2}{\sqrt{5}}, \text{tangent} = \frac{1}{2}.$ |  |
| 11. $\frac{12}{13}, \frac{13}{12}, \frac{77}{85}, \frac{85}{77}.$   | 12. $\frac{3}{5}, \frac{4}{3}, \frac{20}{29}, \frac{29}{20}.$  |

## III. c. PAGE 23.

- |                                    |                                |  |                                    |
|------------------------------------|--------------------------------|--|------------------------------------|
| 1. $\frac{2}{\sqrt{3}}, \sqrt{3}.$ | 2. $\frac{4}{5}, \frac{3}{5}.$ | 3. $\frac{1}{\sqrt{15}}, \frac{\sqrt{15}}{4}.$ | 4. $\frac{\sqrt{5}}{2}, \sqrt{5}.$ |
|------------------------------------|--------------------------------|--|------------------------------------|

5.  $\frac{\sqrt{48}}{7}, \frac{1}{\sqrt{48}}$ . 6.  $\frac{7}{24}, \frac{25}{24}$ . 7.  $\sqrt{1 - \cos^2 A}, \frac{\sqrt{1 - \cos^2 A}}{\cos A}$ .
8.  $\sqrt{1 + \cot^2 \alpha}, \frac{\cot \alpha}{\sqrt{1 + \cot^2 \alpha}}$ . 9.  $\frac{\sqrt{\sec^2 \theta - 1}}{\sec \theta}, \frac{1}{\sqrt{\sec^2 \theta - 1}}$ .
10.  $\operatorname{cosec} A = \frac{1}{\sin A}, \cos A = \sqrt{1 - \sin^2 A}, \sec A = \frac{1}{\sqrt{1 - \sin^2 A}},$   
 $\tan A = \frac{\sin A}{\sqrt{1 - \sin^2 A}}, \cot A = \frac{\sqrt{1 - \sin^2 A}}{\sin A}$ . 11.  $\sqrt{2}$ .
13.  $\frac{p}{q}$ . 14.  $\frac{m^2 - 1}{2m}, \frac{m^2 - 1}{m^2 + 1}$ . 15.  $\frac{p^2 - q^2}{p^2 + q^2}, \frac{p^2 + q^2}{2pq}$ .
16. 3. 17.  $\frac{p^2 - q^2}{p^2 + q^2}$ .

## IV. a. PAGE 26.

1. 5. 2.  $1\frac{1}{2}$ . 3. 0. 4.  $2\frac{1}{3}$ . 5.  $\frac{1}{2}$ .
6.  $1\frac{1}{2}$ . 7. 9. 8. 2. 9.  $2\frac{1}{12}$ . 10.  $\frac{1}{2}$ .
11.  $\frac{3}{4}$ . 12. 0. 13.  $1\frac{11}{12}$ . 14.  $\frac{\sqrt{3}}{2}$ . 15. 6.

## IV. b. PAGE 28.

1.  $22^\circ 30'$ . 2.  $64^\circ 59' 30''$ . 3.  $79^\circ 58' 57''$ . 4.  $45^\circ + A$ .
5.  $45^\circ - B$ . 6.  $60^\circ + B$ . 7.  $50^\circ$ . 8.  $60^\circ$ .
9.  $18^\circ$ . 10.  $9^\circ$ . 11.  $22^\circ 30'$ . 12.  $45^\circ$ .
13.  $30^\circ$ . 14.  $15^\circ$ . 30. 1. 31.  $\tan A$ .

## IV. c. PAGE 31.

1.  $45^\circ$ . 2.  $60^\circ$ . 3.  $60^\circ$ . 4.  $45^\circ$ . 5.  $60^\circ$ .
6.  $30^\circ$ . 7.  $45^\circ$ . 8.  $60^\circ$ . 9.  $45^\circ$ . 10.  $60^\circ$ .
11.  $45^\circ$ . 12.  $60^\circ$ . 13.  $45^\circ$ . 14.  $30^\circ$ . 15.  $45^\circ$ .
16.  $60^\circ$ . 17.  $30^\circ$ . 18.  $60^\circ$ . 19.  $45^\circ$ . 20.  $60^\circ$ .
21.  $30^\circ$ . 22.  $30^\circ$ . 23.  $60^\circ$ . 24.  $45^\circ$ .
25.  $45^\circ$  or  $30^\circ$ .  $[(2 \sin \theta - 1)(\tan \theta - 1) = 0.]$  26.  $60^\circ$ . 28. 1 or  $\frac{1}{2}$ .

## MISCELLANEOUS EXAMPLES. A. PAGE 32.

1. (1)  $\cdot 2537064$ ; (2)  $\cdot 704$ .      3.  $\frac{20}{29}, \frac{29}{21}$ .      4.  $\frac{15}{8}, \frac{17}{8}$ .
6. (1)  $15^\circ 28' 7\cdot5''$ ; (2)  $1' 37\cdot2''$ .      7.  $41, \frac{9}{40}, \frac{41}{9}, \frac{41}{40}$ .
8. (1) possible; (2) impossible; (3) possible.
10.  $\frac{\sqrt{1+\cot^2 a}}{\cot a}, \sqrt{1+\cot^2 a}$ .      11. 6.
12.  $\frac{m}{\sqrt{m^2+n^2}}, \frac{\sqrt{m^2+n^2}}{n}$ .      16.  $\frac{20}{21}, \frac{29}{20}$ .
18.  $10^\circ$ .      20. (1)  $30^\circ$ ; (2)  $45^\circ$ .      22.  $30^\circ$ .
26.  $\frac{5}{14}$ .      29. (1)  $30^\circ$ ; (2)  $30^\circ$ .

## V. a. PAGE 37.

1.  $c=2, B=60^\circ, C=30^\circ$ .      2.  $a=6\sqrt{3}, A=60^\circ, C=30^\circ$ .
3.  $c=8\sqrt{3}, A=30^\circ, B=60^\circ$ .      4.  $c=30\sqrt{3}, B=30^\circ, C=60^\circ$ .
5.  $b=20\sqrt{2}, A=C=45^\circ$ .      6.  $c=10\sqrt{3}, A=30^\circ, B=60^\circ$ .
7.  $a=2\sqrt{2}, B=C=45^\circ$ .      8.  $a=9, A=60^\circ, C=30^\circ$ .
9.  $B=60^\circ, b=27, c=18\sqrt{3}$ .      10.  $C=60^\circ, b=2, c=2\sqrt{3}$ .
11.  $B=30^\circ, a=4\sqrt{3}, b=4$ .      12.  $B=90^\circ, a=3\sqrt{3}, c=3$ .
13.  $A=30^\circ, a=50, c=50\sqrt{3}$ .      14.  $C=90^\circ, a=20, c=40$ .
15.  $A=90^\circ, a=4\sqrt{2}, b=4$ .      16.  $A=90^\circ, b=4, c=4\sqrt{3}$ .
17. 700.      18. 31.      19.  $86\cdot47$ .      20.  $97\cdot8$ .
21.  $C=54^\circ, a=73, b=124$ .      22.  $B=68^\circ 17', C=21^\circ 43', b=93$ .
23.  $C=50^\circ 36', a=34\cdot3875, c=30\cdot435$ .
24.  $c=353, A=39^\circ 36', B=50^\circ 24'$ .

## V. b. PAGE 39.

1.  $10\sqrt{3}$ .      2.  $a=10\sqrt{2}, c=20$ .
3.  $AB=10\sqrt{3}$  ft.,  $AC=10$  ft.,  $AD=5\sqrt{3}$  ft.      4. 12, 4.
5.  $24\sqrt{3}$ .      6.  $20(\sqrt{3}-1)$ .
7.  $20(3+\sqrt{3})$ .      8.  $DC=BD=100$ .

## VI. a. PAGE 42.

- |                        |                      |               |                    |
|------------------------|----------------------|---------------|--------------------|
| 1. 173·2 ft.           | 2. 277·12 ft.        | 3. 60°.       | 4. 50 ft.; 100 ft. |
| 5. 22·5 ft.; 38·97 ft. | 6. 30 ft.            | 7. 200 yds.   |                    |
| 8. 51 yds., 81 yds.    | 9. 86·6 yds.         | 10. 46·19 ft. |                    |
| 11. 273·2 ft.          | 12. Each = 70·98 ft. | 13. 5 miles.  |                    |
| 14. 73·2 ft.           | 15. 64 ft.           | 16. 300 ft.   |                    |
| 17. 1193 yds.          | 18. 277·12 yds.      |               |                    |

## VI. b. PAGE 47.

- |                               |                                     |
|-------------------------------|-------------------------------------|
| 1. 565·6 yds.; 1131·2 yds.    | 2. 3·464 miles; 6 miles.            |
| 3. 29 miles.                  | 4. 10 miles per hour.               |
| 5. 10 miles; 24·14 miles.     | 6. 16 miles; S. 25° W.              |
| 7. 9·656 miles.               | 8. 5·77 miles; 11·54 miles.         |
| 9. 295·1 knots.               | 10. 5·196 miles per hour; 18 miles. |
| 11. 31 minutes past midnight. | 12. 38·97 miles per hour.           |

## VII. a. PAGE 54.

- |                         |                        |                        |                      |                       |
|-------------------------|------------------------|------------------------|----------------------|-----------------------|
| 1. $\frac{\pi}{4}$ .    | 2. $\frac{\pi}{6}$ .   | 3. $\frac{7\pi}{12}$ . | 4. $\frac{\pi}{8}$ . | 5. $\frac{\pi}{10}$ . |
| 6. $\frac{23\pi}{72}$ . | 7. $\frac{2\pi}{25}$ . | 8. $\frac{7\pi}{16}$ . | 9. 4509.             | 10. 6545.             |
| 11. 1·4399.             | 12. 1·1999.            | 13. 2·7489.            | 14. 9163.            |                       |
| 15. 135°.               | 16. 28°.               | 17. 33° 20'.           | 18. 37° 30'.         |                       |
| 19. 22° 30'.            | 20. 30°.               | 21. 37° 30'.           | 22. 165°.            |                       |
| 23. 638.                | 24. 1·332.             | 25. 2·0262.            | 26. 2·9979.          |                       |

## VII. b. PAGE 56.

- |                    |                            |  |  |       |
|--------------------|----------------------------|--|--|-------|
| 1. $\frac{3}{4}$ . | 2. $\frac{1}{3\sqrt{2}}$ . | 3. $\frac{11}{4}$ .                    | 4. $\frac{3}{\sqrt{2}}$ .              | 5. 9. |
| 6. $\frac{3}{4}$ . | 7. 1.                      | 13. $\frac{\pi}{4}, \frac{2\pi}{15}$ . | 14. $\frac{5\pi}{6}, \frac{5\pi}{7}$ . |       |

## VII. c. PAGE 60.

- |                       |            |                                |
|-----------------------|------------|--------------------------------|
| 1. $\frac{1}{5}$ .    | 2. 300 ft. | 3. A radian.                   |
| 4. 5·85 yards.        | 5. 330.    | 6. $\frac{1}{44}$ of a second. |
| 7. 58 $\frac{3}{4}$ . | 8. 40 yds. | 9. 1·15192 miles.              |
| 10. 3·581.            | 11. 2° 6'. | 12. 45 feet.                   |

**MISCELLANEOUS EXAMPLES. B. PAGE 61.**

- |   |   |  |                   |
|---|---|--|-------------------|
| 1. $9^\circ$ .  | 2. $95.26$ .                              | 3. $54^\circ$ .                        | 4. $3438$ inches. |
| 6. $30^\circ$ .                                       | 8. $22\frac{1}{2}^\circ, \frac{\pi}{8}$ . | 9. $67\frac{1}{2}^\circ$ .             |                   |
| 10. $a = 6\sqrt{3}, c = 12$ , perp. $= 3\sqrt{3}$ .   |   | 12. $17.32$ ft.                        |                   |
| 14. $120^\circ, 36^\circ, 24^\circ$ .                 | 15. $-\frac{35}{8}$ .                     |  |                   |
| 17. (1) possible; (2) impossible.                     |   | 18. $8.66$ miles.                      |                   |
| 19. $\frac{\pi}{5}, \frac{\pi}{3}, \frac{7\pi}{15}$ . | 21. $90$ .                                | 24. $4$ miles per hour, $1.732$ miles. |                   |
| 25. $\frac{\pi}{8}$ .                                 | 26. (1) $30^\circ$ ; (2) $30^\circ$ .     | 27. $\frac{5}{13}$ .                   |                   |
| 29. $200$ yards.                                      | 30. $33$ feet.                            |  |                   |

**VIII. a. PAGE 69.**

- |                            |                                      |                                      |                     |
|----------------------------|--------------------------------------|--------------------------------------|---------------------|
| 1. Second.                 | 2. Third.                            | 3. First.                            | 4. Third.           |
| 5. Second.                 | 6. Second.                           | 7. Third.                            | 8. Third.           |
| 9. Sine.                   | 10. Cosine.                          | 11. Tangent.                         | 12. Sine.           |
| 13. Sine.                  | 14. Tangent.                         | 15. Sine.                            | 16. All.            |
| 17. Cosine.                | 18. $60^\circ, \frac{\sqrt{3}}{2}$ . | 19. $30^\circ, \frac{\sqrt{3}}{2}$ . | 20. $45^\circ, 1$ . |
| 21. $45^\circ, \sqrt{2}$ . | 22. $30^\circ, 2$ .                  | 23. $60^\circ, \frac{2}{\sqrt{3}}$ . | 24. $45^\circ, 1$ . |
| 25. $60^\circ, 2$ .        | 26. $60^\circ, \sqrt{3}$ .           |                                      |                     |

**VIII. b. PAGE 72.**

- |  |                                   |   |
|--|-----------------------------------|---|
| 1. $-\sqrt{3}$ .                               | 2. $\frac{1}{\sqrt{2}}$ .         | 3. $-\frac{1}{2}$ .                     |
| 4. $-\frac{12}{13}, \frac{12}{5}$ .            | 5. $-\frac{5}{4}, -\frac{3}{4}$ . | 6. $\frac{4}{3}, -\frac{3}{5}$ .        |
| 7. $\frac{\sqrt{3}}{2}, -\frac{1}{\sqrt{3}}$ . | 8. $1, -\sqrt{2}$ .               | 9. $\pm\frac{5}{13}, \pm\frac{5}{12}$ . |

**IX. PAGE 79.**

1.  $\cot A$  decreases from  $\infty$  to  $0$ , then increases numerically from  $0$  to  $-\infty$ , then decreases from  $\infty$  to  $0$ , then increases numerically from  $0$  to  $-\infty$ .      2.  $\operatorname{cosec} \theta$  decreases from  $\infty$  to  $1$ , then increases from  $1$  to  $\infty$ .      3.  $\cos \theta$  decreases numerically from  $-1$  to  $0$ , then increases from  $0$  to  $1$ .      4.  $\tan A$  decreases from  $\infty$  to  $0$ , then increases numerically from  $0$  to  $-\infty$ .      5.  $\sec \theta$  decreases numerically from  $-\infty$  to  $-1$ , then increases numerically from  $-1$  to  $-\infty$ .      6.  $3$ .      7.  $1$ .      8.  $-2$ .      9.  $2$ .

**MISCELLANEOUS EXAMPLES. C.** PAGE 81.

1.  $\pm \frac{4}{5}$ .      3.  $A=60^\circ, B=30^\circ, a=\frac{21\sqrt{3}}{2}$ .      4.  $\frac{7}{24}$ .
5. 1313 miles, nearly.      6. 301 feet.      7.  $-\frac{7}{10}$ .
8. 12.003 inches.      10. 200 feet.

**X. a.** PAGE 87.

1.  $-\frac{1}{\sqrt{2}}$ .      2.  $\frac{1}{2}$ .      3.  $\sqrt{3}$ .      4.  $-\sqrt{2}$ .
5.  $-\frac{\sqrt{3}}{2}$ .      6. 1.      7. -1.      8.  $-\frac{1}{2}$ .
9. 2.      10. -1.      11.  $-\frac{\sqrt{3}}{2}$ .      12. -2.
13. -2.      14.  $-\frac{1}{\sqrt{2}}$ .      15.  $\sqrt{3}$ .      16.  $\sin A$ .
17.  $\tan A$ .      18.  $-\cos A$ .      19.  $-\sec A$ .      20.  $-\cos A$ .
21.  $-\tan A$ .      22.  $-\cos \theta$ .      23.  $\tan \theta$ .      24.  $-\operatorname{cosec} \theta$ .
25. 1.      26.  $2 \sin A$ .      27. 1.

**X. b.** PAGE 91.

1.  $-\frac{1}{2}$ .      2.  $-\frac{\sqrt{3}}{2}$ .      3.  $\frac{1}{2}$ .      4.  $-\frac{1}{2}$ .      5. -1.
6. 1.      7.  $\frac{2}{\sqrt{3}}$ .      8.  $-\frac{1}{\sqrt{3}}$ .      9.  $-\sqrt{2}$ .      10.  $\frac{1}{\sqrt{2}}$ .
11. 0.      12.  $\frac{1}{\sqrt{2}}$ .      13.  $-\sqrt{2}$ .      14.  $-\frac{1}{2}$ .      15.  $-\sqrt{2}$ .
16.  $-\frac{1}{\sqrt{2}}$ .      17. -1.      18. 2.      19.  $\frac{1}{\sqrt{3}}$ .      20.  $\frac{2}{\sqrt{3}}$ .
21.  $\pm 30^\circ, \pm 330^\circ$ .      22.  $210^\circ, 330^\circ, -30^\circ, -150^\circ$ .
23.  $120^\circ, 300^\circ, -60^\circ, -240^\circ$ .      24.  $135^\circ, 315^\circ, -45^\circ, -225^\circ$ .
30. 3.      31.  $\cot^2 A$ .      32. -1.      34. -4.

**XI. a.** PAGE 97.

4.  $1; \frac{24}{25}$ .      5.  $\frac{33}{65}; \frac{16}{65}$ .      6.  $-\frac{85}{36}$ .



## XI. b. PAGE 100.

1. 1.            2.  $\frac{1}{7}$ .            3.  $0; \frac{12}{35}$ .            4.  $-\frac{278}{29}; \frac{1}{2}$ .
11.  $\cos A \cos B \cos C - \cos A \sin B \sin C - \sin A \cos B \sin C$   
 $- \sin A \sin B \cos C;$   
 $\sin A \cos B \cos C - \cos A \sin B \cos C + \cos A \cos B \sin C$   
 $+ \sin A \sin B \sin C.$
12.  $\frac{\tan A - \tan B - \tan C - \tan A \tan B \tan C}{1 - \tan B \tan C + \tan C \tan A + \tan A \tan B}.$
13.  $\frac{\cot A \cot B \cot C - \cot A - \cot B - \cot C}{\cot B \cot C + \cot C \cot A + \cot A \cot B - 1}.$

## XI. d. PAGE 104.

1.  $-\frac{7}{9}$ .            2.  $\frac{17}{25}$ .            3.  $\frac{24}{25}$ .            4.  $\frac{3}{4}$ .
5.  $\frac{7}{25}; \frac{24}{25}$ .            6.  $\frac{1}{3}$ .            7.  $\frac{1}{7}$ .

## XI. e. PAGE 106.

1.  $-\frac{23}{27}$ .            2.  $\frac{117}{125}$ .            3.  $\frac{9}{13}$ .

## XII. a. PAGE 112.

1.  $\sin 4\theta + \sin 2\theta$ .            2.  $\sin 9\theta - \sin 3\theta$ .            3.  $\cos 12A + \cos 2A$ .
4.  $\cos A - \cos 5A$ .            5.  $\sin 9\theta - \sin \theta$ .            6.  $\sin 12\theta - \sin 4\theta$ .
7.  $\cos 6\theta - \cos 12\theta$ .            8.  $\sin 16\theta - \sin 2\theta$ .            9.  $\cos 13\alpha + \cos 9\alpha$ .
10.  $\cos 5\alpha - \cos 15\alpha$ .            11.  $\frac{1}{2}(\sin 11\alpha - \sin 3\alpha)$ .
12.  $\frac{1}{2}(\cos 2\alpha - \cos 4\alpha)$ .            13.  $\frac{1}{2}(\sin 2A + \sin A)$ .
14.  $\frac{1}{2}(\sin 6A - \sin A)$ .            15.  $\cos \frac{7\theta}{3} + \cos \theta$ .
16.  $\frac{1}{2}\left(\cos \frac{\theta}{2} - \cos \theta\right)$ .            17.  $\cos(\alpha + \beta) + \cos(\alpha - 3\beta)$ .
18.  $\cos(2\alpha - \beta) - \cos(4\alpha + \beta)$ .            19.  $\sin(3\theta - \phi) + \sin(\theta + 3\phi)$ .
20.  $\sin(4\theta - \phi) - \sin(2\theta + 3\phi)$ .            21.  $\frac{1}{2}\left(\frac{\sqrt{3}}{2} - \sin 2\alpha\right)$ .

**XII. b. PAGE 114.**

1.  $2 \sin 6\theta \cos 2\theta$ .
2.  $2 \cos 3\theta \sin 2\theta$ .
3.  $2 \cos 5\theta \cos 2\theta$ .
4.  $2 \sin 10\theta \sin \theta$ .
5.  $2 \cos 6a \sin a$ .
6.  $2 \cos \frac{11a}{2} \cos \frac{5a}{2}$ .
7.  $2 \sin 8a \cos 5a$ .
8.  $-2 \sin 3a \sin 2a$ .
9.  $2 \cos \frac{11A}{2} \cos \frac{7A}{2}$ .
10.  $-2 \cos 7A \sin 4A$ .
11.  $-\sin 20^\circ$ .
12.  $\sqrt{3} \cos 10^\circ$ .

**XIII. a. PAGE 128.**

1.  $60^\circ$ .
2.  $120^\circ$ .
3.  $A=30^\circ, B=120^\circ, C=30^\circ$ .
4.  $45^\circ$ .
5.  $90^\circ$ .
6.  $A=75^\circ, B=45^\circ, C=60^\circ$ .
7.  $A=30^\circ, B=135^\circ, C=15^\circ$ .
8.  $28^\circ 56'$ .
9.  $111^\circ 33'$ .
10.  $\sqrt{6}$ .
11. 7.
12. 8.
13. 14.
14. 9.
15.  $b=2\sqrt{6}, A=75^\circ, C=30^\circ$ .
16.  $a=\sqrt{5}+1, B=36^\circ, C=72^\circ$ .
17.  $C=75^\circ, a=c=2\sqrt{3}+2$ .
18.  $A=105^\circ, a=\sqrt{3}+1, c=\sqrt{3}-1$ .
19.  $C=30^\circ, a=2, b=\sqrt{3}+1$ .
21. 2.
22. 6.
23.  $60^\circ$ .
24.  $105^\circ, 45^\circ, 30^\circ$ .
25.  $105^\circ, 15^\circ, 60^\circ$ .
26.  $\frac{\sqrt{3}}{2}, 105^\circ, 15^\circ$ .

**XIII. b. PAGE 132.**

1.  $B=60^\circ, 120^\circ; C=90^\circ, 30^\circ; c=2, 1$ .
2.  $B=60^\circ, 120^\circ; A=75^\circ, 15^\circ; a=3+\sqrt{3}, 3-\sqrt{3}$ .
3.  $A=45^\circ, B=75^\circ, b=\sqrt{3}+1$ ; no ambiguity.
4. Impossible.
5.  $C=45^\circ, 135^\circ; A=105^\circ, 15^\circ; a=3+\sqrt{3}, 3-\sqrt{3}$ .
6.  $C=75^\circ, 105^\circ; A=45^\circ, 15^\circ; a=2\sqrt{3}, 3-\sqrt{3}$ .
7.  $A=75^\circ, 105^\circ; B=90^\circ, 60^\circ; b=2\sqrt{6}, 3\sqrt{2}$ .
8.  $B=90^\circ, C=72^\circ, c=4\sqrt{5+2\sqrt{5}}$ ; no ambiguity.
9. Impossible.

**XIII. d. PAGE 136.**

1.  $72^\circ, 72^\circ, 36^\circ$ ; each side  $=\sqrt{5}+1$ .
2.  $A=60^\circ, a=9-3\sqrt{3}, b=3(\sqrt{6}-\sqrt{2}), c=3\sqrt{2}$ .
3.  $A=105^\circ, B=15^\circ, C=60^\circ$ .
4.  $B=54^\circ, 126^\circ; C=108^\circ, 36^\circ$ .
5.  $C=60^\circ, 120^\circ; A=90^\circ, 30^\circ; a=75\sqrt{3}$ . No, for  $C=90^\circ$ .
6.  $18^\circ, 126^\circ$ .
8.  $A=90^\circ, B=30^\circ, C=60^\circ; 2c=a\sqrt{3}$ .

**MISCELLANEOUS EXAMPLES. D. PAGE 138.**

2. 43.      3.  $\infty$ , 1.      4. -1.      6.  $a=2$ ,  $B=30^\circ$ ,  $C=105^\circ$ .  
 9.  $A=30^\circ$ ,  $B=75^\circ$ ,  $C=75^\circ$ .

**XIV. a. PAGE 145.**

1. 10, 8,  $-\frac{3}{2}$ ,  $\frac{2}{3}$ ,  $\frac{1}{2}$ , -2.      2.  $\frac{4}{3}$ ,  $\frac{5}{4}$ ,  $-\frac{1}{2}$ ,  $\frac{7}{4}$ .  
 3. 2401,  $\cdot 5$ ,  $\frac{10000}{9}$ , 1,  $\frac{5}{4}$ , 1000, 10000.  
 4. 5, 3, 3, 4, 0.      5. 0, 2,  $\bar{2}$ , 0,  $\bar{4}$ , 3,  $\bar{1}$ .  
 6.  $\bar{1}\cdot 8091488$ ,  $6\cdot 8091488$ ,  $\bar{4}\cdot 8091488$ .      7.  $3\cdot 25$ ,  $325$ ,  $\cdot 000325$ .  
 8.  $2\cdot 8853613$ .      9.  $3\cdot 3714373$ .      10.  $1\cdot 5475286$ .  
 11.  $1\cdot 9163822$ .      12.  $\bar{1}\cdot 4419030$ .      13.  $\bar{2}\cdot 2922560$ .  
 14.  $\bar{1}\cdot 6989700$ .      15.  $\bar{1}\cdot 8125918$ .      16.  $\cdot 0501716$ .  
 17.  $\log 2 = \cdot 3010300$ .      18.  $1 - \log 2 = \cdot 6989700$ .      19.  $1\cdot 320469$ .  
 20.  $\cdot 0260315$ .      21.  $\cdot 2898431$ .      22.  $\bar{7}\cdot 2621538$ .  
 23. 7; 4.      24. 2058.

**XIV. b. PAGE 149.**

1.  $9\cdot 076226$ .      2.  $3\cdot 01824$ .      3.  $2467\cdot 266$ .  
 4.  $2\cdot 23$ .      5.  $3\cdot 54$ .      6.  $1\cdot 72$ .      7.  $96$ ,  $79$ .  
 8.  $22\cdot 2398$ .      9.  $3\cdot 32$ .      10.  $5\cdot 77$ .      11.  $2\cdot 05$ .  
 12.  $x = 2 \log 2 = \cdot 60206$ ,  $y = -2 \log 5 = -1\cdot 39794$ .  
 13.  $x = \frac{\log 3}{\log 3 - \log 2} = 2\cdot 71$ ;  $y = \frac{\log 2}{\log 3 - \log 2} = x - 1 = 1\cdot 71$ .  
 14.  $3(b - a - c + 2)$ ,  $\frac{1}{2}(2a - 3c + 6)$ .  
 15.  $b + c - 2$ ,  $\frac{1}{6}(3a + 2b + 3c - 5)$ .

**MISCELLANEOUS EXAMPLES. E. PAGE 150.**

3.  $b = \sqrt{3} - 1$ ,  $A = 135^\circ$ ,  $C = 30^\circ$ .      8.  $A = 105^\circ$ ,  $B = 45^\circ$ .

**XV. a. PAGE 155.**

1.  $6\cdot 6947485$ .      2.  $\cdot 5404924$ .      3.  $6\cdot 4547860$ .  
 4.  $1\cdot 7606731$ .      5.  $6\cdot 7840083$ .      6.  $55740\cdot 83$ .  
 7.  $673\cdot 5469$ .      8.  $\cdot 0106867$ .      9.  $\cdot 008287771$ .  
 10.  $\cdot 2531925$ .      11.  $2\cdot 031324$ .      12.  $1\cdot 389495$ .  
 13.  $2\cdot 424463$ .      14.  $2\cdot 069138$ .

**XV. b. PAGE 159.**

- |                           |                          |                          |                          |
|---------------------------|--------------------------|--------------------------|--------------------------|
| 1. $\cdot 6164825$ .      | 2. $\cdot 7928863$ .     | 3. $1\cdot 2154838$ .    | 4. $62^\circ 42' 31''$ . |
| 5. $30^\circ 40' 23''$ .  | 6. $48^\circ 45' 44''$ . | 7. $9\cdot 8440554$ .    |                          |
| 8. $10\cdot 1317778$ .    | 9. $9\cdot 7530545$ .    | 10. $44^\circ 17' 8''$ . |                          |
| 11. $55^\circ 30' 39''$ . | 12. $9\cdot 6653952$ .   | 13. $10\cdot 1912872$ .  |                          |

**XV. c. PAGE 161.**

- |   |                          |                          |
|---|--------------------------|--------------------------|
| 1. $2\cdot 36952$ .                                     | 2. $84336$ .             | 3. $33\cdot 27475$ .     |
| 4. $\cdot 03803143$ .                                   | 5. $11218\cdot 4$ .      | 6. $1225\cdot 508$ .     |
| 7. $27\cdot 901983$ .                                   | 8. $\cdot 580303$ .      | 9. $6\cdot 848293$ .     |
| 10. $3\cdot 288754$ , $1\cdot 236122$ .                 |                          | 11. $2273\cdot 54$ .     |
| 12. $\cdot 5095328$ .                                   | 13. $7\cdot 29889$ .     | 14. $\cdot 045800373$ .  |
| 15. $\cdot 08603526$ .                                  | 16. $\cdot 0001706364$ . | 17. $\cdot 644065$ .     |
| 18. $9\cdot 52912$ .                                    | 19. $\cdot 3175272$ .    | 20. $\cdot 335859$ .     |
| 21. $\cdot 4221836$ .                                   | 22. $124272\cdot 2$ .    | 23. $250\cdot 2357$ .    |
| 24. (1) $36^\circ 45' 22''$ ; (2) $19^\circ 28' 16''$ . | 25. $\cdot 441785$ .     | 26. $68^\circ 25' 6''$ . |

**XVI. a. PAGE 166.**

- |                    |                    |
|--------------------|--------------------|
| 6. $\frac{1}{2}$ . | 7. $\frac{3}{2}$ . |
|--------------------|--------------------|

**XVI. b. PAGE 169.**

- |  |                          |                          |
|--|--------------------------|--------------------------|
| 1. $113^\circ 34' 41''$ .  | 2. $49^\circ 28' 26''$ . | 3. $55^\circ 46' 16''$ . |
| 4. $78^\circ 27' 47''$ .   | 5. $64^\circ 37' 23''$ . | 6. $35^\circ 5' 49''$ .  |
| 7. $93^\circ 35'$ .  |                          |                          |
| 8. $A=67^\circ 22' 49''$ , $B=53^\circ 7' 48''$ , $C=59^\circ 29' 23''$ .  |                          |                          |
| 9. $A=46^\circ 34' 3''$ , $B=104^\circ 28' 39''$ , $C=28^\circ 57' 18''$ . |                          |                          |

**XVI. c. PAGE 173.**

- |   |   |
|---|---|
| 1. $A=79^\circ 6' 24''$ , $B=40^\circ 53' 36''$ .                 | 2. $A=6^\circ 1' 54''$ , $B=108^\circ 58' 6''$ .  |
| 3. $A=24^\circ 10' 59''$ , $B=95^\circ 49' 1''$ .                 | 4. $B=78^\circ 48' 52''$ , $C=56^\circ 41' 8''$ . |
| 5. $A=27^\circ 38' 45''$ , $C=117^\circ 38' 45''$ .               | 6. $82^\circ 57' 15''$ , $36^\circ 32' 45''$ .    |
| 7. $A=74^\circ 32' 44''$ , $C=48^\circ 59' 16''$ .                |   |
| 8. $B=100^\circ 47' 1''$ , $C=14^\circ 12' 59''$ .                |   |
| 9. $A=136^\circ 35' 21\cdot 5''$ , $B=13^\circ 14' 33\cdot 5''$ . |   |

**XVI. d.** PAGE 174.

1.  $89.646162$ .      2.  $255.3864$ .      3.  $92.788$ .      4.  $b=185$ ,  $c=192$ .
5.  $321.0793$ .      6.  $a=765.4321$ ,  $c=1035.43$ .
7.  $b=767.72$ ,  $c=1263.58$ .

**XVI. e.** PAGE 176.

1.  $32^{\circ} 25' 35''$ .      2.  $41^{\circ} 41' 28''$  or  $138^{\circ} 18' 32''$ .
3.  $A=100^{\circ} 33'$ ,  $B=34^{\circ} 27'$ .      4.  $51^{\circ} 18' 21''$  or  $128^{\circ} 41' 39''$ .
5.  $A=28^{\circ} 20' 50''$ ,  $C=39^{\circ} 35' 10''$ .      6.  $A=81^{\circ} 45' 2''$ , or  $23^{\circ} 2' 58''$ .
7. (1) Not ambiguous, for  $C=90^{\circ}$ ;  
(2) ambiguous,  $b=60.3893$  ft.;  
(3) not ambiguous.

**XVI. f.** PAGE 180.

1.  $A=58^{\circ} 24' 43''$ ,  $B=48^{\circ} 11' 23''$ ,  $C=73^{\circ} 23' 54''$ .
2.  $112^{\circ} 12' 54''$ ,  $45^{\circ} 53' 33''$ ,  $21^{\circ} 53' 33''$ .      3.  $75^{\circ} 48' 54''$ .
4.  $4227.4718$ .      5.  $B=108^{\circ} 12' 26''$ ,  $C=49^{\circ} 27' 34''$ .
6.  $A=105^{\circ} 38' 57''$ ,  $B=15^{\circ} 38' 57''$ .      7.  $17.1$  or  $3.68$ .
8.  $108^{\circ} 26' 6''$ ,  $53^{\circ} 7' 48''$ ,  $18^{\circ} 26' 6''$ .      9.  $96^{\circ} 27'$ , or  $126^{\circ} 22'$ .
10.  $B=80^{\circ} 46' 26.5''$ ,  $C=63^{\circ} 48' 33.5''$ .      11.  $70^{\circ} 0' 56''$ , or  $109^{\circ} 59' 4''$ .
12.  $2.529823$ .      13.  $41^{\circ} 45' 14''$ .
14.  $A=42^{\circ} 0' 14''$ ,  $B=55^{\circ} 56' 46''$ ,  $C=82^{\circ} 3'$ .
15.  $41^{\circ} 24' 34''$ .      16.  $A=60^{\circ} 5' 34''$ ,  $C=29^{\circ} 54' 26''$ .
17.  $889.2554$  ft.      18.  $72^{\circ} 12' 59''$ ,  $47^{\circ} 47' 1''$ .
19.  $44.4878$  ft.      20.  $A=102^{\circ} 56' 38''$ ,  $B=42^{\circ} 3' 22''$ .
21.  $B=99^{\circ} 54' 23''$ ,  $C=32^{\circ} 50' 37''$ ,  $a=18.7254$ .      22.  $72^{\circ} 26' 26''$ .
23.  $A=27^{\circ} 29' 56''$ ,  $B=98^{\circ} 55'$ ,  $C=53^{\circ} 35' 4''$ .
24.  $B=32^{\circ} 15' 49''$ ,  $C=44^{\circ} 31' 17''$ ,  $a=1180.525$ .
25.  $a=20.9059$ ,  $c=33.5917$ .      26.  $a=2934.124$ ,  $b=3232.846$ .
27.  $B=1^{\circ} 1' 23''$ ,  $C=147^{\circ} 28' 37''$ ,  $a=4390$ .
28.  $A=26^{\circ} 24' 23''$ ,  $B=118^{\circ} 18' 25''$ ,  $b=642.756$ .
29.  $53^{\circ} 17' 55''$ , or  $126^{\circ} 42' 5''$ .
30.  $A=1^{\circ} 51' 56''$ ,  $C=125^{\circ} 49' 4''$ ,  $a=67$ .

31.  $b=4028\cdot5$ ,  $c=2831\cdot7$ .  
 32.  $B=75^\circ 53' 29''$ , or  $104^\circ 6' 31''$ ;  $A=60^\circ 54' 19''$ , or  $32^\circ 41' 17''$ .  
 33. Base= $2\cdot44845$  ft., altitude= $\cdot713322$  ft.  
 34.  $90^\circ$ .      35. (1) impossible; (2) ambiguous; (3) 64.  
 36.  $\theta=72^\circ 31' 53''$ ,  $c=12\cdot8255$ .      37.  $\theta=60^\circ 13' 52''$ ,  $c=19\cdot523977$ .

### XVII. a. PAGE 185.

1.  $146\cdot4$  ft.      2.  $880\sqrt{3}=1524$  ft.      5.  $ab/(a-b)$  ft.  
 6.  $\frac{1}{3}\sqrt{6}=\cdot816$  miles.      7.  $10(\sqrt{10}+\sqrt{2})=45\cdot76$  ft.  
 9.  $1$  or  $\frac{1}{3}$ .      10.  $9\frac{2}{7}$  ft.      12.  $48\sqrt{6}=117\cdot6$  ft.  
 14.  $750\sqrt{6}=1837$  ft.      15.  $2640(3+\sqrt{3})=12492$  ft.

### XVII. b. PAGE 190.

1.  $30$  ft.      2.  $a\sqrt{2}$  ft.      5.  $100$  ft.  
 12.  $\sqrt{500-200\sqrt{3}}=12\cdot4$  ft.

### XVII. c. PAGE 195.

1.  $1060\cdot5$  ft.      2.  $\frac{500\sqrt{6}}{3}=408$  ft.  
 3.  $120\sqrt{6}=294$  ft.      5.  $106$  ft.  
 10. Height= $40\sqrt{6}=98$  ft.; distance= $40(\sqrt{14}+\sqrt{2})=206$  ft.  
 11.  $50\sqrt{120+30\sqrt{6}}=696$  yds.

### XVII. d. PAGE 197.

1.  $5$  miles nearly.  
 2. Height= $19\cdot5375$  yds.; distance= $102\cdot9093$  yds.  
 3.  $200\cdot017$  ft.      4. Height= $418\cdot4045$  ft.; distance= $430$  ft.  
 5. Height= $916\cdot8624$  ft.; distance= $\cdot984808$  miles.  
 6. Height= $46\cdot140214$  ft.; distance= $99\cdot92$  ft.  
 7.  $11\cdot550316$  or  $25\cdot9733$  miles per hour.  
 8. Height= $159\cdot4321$  ft.; distance= $215\cdot6762$  ft.



**XVIII. a.** PAGE 206.

- |  |                            |                      |
|--|----------------------------|----------------------|
| 1. 9000 sq. ft.                          | 2. 15390.                  | 3. $\frac{84}{85}$ . |
| 4. $24, \frac{117}{5}, \frac{936}{25}$ . | 5. 225 sq. ft.             | 6. 672 sq. ft.       |
| 7. 36 yds.                               | 8. $r=4, R=8\frac{1}{3}$ . | 9. 12, 6, 28.        |
| 10. 12, 16, 20.                          |                            |                      |

**XVIII. b.** PAGE 210.

- |                   |                                   |
|-------------------|-----------------------------------|
| 1. 26.46 sq. ft.  | 2. 9.585 yds., 7.18875 sq. yds.   |
| 4. 216.23 sq. ft. | 5. 128.352 in.      6. 203.56 ft. |
| 7. 57.232 ft.     | 8. 61.803 sq. ft.                 |

**XVIII. c.** PAGE 218.

$$17. \quad \frac{\pi}{3} + (-1)^{n-1} \frac{1}{2^{n-1}} \left( A - \frac{\pi}{3} \right),$$

$$\frac{\pi}{3} + (-1)^{n-1} \frac{1}{2^{n-1}} \left( B - \frac{\pi}{3} \right),$$

$$\frac{\pi}{3} + (-1)^{n-1} \frac{1}{2^{n-1}} \left( C - \frac{\pi}{3} \right).$$

**XVIII. d.** PAGE 223.

- |                                   |                                 |
|-----------------------------------|---------------------------------|
| 1. $1\frac{5}{7}, 2\frac{1}{2}$ . | 4. Diagonals 65, 56; area 1764. |
|                                   | 5. $2\sqrt{77} + 6\sqrt{11}$ .  |

**XVIII. e.** PAGE 225.

- |                   |   |                 |
|-------------------|---|-----------------|
| 2. 14142 sq. yds. | 5. $\sqrt{\frac{x}{y} + \frac{y}{z} + \frac{z}{x}}$ . | 13. 20, 21, 29. |
|-------------------|---|-----------------|

**MISCELLANEOUS EXAMPLES. F.** PAGE 228.

- |  |
|--|
| 3. Expression = $\cot A + \cot B + \cot C$ .   |
| 4. $B=45^\circ, 135^\circ; C=105^\circ, 15^\circ; c=\sqrt{6}+\sqrt{2}, \sqrt{6}-\sqrt{2}$ .  |
| 6. 126.      7. 68.3 yds., 35.35 yds.  |
| 11. $C=45^\circ, 135^\circ; A=105^\circ, 15^\circ; a=2\sqrt{3}, 4\sqrt{3}-6$ .   |
| 12. $10(2-\sqrt{3})=2.68$ miles; $10(\sqrt{6}-\sqrt{2})=10.35$ miles.  |
| 24. (1) $90^\circ - \frac{A}{2}, 90^\circ - \frac{B}{2}, 90^\circ - \frac{C}{2}$ ;<br>(2) $180^\circ - 2A, 180^\circ - 2B, 180^\circ - 2C$ . |
| 25. Expression = $\sin^2(\alpha - \beta)$ .      28. 21.3 miles per hour.  |
| 29. 1 hr. 30'; 2 hrs. 16'.   |

**XIX. a.** PAGE 235.

1.  $n\pi + (-1)^n \frac{\pi}{6}$ .
2.  $n\pi + (-1)^n \frac{\pi}{4}$ .
3.  $2n\pi \pm \frac{\pi}{3}$ .
4.  $n\pi + \frac{\pi}{3}$ .
5.  $n\pi - \frac{\pi}{6}$ .
6.  $2n\pi \pm \frac{3\pi}{4}$ .
7.  $n\pi \pm \frac{\pi}{4}$ .
8.  $n\pi \pm \frac{\pi}{6}$ .
9.  $n\pi \pm \frac{\pi}{3}$ .
10.  $2n\pi \pm a$ .
11.  $n\pi \pm a$ .
12.  $n\pi \pm a$ .
13.  $n\pi$ .
14.  $\frac{n\pi}{3} + (-1)^n a$ .
15.  $2n\pi$ , or  $\frac{2n\pi}{5}$ .
16.  $\frac{n\pi}{3}$ , or  $n\pi \pm \frac{\pi}{6}$ .
17.  $\frac{n\pi}{4}$ , or  $\frac{n\pi}{3} + (-1)^n \frac{\pi}{18}$ .
18.  $\frac{(2n+1)\pi}{2}$ , or  $2n\pi$ , or  $\frac{(2n+1)\pi}{5}$ .
19.  $\frac{(2n+1)\pi}{2}$ , or  $\frac{(2n+1)\pi}{4}$ , or  $\frac{(2n+1)\pi}{8}$ .
20.  $n\pi$ , or  $\frac{(2n+1)\pi}{14}$ .
21.  $\frac{n\pi}{6}$ , or  $\frac{n\pi}{9}$ .
22.  $\frac{(2n+1)\pi}{6}$ , or  $n\pi \pm \frac{\pi}{8}$ .
23.  $\frac{(2n+1)\pi}{8}$ , or  $\frac{n\pi}{3} + (-1)^n \frac{\pi}{9}$ .
24.  $(2n+1)\pi$ , or  $2n\pi \pm \frac{\pi}{3}$ .
25.  $2n\pi$ , or  $2n\pi \pm \frac{2\pi}{3}$ .
26.  $n\pi + (-1)^n \frac{\pi}{6}$ , or  $2n\pi + \frac{3\pi}{2}$ .
27.  $\frac{n\pi}{2} + \frac{\pi}{8}$ .
28.  $2n\pi + \frac{2\pi}{3}$ .
29.  $2n\pi - \frac{\pi}{4}$ .

**XIX. b.** PAGE 237.

1.  $\frac{(2n+1)\pi}{2(p+q)}$ .
2.  $\frac{(4k+1)\pi}{2(n-m)}$ , or  $\frac{(4k-1)\pi}{2(n+m)}$ .
3.  $2n\pi$ , or  $2n\pi - \frac{2\pi}{3}$ .
4.  $2n\pi + \frac{\pi}{2}$ , or  $(2n+1)\pi + \frac{\pi}{6}$ .
5.  $2n\pi + \frac{\pi}{2}$ , or  $2n\pi + \frac{\pi}{6}$ .
6.  $2n\pi + \frac{5\pi}{12}$ , or  $2n\pi - \frac{\pi}{12}$ .
7.  $2n\pi + \frac{\pi}{12}$ , or  $2n\pi - \frac{7\pi}{12}$ .
8.  $2n\pi + \frac{5\pi}{4}$ , or  $2n\pi - \frac{3\pi}{4}$ .
9.  $2n\pi + \frac{\pi}{3}$ , or  $(2n+1)\pi$ .
10.  $\frac{n\pi}{2} + (-1)^n \frac{\pi}{12}$ .
11.  $\frac{(2n+1)\pi}{4}$ , or  $n\pi \pm \frac{\pi}{6}$ .

12.  $n\pi$ , or  $n\pi \pm \frac{\pi}{6}$ .

13.  $n\pi$ , or  $\frac{n\pi}{2}$ .

14.  $n\pi + \frac{\pi}{4}$ , or  $2n\pi$ , or  $2n\pi + \frac{\pi}{2}$ .

[In some of the following examples, the equations have to be squared, so that extraneous solutions are introduced.]

15.  $n\pi + \frac{\pi}{4}$ , or  $\frac{n\pi}{2} + (-1)^{n+1} \frac{\pi}{12}$ .

16.  $n\pi - \frac{\pi}{4}$ , or  $\frac{n\pi}{2} + (-1)^n \frac{\pi}{12}$ .

17.  $\frac{(2n+1)\pi}{10}$ , or  $\frac{(2n+1)\pi}{2}$ .

18.  $\frac{(2n+1)\pi}{2}$ , or  $n\pi \pm \frac{\pi}{3}$ .

19.  $n\pi + \frac{\pi}{4}$ , or  $n\pi + \frac{\pi}{6}$ .

20.  $\frac{n\pi}{2} + (-1)^{n+1} \frac{\pi}{12}$ , or  $\frac{n\pi}{2}$ .

21.  $\theta = n\pi \pm \frac{\pi}{4}$ ,  $\phi = n\pi \pm \frac{\pi}{6}$ .

22.  $\theta = n\pi \pm \frac{\pi}{6}$ ,  $\phi = n\pi \pm \frac{\pi}{3}$ .

23.  $\theta = n\pi \pm \frac{\pi}{4}$ ,  $\phi = n\pi \pm \frac{\pi}{3}$ .

## XIX. d. PAGE 244.

1.  $\pm \frac{1}{\sqrt{2}}$ .

2.  $\pm 1$ .

3.  $\pm 2$ .

4.  $\frac{-3 \pm \sqrt{17}}{4}$ .

5. 1, or  $\frac{1}{2}$ .

6. 0, or  $\pm \frac{1}{2}$ .

7.  $\pm \frac{1}{\sqrt{2}}$ .

8.  $\frac{25}{24}$ .

9.  $\frac{1}{2}$ .

10.  $\frac{a-b}{1+ab}$ .

11.  $\frac{b-a}{1+ab}$ .

12.  $\sqrt{3}$ .

13.  $x = ac - bd$ ,  $y = bc + ad$ .

14.  $\pm 1$ , or  $\pm (1 \pm \sqrt{2})$ .

15.  $n\pi$ , or  $n\pi + \frac{\pi}{4}$ .

19.  $x = 1$ ,  $y = 2$ ;  $x = 2$ ,  $y = 7$ .

## MISCELLANEOUS EXAMPLES. G. PAGE 246.

2. (1)  $\frac{(2n+1)\pi}{2}$ ,  $2n\pi \pm \frac{\pi}{3}$ ; (2)  $2n\pi \pm \frac{\pi}{3}$ .

6.  $78^\circ 27' 4''$ .

9. 6.

10. 800 yds, 146.4 yds, 546.4 yds.

## XX. a. PAGE 255.

2.  $\sin \frac{A}{2} = \frac{5}{13}$ ,  $\cos \frac{A}{2} = -\frac{12}{13}$ .

3.  $\sin \frac{A}{2} = -\frac{15}{17}$ ,  $\cos \frac{A}{2} = \frac{8}{17}$ .

4.  $2 \sin \frac{A}{2} = -\sqrt{1 + \sin A} + \sqrt{1 - \sin A}$ ;  
 $2 \cos \frac{A}{2} = -\sqrt{1 + \sin A} - \sqrt{1 - \sin A}$ .
5.  $2 \sin \frac{A}{2} = -\sqrt{1 + \sin A} - \sqrt{1 - \sin A}$ ;  
 $2 \cos \frac{A}{2} = -\sqrt{1 + \sin A} + \sqrt{1 - \sin A}$ .
6.  $2 \sin \frac{A}{2} = +\sqrt{1 + \sin A} + \sqrt{1 - \sin A}$ ;  
 $2 \cos \frac{A}{2} = +\sqrt{1 + \sin A} - \sqrt{1 - \sin A}$ .
7.  $\sin \frac{A}{2} = \frac{4}{5}$ ,  $\cos \frac{A}{2} = \frac{3}{5}$ .      8.  $\sin \frac{A}{2} = \frac{15}{17}$ ,  $\cos \frac{A}{2} = -\frac{8}{17}$ .
9. (1)  $2n\pi - \frac{\pi}{4}$  and  $2n\pi + \frac{\pi}{4}$ ; (2)  $2n\pi + \frac{5\pi}{4}$  and  $2n\pi + \frac{7\pi}{4}$ ;  
 (3)  $2n\pi + \frac{3\pi}{4}$  and  $2n\pi + \frac{5\pi}{4}$ .
10. No;  $2 \sin \frac{A}{2} = \sqrt{1 + \sin A} + \sqrt{1 - \sin A}$ .
14. (1)  $= \sqrt{2} \cos \left( \theta - \frac{\pi}{4} \right)$ ; (2)  $= 2 \sin \left( \theta - \frac{\pi}{3} \right)$ .
15. (1)  $= -\sec 2\theta$ ; (2)  $= \tan^2 \frac{\theta}{2}$ .

## XX. b. PAGE 260.

3.  $\frac{1}{5}$ .      4.  $\frac{1}{3}$ .

## XXI. a. PAGE 267.

1. 1440 yards.      2.  $342\frac{6}{7}$  yards.      3. 22 yards.  
 4.  $6' 34''$ .      5.  $13' 45''$ .      6. 11 ft. 11 in.  
 7. 210 yards.      8.  $9' 49''$ .      10. 50 ft.  
 11. (1)  $\frac{\pi}{10800}$ ; (2)  $\frac{\pi}{648000}$ .      12.  $\pi r^2$ .  
 13.  $\frac{1}{2}$ .      14.  $m - n$ .      15.  $\frac{1}{2} - \frac{\sqrt{3}}{200} = .491$ .  
 16.  $\frac{1}{2} + \frac{11\sqrt{3}}{7200} = .503$ .      17.  $\frac{21\sqrt{3}}{55} = 39.7'$ .

**XXI. b. PAGE 271.**

1. 12 miles.      2. 150 ft.      3. 15 miles.      4. 80 ft. 8 in.
5. 204 ft. 2 in.      6. 104 ft. 2 in.      7. 54' 33".
8. 10560 ft.      9. 610 ft.,  $\frac{5}{2}\sqrt{110}$  minutes = 26' 13".
11. 8.      12. -1.      13. (1)  $\cos \alpha$ ; (2)  $-\sin \alpha$ .
14.  $45^\circ 54' 33''$ ,  $44^\circ 5' 27''$ .

**MISCELLANEOUS EXAMPLES. H. PAGE 283.**

3.  $18^\circ 26' 6''$ .      6. 35 miles or 13 miles per hour.

**XXIII. a. PAGE 291.**

1.  $\frac{\sin 2n\alpha}{2 \sin \alpha}$ .      2.  $\sin \frac{n\beta}{2} \cos \left( \alpha - \frac{n-1}{2} \beta \right) / \sin \frac{\beta}{2}$ .
3.  $-\cos \left( \alpha + \frac{\pi}{2n} \right) / \sin \frac{\pi}{2n}$ .      4.  $\sin \frac{n\pi}{2k} \cos \frac{(n+1)\pi}{2k} / \sin \frac{\pi}{2k}$ .
5.  $\frac{1}{2}$ .      6.  $-\frac{1}{2}$ .      7.  $\cot \frac{\pi}{2n}$ .      8.  $-\cos \frac{2\pi}{n}$ .
9.  $\sin n\alpha$ .      10.  $\sin \frac{n(\theta + \pi)}{2} \sin \left( \frac{n+1}{2} \theta + \frac{n-1}{2} \pi \right) / \sin \frac{\theta + \pi}{2}$ .
11.  $\sin \frac{n(\pi - \beta)}{2} \cos \left\{ \alpha + \frac{(n-1)(\pi - \beta)}{2} \right\} / \sin \frac{\pi - \beta}{2}$ .
12.  $\sin \frac{n(\pi - 2\beta)}{4} \cos \left\{ \alpha + \frac{(n-1)(\pi - 2\beta)}{4} \right\} / \sin \frac{\pi - 2\beta}{4}$ .
13.  $\frac{n \cos \theta}{2} - \frac{\sin n\theta \cos (n+2)\theta}{2 \sin \theta}$ .
14.  $\frac{\sin 2n\alpha \sin 2(n+1)\alpha}{2 \sin 2\alpha} - \frac{n \sin 2\alpha}{2}$ .
15.  $\operatorname{cosec} \alpha \{ \tan (n+1)\alpha - \tan \alpha \}$ .
16.  $\operatorname{cosec} 2\theta \{ \cot \theta - \cot (2n+1)\theta \}$ .      17.  $\tan \alpha - \tan \frac{\alpha}{2^n}$ .
18.  $\frac{1}{2} (\operatorname{cosec} \alpha - \operatorname{cosec} 3^n \alpha)$ .      19.  $\frac{1}{2} (\tan 3^n \alpha - \tan \alpha)$ .

## XXIII. b. PAGE 294.

1.  $\frac{n}{2} + \frac{\sin 4n\theta}{4 \sin 2\theta}.$

2.  $\frac{n}{2}.$

3.  $\frac{n}{2}.$

4. 
$$\frac{3 \sin \frac{n\theta}{2} \sin \frac{(n+1)\theta}{2}}{4 \sin \frac{\theta}{2}} - \frac{\sin \frac{3n\theta}{2} \cos \frac{3(n+1)\theta}{2}}{4 \sin \frac{3\theta}{2}}.$$

5. 0.      6. 0.      7.  $\cot \theta - 2^n \cot 2^n \theta.$

8.  $\frac{1}{2} \operatorname{cosec} a \{ \tan (n+1) a - \tan a \}.$

9.  $\frac{\sin 2\theta}{2} - \frac{\sin 2^{n+1} \theta}{2^{n+1}}.$

10.  $\sin^2 \theta - 2^n \sin^2 \frac{\theta}{2^n}.$

11.  $\tan^{-1} x - \tan^{-1} \frac{x}{n+1}.$

12.  $\tan^{-1} (n+1) - \frac{\pi}{4}.$

13.  $\tan^{-1} \{ 1 + n(n+1) \} - \frac{\pi}{4}.$

14.  $\tan^{-1} n(n+1).$

## XXIV. a. PAGE 301.

2.  $x = a \cos \frac{\alpha + \beta}{2} \Big/ \cos \frac{\alpha - \beta}{2}, \quad y = b \sin \frac{\alpha + \beta}{2} \Big/ \cos \frac{\alpha - \beta}{2}.$

3.  $x = a (\cos \alpha + \sin \alpha), \quad y = b (\sin \alpha - \cos \alpha).$

13.  $4 \sin \frac{\alpha + \beta + \gamma}{2} \sin \frac{\beta + \gamma - \alpha}{2} \sin \frac{\gamma + \alpha - \beta}{2} \sin \frac{\alpha + \beta - \gamma}{2}.$

14.  $4 \sin \frac{\alpha + \beta + \gamma}{2} \sin \frac{\alpha + \beta - \gamma}{2} \cos \frac{\beta + \gamma - \alpha}{2} \cos \frac{\gamma + \alpha - \beta}{2}.$

15.  $-4 \cos \left( \frac{\alpha + \beta + \gamma}{2} - \frac{\pi}{4} \right) \Pi \cos \left( \frac{\beta + \gamma - \alpha}{2} + \frac{\pi}{4} \right).$

22. (1)  $(a^2 + b^2)x^2 - 2bcx + c^2 - a^2 = 0;$

(2)  $(a^2 + b^2)^2 x^2 - 2(a^2 - b^2)(2c^2 - a^2 - b^2)x + a^4 + b^4 + 4c^4 - 2a^2b^2 - 4a^2c^2 - 4b^2c^2 = 0.$

[Use  $\cos 2\alpha \cos 2\beta = \cos^2(\alpha - \beta) - \sin^2(\alpha + \beta).$ ]



**XXV. a.** PAGE 318.

1.  $2\sqrt{pq}$ .      2. 4.      3. 24.      4. 2.      7.  $\sqrt{2}$ .      8. 2.
9.  $\sqrt{a^2 - 2ab \sin \alpha + b^2}$ .      10.  $\sqrt{p^2 + 2pq \sin \alpha + q^2}$ .
11. Maximum  $= 2 \sin \frac{\sigma}{2}$ .      12. Maximum  $= \sin^2 \frac{\sigma}{2}$ .
13. Minimum  $= 2 \tan \frac{\sigma}{2}$ .      14. Minimum  $= 2 \operatorname{cosec} \frac{\sigma}{2}$ .
15. Maximum  $= \frac{1}{8}$ .      16. Minimum  $= \sqrt{3}$ .
17. Minimum  $= \frac{3}{4}$ .      18. Minimum  $= 6$ .
19. Minimum  $= 1$ .      20. Minimum  $= 1$ .
21.  $\frac{1}{2}(a+c) \pm \frac{1}{2}\sqrt{b^2 + (a-c)^2}$ .      25.  $\frac{5}{3}$ .
26.  $k^2/(a^2 + b^2 + c^2)$ ;  $k^2/(a+b+c)$ .

**XXV. b.** PAGE 324.

1.  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 2$ .      2.  $x^2 + y^2 = a^2 + b^2$ .      3.  $b^2 = a^2(2 - a^2)$ .
4.  $y(x^2 - 1) = 2$ .      5.  $(a^2 - b^2)^2 = 16ab$ .      6.  $x^{\frac{4}{3}}y^{\frac{2}{3}} - x^{\frac{2}{3}}y^{\frac{4}{3}} = 1$ .
7.  $a^2b^2(a^2 + b^2) = 1$ .      8.  $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$ .      9.  $x^{\frac{4}{5}}y^{\frac{6}{5}} - x^{\frac{6}{5}}y^{\frac{4}{5}} = a^2$ .
10.  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ .      12.  $x^{\frac{1}{2}} + y^{\frac{1}{2}} = 2$ .      13.  $(x+y)^{\frac{2}{3}} + (x-y)^{\frac{2}{3}} = 2$ .
16.  $a^2 + b^2 = 2c^2$ .      20.  $(x+y)^{\frac{2}{3}} + (x-y)^{\frac{2}{3}} = 2a^{\frac{2}{3}}$ .      21.  $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$ .
22.  $\frac{x^2}{a} + \frac{y^2}{b} = a + b$ , or  $\{a(y^2 - b^2) - b(x^2 - a^2)\}^2 = -4abx^2y^2$ .
24.  $xy = (y - x) \tan \alpha$ .      25.  $a^2 + b^2 - 2 \cos \alpha = 2$ .      26.  $a + b = 2ab$ .
29.  $(a+b)(m+n) = 2mn$ .      30.  $x^2 + y^2 = 16a^2$ .
31.  $(a-b)\{c^2 - (a+b)^2\} = 4abcm$ .

## XXV. c. PAGE 334.

1.  $2 \cos 20^\circ, -2 \cos 40^\circ, -2 \cos 80^\circ.$
2.  $2 \sin 10^\circ, 2 \sin 50^\circ, -2 \sin 70^\circ.$
3.  $2 \cos 10^\circ, -2 \cos 50^\circ, -2 \cos 70^\circ.$
4.  $\sin 15^\circ, \sin 45^\circ, -\sin 75^\circ.$
5.  $\frac{1}{a} \sin A, \frac{1}{a} \sin (60^\circ - A), -\frac{1}{a} \sin (60^\circ + A).$
6.  $2a \cos A, 2a \cos (120^\circ \pm A).$
16. (1)  $8x^3 - 4x^2 - 4x + 1 = 0$ ; (2)  $64y^3 - 80y^2 + 24y - 1 = 0.$
17.  $64y^3 - 112y^2 + 56y - 7 = 0.$
18. (1)  $16x^4 + 8x^3 - 12x^2 - 4x + 1 = 0$ ; (2)  $16x^4 - 8x^3 - 12x^2 + 4x + 1 = 0.$
19.  $256y^4 - 448y^3 + 240y^2 - 40y + 1 = 0.$
20.  $t^3 - 36t^6 + 126t^4 - 84t^2 + 9 = 0.$
21.  $32x^5 - 16x^4 - 32x^3 + 12x^2 + 6x - 1 = 0.$

# MATHEMATICAL WORKS

BY

Messrs HALL AND KNIGHT,

PUBLISHED BY

MACMILLAN AND CO.

---

**ALGEBRA FOR BEGINNERS.** By H. S. HALL, M.A., formerly Scholar of Christ's College, Cambridge, Master of the Military Side, Clifton College; and S. R. KNIGHT, B.A., M.B., Ch.B., formerly Scholar of Trinity College, Cambridge, late Assistant Master at Marlborough College. Globe 8vo. 2s. With Answers. 2s. 6d.

*Educational Times*:—"We pronounce this book the best of its size which we have seen."

*University Correspondent*:—"Those masters who have already adopted Messrs Hall and Knight's *Elementary Algebra* in their schools, will welcome this new work for the use of their junior classes . . . The numerous exercises for the student are excellent in quality and entirely new. We can unhesitatingly recommend the book to the notice of both teachers and students."

## ELEMENTARY ALGEBRA FOR SCHOOLS.

Sixth Edition. Revised and Enlarged. Globe 8vo. (bound in maroon-coloured cloth), 3s. 6d. With Answers (bound in green-coloured cloth), 4s. 6d. ANSWERS, 1s.

### PREFACE TO THE SIXTH EDITION.

In previous editions the book has been enlarged by introducing chapters on the Theory of Quadratic Equations, Permutations, and Combinations, the Binomial Theorem, Logarithms and Scales of Notation; the number of Miscellaneous Examples has also been increased. In the present edition we have added chapters on Exponential and Logarithmic Series, Miscellaneous Equations, and Interest and Annuities. For a fuller and more exhaustive treatment of these later chapters readers are referred to our *Higher Algebra*. The very favourable reception accorded to previous editions leads us to hope that in its present more complete form the work will be found suitable as a first

---

MACMILLAN AND CO., LONDON.

## *Works by H. S. HALL, M.A. and S. R. KNIGHT, B.A.*

text-book for every class of student, and amply sufficient for all whose study of Algebra does not extend much beyond the Binomial Theorem.

*Academy*:—"We will not say that this is the best Elementary Algebra for school use that we have come across, but we can say that we do not remember to have seen a better . . . . It is the outcome of a long experience of school teaching, and so is a thoroughly practical book. All others that we have in our eye are the works of men who have had considerable experience with senior and junior students at the universities, but have had little, if any, acquaintance with the poor creatures who are just stumbling over the threshold of Algebra . . . . Buy or borrow the book for yourselves and judge, or write a better . . . . A higher text-book is on its way. This occupies sufficient ground for the generality of boys."

*Educational Times*: ". . . . A very good book. The explanations are concise and clear, and the examples both numerous and well chosen."

**KEY TO ELEMENTARY ALGEBRA FOR SCHOOLS.** Adapted to the Sixth Enlarged Edition of the Elementary Algebra. Second Edition, revised and corrected. Crown 8vo. 8s. 6d.

**HIGHER ALGEBRA.** A Sequel to Elementary Algebra for Schools. Fourth Edition, revised and enlarged. Crown 8vo. 7s. 6d.

\* \* *The Fourth edition contains a collection of three hundred Miscellaneous Examples which will be found useful for advanced Students. These examples have been selected mainly from recent Scholarship or Senate House papers.*

*The Athenæum*:—"The *Elementary Algebra* by the same authors, which has already reached a third edition, is a work of such exceptional merit that those acquainted with it will form high expectations of the sequel to it now issued. Nor will they be disappointed. Of the authors' *Higher Algebra* as of their *Elementary Algebra*, we unhesitatingly assert that it is by far the best work of its kind with which we are acquainted. It supplies a want much felt by teachers."

---

MACMILLAN AND CO., LONDON.

*Works by H. S. HALL, M.A. and S. R. KNIGHT, B.A.*

**KEY TO HIGHER ALGEBRA.** Adapted to the Fourth and enlarged Edition of the Higher Algebra. Crown 8vo. 10s. 6d.

**ALGEBRAICAL EXERCISES AND EXAMINATION PAPERS.** With or without Answers. Third Edition. Globe 8vo. 2s. 6d.

This book has been compiled as a suitable companion to the *Elementary Algebra* by the same Authors. It consists of one hundred and twenty progressive Miscellaneous Exercises, followed by a comprehensive collection of Papers set at recent Examinations.

**ARITHMETICAL EXERCISES AND EXAMINATION PAPERS.** With an Appendix containing Questions in LOGARITHMS AND MENSURATION. With or without Answers. Third Edition, enlarged. Globe 8vo. 2s. 6d.

*In the Second Edition the Appendix has been increased by a new series of Examples which are intended to be worked by the aid of Logarithmic Tables. In view of the increasing importance of logarithmic calculation in many Examinations this last section will be found especially useful.*

**ELEMENTARY TRIGONOMETRY.** Globe 8vo. 4s. 6d. [Ready in December.

*EXTRACT FROM THE PREFACE.*

“This work will be found to comprise all the parts of Elementary Trigonometry which can conveniently be treated without the use of infinite series and imaginary quantities.

“The manuscript and proof sheets have been tested with pupils in all stages of proficiency; we hope, therefore, that our treatment of the subject will be found sufficiently simple for beginners, and at the same time complete enough for all who do not require to study the more advanced parts of the subject.

---

MACMILLAN AND CO., LONDON.



*Works by H. S. HALL, M.A., S. R. KNIGHT, B.A.  
and F. H. Stevens, M.A.*

"In the early pages special prominence has been given to easy identities and equations, so as to familiarise the reader as soon as possible with the Trigonometrical Ratios and their application; with the same end in view the subject of Radian Measure has been postponed later than usual. The sections which treat of Logarithms and Problems in Heights and Distances have been dealt with very fully, and illustrated by a selection of examples carefully classified and graduated. Throughout the work the chapters have been divided into short sections, and the examples, which are very numerous, have been arranged with considerable care; the easier ones in each section being simple enough for any student to whom the subject is new, while the later chapters, and the miscellaneous examples scattered through the book, will furnish good practice for readers of some Mathematical ability."

---

**HIGHER TRIGONOMETRY.** By S. R. KNIGHT,  
B.A., M.B., Ch.B. *[In preparation.]*

---

**A TEXT-BOOK OF EUCLID'S ELEMENTS,** including Alternative Proofs, together with additional Theorems and Exercises, classified and arranged. By H. S. HALL, M.A., and F. H. STEVENS, M.A., Masters of the Military Side, Clifton College.

**THIRD EDITION, ENLARGED.**

Books I.-VI. and XI. Globe 8vo. 4s. 6d.

Also in parts separately as follows:—

Book I. . . . . 1s.	Books III.-VI. . . . . 3s.
Books I. and II. . . . 1s. 6d.	Books I.-IV. . . . . 3s.
Books II. and III. <i>[In the Press.]</i>	Books V., VI., and XI. . 2s. 6d.
Books III. and IV. . . 2s.	Book XI. . . . . 1s.

**KEY TO A TEXT-BOOK OF EUCLID'S ELEMENTS.**

Books I.-IV. Crown 8vo. 6s. 6d.

Books VI. and XI. Crown 8vo. 3s. 6d.

Books I.-VI. and XI. Crown 8vo. 8s. 6d.

---

MACMILLAN AND CO., LONDON.



# A Catalogue OF Educational Books

PUBLISHED BY

## Macmillan & Co.

BEDFORD STREET, STRAND, LONDON

*For books of a less educational character on the subjects named below, see  
Macmillan and Co.'s Classified Catalogue of Books in General Literature.*

### CONTENTS

GREEK AND LATIN	PAGE		PAGE
<b>CLASSICS—</b>		ASTRONOMY . . . . .	30
ELEMENTARY CLASSICS . . . . .	2	HISTORICAL . . . . .	30
CLASSICAL SERIES . . . . .	4	<b>NATURAL SCIENCES—</b>	
CLASSICAL LIBRARY; Texts, Com- mentaries, Translations . . . . .	6	CHEMISTRY . . . . .	30
GRAMMAR, COMPOSITION, AND PHI- LOLOGY . . . . .	8	PHYSICAL GEOGRAPHY, GEOLOGY, AND MINERALOGY . . . . .	32
ANTIQUITIES, ANCIENT HISTORY, AND PHILOSOPHY . . . . .	11	BIOLOGY—	
<b>MODERN LANGUAGES AND</b>		Botany . . . . .	33
<b>LITERATURE—</b>		Zoology . . . . .	33
ENGLISH . . . . .	13	General Biology . . . . .	34
FRENCH . . . . .	17	Physiology . . . . .	35
GERMAN . . . . .	19	MEDICINE . . . . .	35
MODERN GREEK . . . . .	20	<b>HUMAN SCIENCES—</b>	
ITALIAN . . . . .	20	MENTAL AND MORAL PHILOSOPHY . . . . .	36
SPANISH . . . . .	20	POLITICAL ECONOMY . . . . .	37
<b>MATHEMATICS—</b>		LAW AND POLITICS . . . . .	38
ARITHMETIC . . . . .	21	ANTHROPOLOGY . . . . .	38
BOOK-KEEPING . . . . .	22	EDUCATION . . . . .	39
ALGEBRA . . . . .	22	<b>TECHNICAL KNOWLEDGE—</b>	
EUCLID AND PURE GEOMETRY . . . . .	22	CIVIL AND MECHANICAL ENGINEER- ING . . . . .	39
GEOMETRICAL DRAWING . . . . .	23	MILITARY AND NAVAL SCIENCE . . . . .	39
MENSURATION . . . . .	24	AGRICULTURE AND FORESTRY . . . . .	40
TRIGONOMETRY . . . . .	24	DOMESTIC ECONOMY . . . . .	40
ANALYTICAL GEOMETRY . . . . .	25	HYGIENE . . . . .	40
PROBLEMS AND QUESTIONS IN MA- THEMATICS . . . . .	25	COMMERCE . . . . .	41
HIGHER PURE MATHEMATICS . . . . .	25	TECHNOLOGY . . . . .	41
MECHANICS . . . . .	26	<b>GEOGRAPHY . . . . .</b>	41
PHYSICS . . . . .	28	<b>HISTORY . . . . .</b>	42
		<b>ART . . . . .</b>	44
		<b>DIVINITY . . . . .</b>	44

# GREEK AND LATIN CLASSICS.

Elementary Classics; Classical Series; Classical Library, (1) Texts, (2) Translations; Grammar, Composition, and Philology; Antiquities, Ancient History, and Philosophy.

## \*ELEMENTARY CLASSICS.

18mo, Eighteenpence each.

The following contain Introductions, Notes, and Vocabularies, and in some cases Exercises.

**ACCIDENCE, LATIN, AND EXERCISES ARRANGED FOR BEGINNERS.**—By W. WELCH, M.A., and C. G. DUFFIELD, M.A.

**AESCHYLUS.—PROMETHEUS VINCTUS.** By Rev. H. M. STEPHENSON, M.A.

**ARRIAN.—SELECTIONS.** With Exercises. By Rev. JOHN BOND, M.A., and Rev. A. S. WALPOLE, M.A.

**AULUS GELLIUS, STORIES FROM.**—Adapted for Beginners. With Exercises. By Rev. G. H. NALL, M.A., Assistant Master at Westminster.

**CÆSAR.—THE HELVETIAN WAR.** Selections from Book I., adapted for Beginners. With Exercises. By W. WELCH, M.A., and C. G. DUFFIELD, M.A.

**THE INVASION OF BRITAIN.** Selections from Books IV. and V., adapted for Beginners. With Exercises. By the same.

**SCENES FROM BOOKS V. AND VI.** By C. COLBECK, M.A.

**SCENES FROM THE CIVIL WAR.** By C. H. KEENE.

**THE GALLIC WAR. BOOK I.** By Rev. A. S. WALPOLE, M.A.

**BOOKS II. AND III.** By the Rev. W. G. RUTHERFORD, M.A., LL.D.

**BOOK IV.** By CLEMENT BRYANS, M.A., Assistant Master at Dulwich College.

**BOOK V.** By C. COLBECK, M.A., Assistant Master at Harrow.

**BOOK VI.** By C. COLBECK, M.A.

**BOOK VII.** By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

**THE CIVIL WAR. BOOK I.** By M. MONTGOMREY, M.A.

**CICERO.—DE SENECTUTE.** By E. S. SHUCKBURGH, M.A.

**DE AMICITIA.** By the same.

**STORIES OF ROMAN HISTORY.** Adapted for Beginners. With Exercises. By Rev. G. E. JEANS, M.A., and A. V. JONES, M.A.

**CURTIUS (Quintus).—SELECTIONS.** Adapted for Beginners. With Notes, Vocabulary, and Exercises. By F. COVERLEY SMITH. [*In preparation.*]

**EURIPIDES.—ALCESTIS.** By Rev. M. A. BAYFIELD, M.A.

**MEDEA.** By Rev. M. A. BAYFIELD, M.A.

**HECUBA.** By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

**EUTROPIUS.**—Adapted for Beginners. With Exercises. By W. WELCH, M.A., and C. G. DUFFIELD, M.A.

**BOOKS I. and II.** By the same.

**EXERCISES IN UNSEEN TRANSLATION IN LATIN.** By W. WELCH, M.A., and Rev. C. G. DUFFIELD, M.A.

**HERODOTUS, TALES FROM.** Atticised. By G. S. FARNELL, M.A.

**HOMER.—ILIAD. BOOK I.** By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

**BOOK VI.** By WALTER LEAF, Litt.D., and Rev. M. A. BAYFIELD.

**BOOK XVIII.** By S. R. JAMES, M.A., Assistant Master at Eton.

**BOOK XXIV.** By W. LEAF, Litt.D., and Rev. M. A. BAYFIELD, M.A. [*In the Press.*]

**ODYSSEY. BOOK I.** By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

**HORACE.—ODES. BOOKS I. II. III. AND IV.** separately. By T. E. PAGE, M.A., Assistant Master at the Charterhouse. Each 1s. 6d.

**LIVY.—BOOK I.** By H. M. STEPHENSON, M.A.

**BOOK V.** By M. ALFORD.

**BOOK XXI.** Adapted from Mr. Capes's Edition. By J. E. MELHUISE, M.A.

- BOOK XXII. Adapted from Mr. CAPES's Edition. By J. E. MELHUISE, M.A.  
 SELECTIONS FROM BOOKS V. and VI. By W. CECIL LAMING, M.A.  
 THE HANNIBALIAN WAR. BOOKS XXI. and XXII. adapted by G. C. MACAULAY, M.A.  
 BOOKS XXIII. and XXIV. adapted by the same. *[In preparation.]*  
 THE SIEGE OF SYRACUSE. Being part of the XXIV. and XXV. BOOKS OF LIVY, adapted for Beginners. With Exercises. By G. RICHARDS, M.A., and Rev. A. S. WALPOLE, M.A.  
 LEGENDS OF ANCIENT ROME. Adapted for Beginners. With Exercises. By H. WILKINSON, M.A.  
 LUCIAN.—EXTRACTS FROM LUCIAN. With Exercises. By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.  
 NEPOS.—SELECTIONS ILLUSTRATIVE OF GREEK AND ROMAN HISTORY. With Exercises. By G. S. FARNELL, M.A.  
 OVID.—SELECTIONS. By E. S. SHUCKBURGH, M.A.  
 EASY SELECTIONS FROM OVID IN ELEGIAC VERSE. With Exercises. By H. WILKINSON, M.A.  
 METAMORPHOSES. BOOK I. By CHARLES SIMMONS, M.A. *[In preparation.]*  
 STORIES FROM THE METAMORPHOSES. With Exercises. By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.  
 TRISTIA.—BOOK I. By E. S. SHUCKBURGH, M.A. *[In preparation.]*  
 BOOK III. By E. S. SHUCKBURGH, M.A. *[In preparation.]*  
 PHÆDRUS.—SELECT FABLES. Adapted for Beginners. With Exercises. By Rev. A. S. WALPOLE, M.A.  
 THUCYDIDES.—THE RISE OF THE ATHENIAN EMPIRE. BOOK I. CHS. 89-117 and 228-238. With Exercises. By F. H. COLSON, M.A.  
 SELECTIONS FROM BOOKS II. AND III. THE SIEGE OF PLATÆA. By W. T. SUTHERY, M.A., and A. S. GRAVES, M.A. *[In the Press.]*  
 VIRGIL.—SELECTIONS. By E. S. SHUCKBURGH, M.A.  
 BUCOLICS. By T. E. PAGE, M.A.  
 GEORGICS. BOOK I. By T. E. PAGE, M.A.  
 BOOK II. By Rev. J. H. SKRINE, M.A.  
 BOOK III. By T. E. PAGE, M.A. *[In preparation.]*  
 BOOK IV. By T. E. PAGE, M.A. *[In preparation.]*  
 ÆNEID. BOOK I. By Rev. A. S. WALPOLE, M.A.  
 BOOK I. By T. E. PAGE, M.A.  
 BOOK II. By T. E. PAGE, M.A.  
 BOOK III. By T. E. PAGE, M.A.  
 BOOK IV. By Rev. H. M. STEPHENSON, M.A.  
 BOOK V. By Rev. A. CALVERT, M.A.  
 BOOK VI. By T. E. PAGE, M.A.  
 BOOK VII. By Rev. A. CALVERT, M.A.  
 BOOK VIII. By Rev. A. CALVERT, M.A.  
 BOOK IX. By Rev. H. M. STEPHENSON, M.A.  
 BOOK X. By S. G. OWEN, M.A.  
 XENOPHON.—ANABASIS. Selections, adapted for Beginners. With Exercises. By W. WELCH, M.A., and C. G. DUFFIELD, M.A.  
 BOOK I. With Exercises. By E. A. WELLS, M.A.  
 BOOK I. By Rev. A. S. WALPOLE, M.A.  
 BOOK II. By Rev. A. S. WALPOLE, M.A.  
 BOOK III. By Rev. G. H. NALL, M.A.  
 BOOK IV. By Rev. E. D. STONE, M.A.  
 BOOK V. By Rev. G. H. NALL, M.A.  
 BOOK VI. By Rev. G. H. NALL, M.A.  
 SELECTIONS FROM BOOK IV. With Exercises. By Rev. E. D. STONE, M.A.  
 SELECTIONS FROM THE CYROPÆDIA. With Exercises. By A. H. COOKE, M.A.  
 TALES FROM THE CYROPÆDIA. With Exercises. By CHARLES H. KEENE.
- The following contain Introductions and Notes, but no Vocabulary:—  
 CICERO.—SELECT LETTERS. By Rev. G. E. JEANS, M.A.  
 HERODOTUS.—SELECTIONS FROM BOOKS VII. AND VIII. THE EXPEDITION OF XERXES. By A. H. COOKE, M.A.

- HORACE.**—SELECTIONS FROM THE SATIRES AND EPISTLES. By Rev. W. J. V. BAKER, M.A.  
**SELECT EPODES AND ARS POETICA.** By H. A. DALTON, M.A.  
**PLATO.**—EUTHYPHRO AND MENEXENUS. By C. E. GRAVES, M.A.  
**TERENCE.**—SCENES FROM THE ANDRIA. By F. W. CORNISH, M.A., Assistant Master at Eton.  
**THE GREEK ELEGIAC POETS.**—FROM CALLINUS TO CALLIMACHUS. Selected by Rev. HERBERT KYNASTON, D.D.  
**THUCYDIDES.**—BOOK IV. CHS. 1-41. THE CAPTURE OF SPHACTERIA. By C. E. GRAVES, M.A.

## CLASSICAL SERIES FOR COLLEGES AND SCHOOLS.

Fcap. 8vo.

- ÆSCHINES.**—IN CTESIPHONTA. By Rev. T. GWATKIN, M.A., and E. S. SHUCKBURGH, M.A. 5s.  
**ÆSCHYLUS.**—PERSÆ. By A. O. PRICKARD, M.A., Fellow and Tutor of New College, Oxford. With Map. 2s. 6d.  
**SEVEN AGAINST THEBES.** SCHOOL EDITION. By A. W. VERRALL, Litt.D., and M. A. BAYFIELD, M.A. 2s. 6d.  
**ANDOCIDES.**—DE MYSTERIIS. By W. J. HICKIE, M.A. 2s. 6d.  
**ATTIC ORATORS.**—Selections from ANTIPHON, ANDOCIDES, LYSIAS, ISO-CRATES, and ISAEUS. By R. C. JEBB, Litt.D., Regius Professor of Greek in the University of Cambridge. 5s.  
**\*CÆSAR.**—THE GALLIC WAR. By Rev. JOHN BOND, M.A., and Rev. A. S. WALPOLE, M.A. With Maps. 4s. 6d.  
**CATULLUS.**—SELECT POEMS. By F. P. SIMPSON, B.A. 3s. 6d. The Text of this Edition is carefully expurgated for School use.  
**\*CICERO.**—THE CATILINE ORATIONS. By A. S. WILKINS, Litt.D., Professor of Latin, Owens College, Manchester. 2s. 6d.  
**PRO LEGE MANILIA.** By Prof. A. S. WILKINS, Litt.D. 2s. 6d.  
**THE SECOND PHILIPPIC ORATION.** By JOHN E. B. MAYOR, M.A., Professor of Latin in the University of Cambridge. 3s. 6d.  
**PRO ROSCIO AMERINO.** By E. H. DONKIN, M.A. 2s. 6d.  
**PRO P. SESTIO.** By Rev. H. A. HOLDEN, Litt.D. 3s. 6d.  
**PRO MILONE.** By F. H. COLSON, M.A. 2s. 6d.  
**SELECT LETTERS.** By R. Y. TYRRELL, M.A. 4s. 6d.  
**DEMOSTHENES.**—DE CORONA. By B. DRAKE, M.A. 7th Edition, revised by E. S. SHUCKBURGH, M.A. 3s. 6d.  
**ADVERSUS LEPTINEM.** By Rev. J. R. KING, M.A., Fellow and Tutor of Oriel College, Oxford. 2s. 6d.  
**THE FIRST PHILIPPIC.** By Rev. T. GWATKIN, M.A. 2s. 6d.  
**IN MIDIAM.** By Prof. A. S. WILKINS, Litt.D., and HERMAN HAGER, Ph.D., the Owens College, Victoria University, Manchester. *[In preparation.]*  
**EURIPIDES.**—HIPPOLYTUS. By Rev. J. P. MAHAFFY, D.D., Fellow of Trinity College, and Professor of Ancient History in the University of Dublin, and J. B. BURY, M.A., Fellow of Trinity College, Dublin. 2s. 6d.  
**MEDEA.** By A. W. VERRALL, Litt.D., Fellow of Trinity College, Cambridge. 2s. 6d.  
**IPHIGENIA IN TAURIS.** By E. B. ENGLAND, M.A. 3s.  
**ION.** By M. A. BAYFIELD, M.A., Headmaster of Christ College, Brecon. 2s. 6d.  
**BACCHAE.** By R. Y. TYRRELL, M.A., Regius Professor of Greek in the University of Dublin. 3s. 6d.  
**HERODOTUS.**—BOOK III. By G. C. MACAULAY, M.A. 2s. 6d.  
**BOOK V.** By J. STRACHAN, M.A., Professor of Greek, Owens College, Manchester. *[In preparation.]*  
**BOOK VI.** By the same. 3s. 6d.  
**BOOK VII.** By Mrs. MONTAGU BUTLER. 3s. 6d.



- HOMER.**—**ILIAD.** In 4 vols. Edited by W. LEAF, Litt.D., and Rev. M. A. BAYFIELD, M.A. [Vol. I. *in the Press.*]  
**ILIAD.** BOOKS I., IX., XI., XVI.—XXIV. **THE STORY OF ACHILLES.** By the late J. H. PRATT, M.A., and WALTER LEAF, Litt.D., Fellows of Trinity College, Cambridge. 5s.  
**ODYSSEY.** BOOK IX. By Prof. JOHN E. B. MAYOR. 2s. 6d.  
**ODYSSEY.** BOOKS XXI.—XXIV. **THE TRIUMPH OF ODYSSEUS.** By S. G. HAMILTON, M.A., Fellow of Hertford College, Oxford. 2s. 6d.  
**HORACE.**—\***THE ODES.** By T. E. PAGE, M.A., Assistant Master at the Charterhouse. 5s. (BOOKS I. II. III. and IV. separately, 2s. each.)  
**THE SATIRES.** By ARTHUR PALMER, M.A., Professor of Latin in the University of Dublin. 5s.  
**THE EPISTLES AND ARS POETICA.** By Prof. A. S. WILKINS, Litt.D. 5s.  
**JUVENAL.**—\***THIRTEEN SATIRES.** By E. G. HARDY, M.A. 5s. The Text is carefully expurgated for School use.  
**SELECT SATIRES.** By Prof. JOHN E. B. MAYOR. XII.—XVI. 4s. 6d.  
**LIVY.**—\***BOOKS II. and III.** By Rev. H. M. STEPHENSON, M.A. 3s. 6d.  
 \***BOOKS XXI. and XXII.** By Rev. W. W. CAPE, M.A. With Maps. 4s. 6d.  
 \***BOOKS XXIII. and XXIV.** By G. C. MACAULAY, M.A. With Maps. 3s. 6d.  
 \***THE LAST TWO KINGS OF MACEDON. EXTRACTS FROM THE FOURTH AND FIFTH DECADES OF LIVY.** By F. H. RAWLINS, M.A., Assistant Master at Eton. With Maps. 2s. 6d.  
**LUCRETIUS.**—**BOOKS I.—III.** By J. H. WARBURTON LEE, M.A., late Assistant Master at Rossall. 3s. 6d.  
**LYSIAS.**—**SELECT ORATIONS.** By E. S. SHUCKBURGH, M.A. 5s.  
**MARTIAL.**—**SELECT EPIGRAMS.** By Rev. H. M. STEPHENSON, M.A. 5s.  
 \***OVID.**—**FASTI.** By G. H. HALLAM, M.A., Assistant Master at Harrow. 3s. 6d.  
 \***HEROIDUM EPISTULÆ XIII.** By E. S. SHUCKBURGH, M.A. 3s. 6d.  
**METAMORPHOSES.** BOOKS I.—III. By C. SIMMONS, M.A. [*In preparation.*]  
 BOOKS XIII. and XIV. By the same. 3s. 6d.  
**PLATO.**—**LACHES.** By M. T. TATHAM, M.A. 2s. 6d.  
**THE REPUBLIC.** BOOKS I.—V. By T. H. WARREN, M.A., President of Magdalen College, Oxford. 5s.  
**PLAUTUS.**—**MILES GLORIOSUS.** By R. Y. TYRRELL, M.A., Regius Professor of Greek in the University of Dublin. 2nd Ed., revised. 3s. 6d.  
**AMPHITRUO.** By Prof. ARTHUR PALMER, M.A. 3s. 6d.  
**CAPTIVI.** By A. R. S. HALLIDIE, M.A. 3s. 6d.  
**PLINY.**—**LETTERS.** BOOKS I. and II. By J. COWAN, M.A., Assistant Master at the Manchester Grammar School. 3s.  
**LETTERS.** BOOK III. By Prof. JOHN E. B. MAYOR. With Life of Pliny by G. H. RENDALL, M.A. 3s. 6d.  
**PLUTARCH.**—**LIFE OF THEMISTOKLES.** By Rev. H. A. HOLDEN, Litt.D. 3s. 6d.  
**LIVES OF GALBA AND OTHO.** By E. G. HARDY, M.A. 5s.  
**LIFE OF PERICLES.** By Rev. H. A. HOLDEN, Litt.D. [*In the Press.*]  
**POLYBIUS.**—**THE HISTORY OF THE ACHÆAN LEAGUE AS CONTAINED IN THE REMAINS OF POLYBIUS.** By Rev. W. W. CAPE, M.A. 5s.  
**PROPERTIUS.**—**SELECT POEMS.** By Prof. J. P. POSTGATE, Litt.D. 2nd Ed. 5s.  
**SALLUST.**—\***CATILINA and JUGURTHA.** By C. MERIVALE, D.D., Dean of Ely. 3s. 6d. Or separately, 2s. each.  
 \***BELLUM CATULINÆ.** By A. M. COOK, M.A. 2s. 6d.  
**JUGURTHA.** By the same. [*In preparation.*]  
**TACITUS.**—**THE ANNALS.** BOOKS I. and II. By J. S. REID, Litt.D. [*In prep.*]  
 BOOK VI. By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. 2s.  
**THE HISTORIES.** BOOKS I. and II. By A. D. GODLEY, M.A. 3s. 6d.  
 BOOKS III.—V. By the same. 3s. 6d.  
**AGRICOLA and GERMANIA.** By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. 3s. 6d. Or separately, 2s. each.  
**AGRICOLA and GERMANIA (separately).** By F. J. HAVERFIELD, M.A., Student of Christ Church, Oxford. [*In preparation.*]  
**TERENCE.**—**HAUTON TIMORUMENOS.** By E. S. SHUCKBURGH, M.A. 2s. 6d.  
 With Translation. 3s. 6d.  
**PHORMIO.** By Rev. JOHN BOND, M.A., and Rev. A. S. WALPOLE, M.A. 2s. 6d.  
**ADELPHOE.** By Prof. S. G. ASHMORE. 3s. 6d.

- THUCYDIDES.**—BOOK I. By CLEMENT BRYANS, M.A. [*In preparation.*]  
 BOOK II. By E. C. MARCHANT, M.A., Fellow of St. Peter's Coll., Cam. 3s. 6d.  
 BOOK III. By E. C. MARCHANT, M.A. [*In preparation.*]  
 BOOK IV. By C. E. GRAVES, M.A., Classical Lecturer at St. John's College, Cambridge. 3s. 6d.  
 BOOK V. By C. E. GRAVES, M.A. 3s. 6d.  
 BOOKS VI. AND VII. By Rev. PERCIVAL FROST, M.A. With Map. 3s. 6d.  
 BOOK VI. By E. C. MARCHANT, M.A. [*In preparation.*]  
 BOOK VII. By E. C. MARCHANT, M.A. 3s. 6d.  
 BOOK VIII. By Prof. T. G. TUCKER, Litt.D. 3s. 6d.  
**TIBULLUS.**—SELECT POEMS. By Prof. J. P. POSTGATE, Litt.D. [*In preparation.*]  
**VIRGIL.**—ÆNEID. BOOKS II. AND III. THE NARRATIVE OF ÆNEAS. By E. W. HOWSON, M.A., Assistant Master at Harrow. 2s.  
**XENOPHON.**—\*THE ANABASIS. BOOKS I.—IV. By Profs. W. W. GOODWIN and J. W. WHITE. Adapted to Goodwin's Greek Grammar. With Map. 3s. 6d.  
 BOOKS V.—VII. By Rev. G. H. NALL, M.A. [*In preparation.*]  
 HELLENICA. BOOKS I. AND II. By H. HAILSTONE, B.A. With Map. 2s. 6d.  
 BOOK III. By H. G. DAKYNS, M.A. [*In the Press.*]  
 BOOK IV. By the same. [*In preparation.*]  
 CYROPÆDIA. BOOKS VII. AND VIII. By A. GOODWIN, M.A. 2s. 6d.  
 MEMORABILIA SOCRATIS. By A. R. CLUER, B.A., Balliol College, Oxford. 5s.  
 HERODOTUS. By Rev. H. A. HOLDEN, Litt.D. 2s. 6d.  
 OECONOMICUS. By the same. With Lexicon. 5s.

## CLASSICAL LIBRARY.

**Texts, Edited with Introductions and Notes, for the use of Advanced Students; Commentaries and Translations.**

- ÆSCHYLUS.**—THE SUPPLICES. A Revised Text, with Translation. By T. G. TUCKER, Litt.D., Professor of Classical Philology in the University of Melbourne. 8vo. 10s. 6d.  
 THE SEVEN AGAINST THEBES. With Translation. By A. W. VERRALL, Litt.D., Fellow of Trinity College, Cambridge. 8vo. 7s. 6d.  
 AGAMEMNON. With Translation. By A. W. VERRALL, Litt.D. 8vo. 12s.  
 THE CHOEPHORI. With Translation. By A. W. VERRALL, Litt.D. 8vo. 12s.  
 AGAMEMNON, CHOEPHORI, AND EUMENIDES. By A. O. PRICKARD, M.A., Fellow and Tutor of New College, Oxford. 8vo. [*In preparation.*]  
 THE EUMENIDES. With Verse Translation. By B. DRAKE, M.A. 8vo. 5s.  
 ÆSCHYLUS. Translated into English Prose by Prof. T. G. TUCKER. Cr. 8vo. [*In preparation.*]  
**ANTONINUS, MARCUS AURELIUS.**—BOOK IV. OF THE MEDITATIONS. With Translation. By HASTINGS CROSSLEY, M.A. 8vo. 6s.  
**ARISTOPHANES.**—THE BIRDS. Translated into English Verse. By B. H. KENNEDY, D.D. Cr. 8vo. 6s. Help Notes to the Same, for the Use of Students. 1s. 6d.  
 SCHOLIA ARISTOPHANICA; being such Comments adscript to the text of Aristophanes as are preserved in the Codex Ravennas, arranged, emended, and translated. By Rev. W. G. RUTHERFORD, M.A., LL.D. 8vo. [*In the Press.*]  
**ARISTOTLE.**—THE METAPHYSICS. BOOK I. Translated by a Cambridge Graduate. 8vo. 5s.  
 THE POLITICS. By R. D. HICKS, M.A., Fellow of Trinity College, Cambridge. 8vo. [*In the Press.*]  
 THE POLITICS. Translated by Rev. J. E. C. WELLDON, M.A., Headmaster of Harrow. Cr. 8vo. 10s. 6d.  
 THE RHETORIC. Translated by the same. Cr. 8vo. 7s. 6d.  
 AN INTRODUCTION TO ARISTOTLE'S RHETORIC. With Analysis, Notes, and Appendices. By E. M. COPE, Fellow and late Tutor of Trinity College, Cambridge. 8vo. 14s.  
 THE NICOMACHEAN ETHICS. Translated by Rev. J. E. C. WELLDON, M.A. Cr. 8vo. 7s. 6d.  
 THE SOPHISTICI ELENCHI. With Translation. By E. POSTE, M.A., Fellow of Oriel College, Oxford. 8vo. 8s. 6d.



- ON THE CONSTITUTION OF ATHENS. By J. E. SANDYS, Litt.D. 8vo. 15s.  
ON THE CONSTITUTION OF ATHENS. Translated by E. POSTE, M.A. 2nd Ed. Cr. 8vo. 3s. 6d.
- ON THE ART OF POETRY. A Lecture. By A. O. PRICKARD, M.A., Fellow and Tutor of New College, Oxford. Cr. 8vo. 3s. 6d.
- ATTIC ORATORS.—FROM ANTIPHON TO ISAEOS. By R. C. JEBB, Litt.D., Regius Professor of Greek in the University of Cambridge. 2 vols. 8vo. 25s.
- BABRIUS.—With Lexicon. By Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. 8vo. 12s. 6d.
- CATULLUS. By Prof. ARTHUR PALMER. *[In preparation.]*
- CICERO.—THE ACADEMICA. By J. S. REID, Litt.D., Fellow of Caius College, Cambridge. 8vo. 15s.
- THE ACADEMICS. Translated by the same. 8vo. 5s. 6d.
- SELECT LETTERS. After the Edition of ALBERT WATSON, M.A. Translated by G. E. JEANS, M.A., Fellow of Hertford College, Oxford. Cr. 8vo. 10s. 6d.
- EURIPIDES.—MEDEA. By A. W. VERRALL, Litt.D. 8vo. 7s. 6d.
- IPHIGENEIA AT AULIS. By E. B. ENGLAND, Litt.D. 8vo. 7s. 6d.
- \*INTRODUCTION TO THE STUDY OF EURIPIDES. By Professor J. P. MAHAFFY. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)
- HERODOTUS.—BOOKS I.—III. THE ANCIENT EMPIRES OF THE EAST. By A. H. SAYCE, Deputy-Professor of Comparative Philology in the University of Oxford. 8vo. 16s.
- BOOKS IV.—IX. By R. W. MACAN, M.A., Reader in Ancient History in the University of Oxford. 8vo. *[In the Press.]*
- THE HISTORY. Translated by G. C. MACAULAY, M.A. 2 vols. Cr. 8vo. 18s.
- HOMER.—THE ILIAD. By WALTER LEAF, Litt.D. 8vo. Books I.—XII. 14s. Books XIII.—XXIV. 14s.
- COMPANION TO THE ILIAD FOR ENGLISH READERS. By the same. Cr. 8vo. 7s. 6d.
- THE ILIAD. Translated into English Prose by ANDREW LANG, M.A., WALTER LEAF, Litt.D., and ERNEST MYERS, M.A. Cr. 8vo. 12s. 6d.
- THE ODYSSEY. Done into English by S. H. BUTCHER, M.A., Professor of Greek in the University of Edinburgh, and ANDREW LANG, M.A. Cr. 8vo. 6s.
- \*INTRODUCTION TO THE STUDY OF HOMER. By the Right Hon. W. E. GLADSTONE. 18mo. 1s. (*Literature Primers.*)
- HOMERIC DICTIONARY. Translated from the German of Dr. G. AUTENRIETH by R. P. KEEP, Ph.D. Illustrated. Cr. 8vo. 6s.
- HORACE.—Translated by J. LONSDALE, M.A., and S. LEE, M.A. Gl. 8vo. 3s. 6d.
- JUVENAL.—THIRTEEN SATIRES OF JUVENAL. By JOHN E. B. MAYOR, M.A., Professor of Latin in the University of Cambridge. Cr. 8vo. 2 vols. 10s. 6d. each.
- THIRTEEN SATIRES. Translated by ALEX. LEEPER, M.A., LL.D., Warden of Trinity College, Melbourne. Revised Ed. Cr. 8vo. 3s. 6d.
- KTESIAS.—THE FRAGMENTS OF THE PERSIKA OF KTESIAS. By JOHN GILMORE, M.A. 8vo. 8s. 6d.
- LIVY.—BOOKS I.—IV. Translated by Rev. H. M. STEPHENSON, M.A. *[In prep.]*
- BOOKS XXI.—XXV. Translated by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. Cr. 8vo. 7s. 6d.
- \*INTRODUCTION TO THE STUDY OF LIVY. By Rev. W. W. CAPES, M.A. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)
- LONGINUS.—ON THE SUBLIME. Translated by H. L. HAVELL, B.A. With Introduction by ANDREW LANG. Cr. 8vo. 4s. 6d.
- MARTIAL.—BOOKS I. AND II. OF THE EPIGRAMS. By Prof. JOHN E. B. MAYOR, M.A. 8vo. *[In the Press.]*
- MELEAGER.—FIFTY POEMS OF MELEAGER. Translated by WALTER HEADLAM. Fcap. 4to. 7s. 6d.
- PAUSANIAS.—DESCRIPTION OF GREECE. Translated with Commentary by J. G. FRAZER, M.A., Fellow of Trinity College, Cambridge. *[In prep.]*
- PHRYNICHUS.—THE NEW PHRYNICHUS; being a Revised Text of the Ecloga of the Grammarians Phrynichus. With Introduction and Commentary by Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. 8vo. 18s.
- PINDAR.—THE EXTANT ODES OF PINDAR. Translated by ERNEST MYERS, M.A. Cr. 8vo. 5s.

- THE OLYMPIAN AND PYTHIAN ODES. Edited, with an Introductory Essay, by BASIL GILDERSLEEVE, Professor of Greek in the Johns Hopkins University, U.S.A. Cr. 8vo. 7s. 6d.
- THE NEMEAN ODES. By J. B. BURY, M.A., Fellow of Trinity College, Dublin. 8vo. 12s.
- THE ISTHMIAN ODES. By the same Editor. 8vo. 10s. 6d.
- PLATO.—PHÆDO. By R. D. ARCHER-HIND, M.A., Fellow of Trinity College, Cambridge. 8vo. 8s. 6d.
- PHÆDO. By Sir W. D. GEDDES, LL.D., Principal of the University of Aberdeen. 8vo. 8s. 6d.
- TIMÆUS. With Translation. By R. D. ARCHER-HIND, M.A. 8vo. 16s.
- THE REPUBLIC OF PLATO. Translated by J. LL. DAVIES, M.A., and D. J. VAUGHAN, M.A. 18mo. 2s. 6d. net.
- EUTHYPHRO, APOLOGY, CRITO, AND PHÆDO. Translated by F. J. CHURCH. 18mo. 2s. 6d. net.
- PHÆDRUS, LYSIS, AND PROTAGORAS. Translated by J. WRIGHT, M.A. 18mo. 2s. 6d. net.
- PLAUTUS.—THE MOSTELLARIA. By WILLIAM RAMSAY, M.A. Ed. by G. G. RAMSAY, M.A., Professor of Humanity, University of Glasgow. 8vo. 14s.
- PLINY.—CORRESPONDENCE WITH TRAJAN. C. Plinii Caecilii Secundi Epistolæ ad Traianum Imperatorem cum Eiusdem Responsis. By E. G. HARDY, M.A. 8vo. 10s. 6d.
- POLYBIUS.—THE HISTORIES OF POLYBIUS. Translated by E. S. SHUCK-BURGH, M.A. 2 vols. Cr. 8vo. 24s.
- SALLUST.—CATILINE AND JUGURTHA. Translated by A. W. POLLARD, B.A. Cr. 8vo. 6s. THE CATILINE (separately). 3s.
- SOPHOCLES.—ŒDIPUS THE KING. Translated into English Verse by E. D. A. MORSEHEAD, M.A., Assistant Master at Winchester. Fcap. 8vo. 3s. 6d.
- TACITUS.—THE ANNALS. By G. O. HOLBROOKE, M.A., Professor of Latin in Trinity College, Hartford, U.S.A. With Maps. 8vo. 16s.
- THE ANNALS. Translated by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Maps. Cr. 8vo. 7s. 6d.
- THE HISTORIES. By Rev. W. A. SPOONER, M.A., Fellow and Tutor of New College, Oxford. 8vo. 16s.
- THE HISTORY. Translated by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Map. Cr. 8vo. 6s.
- THE AGRICOLA AND GERMANY, WITH THE DIALOGUE ON ORATORY. Translated by the same. With Maps. Cr. 8vo. 4s. 6d.
- \*INTRODUCTION TO THE STUDY OF TACITUS. By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)
- THEOCRITUS, BION, AND MOSCHUS. Translated by A. LANG, M.A. 18mo. 2s. 6d. net. Also an Edition on Large Paper. Cr. 8vo. 9s.
- THUCYDIDES.—BOOK IV. A Revision of the Text, Illustrating the Principal Causes of Corruption in the Manuscripts of this Author. By Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. 8vo. 7s. 6d.
- BOOK VIII. By H. C. GOODHART, M.A., Professor of Latin in the University of Edinburgh. 8vo. 9s.
- VIRGIL.—Translated by J. LONSDALE, M.A., and S. LEE, M.A. Gl. 8vo. 3s. 6d.
- THE ÆNEID. Translated by J. W. MACKAIL, M.A., Fellow of Balliol College, Oxford. Cr. 8vo. 7s. 6d.
- XENOPHON.—Translated by H. G. DAKYNS, M.A. In four vols. Cr. 8vo. Vol. I. "The Anabasis" and "The Hellenica I. and II." 10s. 6d. Vol. II. "Hellenica" III.—VII. "Agesilaus," the "Politics," and "Revenues." 10s. 6d.

[Vol. III. in the Press]

## GRAMMAR, COMPOSITION, & PHILOLOGY.

### Latin.

- \*BELCHER.—SHORT EXERCISES IN LATIN PROSE COMPOSITION AND EXAMINATION PAPERS IN LATIN GRAMMAR. Part I. By Rev. H. BELCHER, LL.D., Rector of the High School, Dunedin, N.Z. 18mo. 1s. 6d. KEY, for Teachers only. 18mo. 3s. 6d.
- \*Part II. On the Syntax of Sentences, with an Appendix, including EXERCISES IN LATIN IDIOMS. etc. 18mo. 2s. KEY, for Teachers only. 18mo. 3s.

- \***BRYANS.**—LATIN PROSE EXERCISES BASED UPON CÆSAR'S GALLIC WAR. With a Classification of Cæsar's Chief Phrases and Grammatical Notes on Cæsar's Usages. By CLEMENT BRYANS, M.A., Assistant Master at Dulwich College. Ex. fcap. 8vo. 2s. 6d. KEY, for Teachers only. 4s. 6d.
- CORNELL UNIVERSITY STUDIES IN CLASSICAL PHILOLOGY.** Edited by I. FLAGG, W. G. HALE, and B. I. WHEELER. I. The *CUM*-Constructions: their History and Functions. By W. G. HALE. Part I. Critical. 1s. 8d. net. Part 2. Constructive. 3s. 4d. net. II. Analogy and the Scope of its Application in Language. By B. I. WHEELER. 1s. 3d. net.
- \***ENGLAND.**—EXERCISES ON LATIN SYNTAX AND IDIOM. ARRANGED WITH REFERENCE TO ROBY'S SCHOOL LATIN GRAMMAR. By E. B. ENGLAND, Litt.D., Assistant Lecturer at the Owens College, Manchester. Cr. 8vo. 2s. 6d. KEY, for Teachers only. 2s. 6d.
- GILES.**—A SHORT MANUAL OF PHILOLOGY FOR CLASSICAL STUDENTS. By P. GILES, M.A., Reader in Comparative Philology in the University of Cambridge. Cr. 8vo. *[In the Press.]*
- HADLEY.**—ESSAYS, PHILOLOGICAL AND CRITICAL. By JAMES HADLEY, late Professor in Yale College. 8vo. 16s.
- HODGSON.**—MYTHOLOGY FOR LATIN VERSIFICATION. Fables for rendering into Latin Verse. By F. HODGSON, B.D., late Provost of Eton. New Ed., revised by F. C. HODGSON, M.A. 18mo. 3s.
- JANNARIS.**—HISTORICAL GRAMMAR OF THE GREEK LANGUAGE. By Prof. A. N. JANNARIS. 8vo. *[In preparation.]*
- LUPTON.**—\*AN INTRODUCTION TO LATIN ELEGIAC VERSE COMPOSITION. By J. H. LUPTON, Sur-Master of St. Paul's School. Gl. 8vo. 2s. 6d. KEY TO PART II. (XXV.—C.) for Teachers only. Gl. 8vo. 3s. 6d.
- \*AN INTRODUCTION TO LATIN LYRIC VERSE COMPOSITION. By the same. Gl. 8vo. 3s. KEY, for Teachers only. Gl. 8vo. 4s. 6d.
- \***MACMILLAN.**—FIRST LATIN GRAMMAR. By M. C. MACMILLAN, M.A. Fcap. 8vo. 1s. 6d.
- MACMILLAN'S LATIN COURSE.** Globe 8vo.
- \***FIRST PART.** By A. M. COOK, M.A., Assistant Master at St. Paul's School. Gl. 8vo. 3s. 6d.
- \***SECOND PART.** By A. M. COOK, M.A., and W. E. P. PANTIN, M.A. New and Enlarged Edition. Gl. 8vo. 4s. 6d.
- \***MACMILLAN'S SHORTER LATIN COURSE.** Gl. 8vo.
- FIRST PART.** By A. M. COOK, M.A. 1s. 6d. KEY, for Teachers only. 4s. 6d.
- SECOND PART.** By A. M. COOK, M.A., and W. E. P. PANTIN, M.A. *[In January. Key shortly.]*
- \***MACMILLAN'S LATIN READER.**—A LATIN READER FOR THE LOWER FORMS IN SCHOOLS. By H. J. HARDY, M.A., Assistant Master at Winchester. Gl. 8vo. 2s. 6d.
- NALL.**—A LATIN-ENGLISH DICTIONARY. By Rev. G. H. NALL. *[In preparation.]*
- NIXON.**—PARALLEL EXTRACTS, Arranged for Translation into English and Latin, with Notes on Idioms. By J. E. NIXON, M.A., Fellow and Classical Lecturer, King's College, Cambridge. Part I.—Historical and Epistolary. Cr. 8vo. 3s. 6d.
- PROSE EXTRACTS, Arranged for Translation into English and Latin, with General and Special Prefaces on Style and Idiom. By the same. I. Oratorical. II. Historical. III. Philosophical. IV. Anecdotes and Letters. 2nd Ed., enlarged to 280 pp. Cr. 8vo. 4s. 6d. SELECTIONS FROM THE SAME. 2s. 6d. Translations of about 70 Extracts can be supplied to Schoolmasters (2s. 6d.), on application to the Author: and about 40 similarly of "Parallel Extracts." 1s. 6d. post free.
- NIXON—SMITH.**—PARALLEL VERSE EXTRACTS FOR TRANSLATION INTO ENGLISH AND LATIN. By J. E. NIXON, M.A., and E. H. C. SMITH, M.A. Cr. 8vo. 5s. 6d.
- \***PANTIN.**—A FIRST LATIN VERSE BOOK. By W. E. P. PANTIN, M.A. Assistant Master at St. Paul's School. Gl. 8vo. 1s. 6d. KEY, for Teachers only. 4s. net.
- \***PEILE.**—A PRIMER OF PHILOLOGY. By J. PEILE, Litt.D., Master of Christ's College, Cambridge. 18mo. 1s.



- \***POSTGATE**.—**SERMO LATINUS**. A short Guide to Latin Prose Composition. By Prof. J. P. POSTGATE, Litt.D., Fellow of Trinity College, Cambridge. Gl. 8vo. 2s. 6d. KEY to "Selected Passages." Gl. 8vo. 4s. 6d. net.
- POTTS**.—\***HINTS TOWARDS LATIN PROSE COMPOSITION**. By A. W. POTTS, M.A., LL.D., late Fellow of St. John's College, Cambridge. Ex. fcap. 8vo. 3s.
- \***PASSAGES FOR TRANSLATION INTO LATIN PROSE**. Edited with Notes and References to the above. Ex. fcap. 8vo. 2s. 6d. KEY, for Teachers only. 2s. 6d.
- \***PRESTON**.—**EXERCISES IN LATIN VERSE OF VARIOUS KINDS**. By Rev. G. PRESTON. Gl. 8vo. 2s. 6d. KEY, for Teachers only. Gl. 8vo. 5s.
- REID**.—**A GRAMMAR OF TACITUS**. By J. S. REID, Litt.D., Fellow of Caius College, Cambridge. *[In preparation.]*  
A GRAMMAR OF VIRGIL. By the same. *[In preparation.]*
- ROBY**.—Works by H. J. ROBY, M.A., late Fellow of St. John's College, Cambridge.  
A GRAMMAR OF THE LATIN LANGUAGE, from Plautus to Suetonius. Part I. Sounds, Inflections, Word-formation, Appendices. Cr. 8vo. 9s. Part II. Syntax, Prepositions, etc. 10s. 6d.
- \***SCHOOL LATIN GRAMMAR**. Cr. 8vo. 5s.
- \***ROBY—WILKINS**. **AN ELEMENTARY LATIN GRAMMAR**. By H. J. ROBY, M.A., and Prof. A. S. WILKINS, Litt.D. Gl. 8vo. 2s. 6d.
- \***RUST**.—**FIRST STEPS TO LATIN PROSE COMPOSITION**. By Rev. G. RUST M.A. 18mo. 1s. 6d. KEY, for Teachers only. By W. M. YATES. 18mo. 3s. 6d.
- \***SIMPSON**.—**LATIN PROSE AFTER THE BEST AUTHORS: Cæsar's Prose**. By F. P. SIMPSON, B.A. Ex. fcap. 8vo. 2s. 6d. KEY, for Teachers only. 5s.
- SMITH**.—**THE THEORY OF CONDITIONAL SENTENCES IN LATIN AND GREEK**. By R. HORTON SMITH, Q.C. 8vo. *[In the Press.]*
- STRACHAN—WILKINS**.—**ANALECTA**. Selected Passages for Translation. By J. S. STRACHAN, M.A., Professor of Greek, and A. S. WILKINS, Litt.D., Professor of Latin, Owens College, Manchester. Cr. 8vo. 5s. Also in two parts, 2s. 6d. each. Indexes to Greek and Latin passages, 6d. each.
- THRING**.—**A LATIN GRADUAL**. By the Rev. E. THRING, M.A., late Headmaster of Uppingham. A First Latin Construing Book. Fcap. 8vo. 2s. 6d.
- A **MANUAL OF MOOD CONSTRUCTIONS**. Fcap. 8vo. 1s. 6d.
- WELCH—DUFFIELD**.—\***LATIN ACCIDENCE AND EXERCISES ARRANGED FOR BEGINNERS**. By W. WELCH and C. G. DUFFIELD. 18mo. 1s. 6d.
- \***EXERCISES IN UNSEEN TRANSLATION IN LATIN**. By the same. 18mo. 1s. 6d.
- WRIGHT**.—Works by J. WRIGHT, M.A., late Headmaster of Sutton Coldfield School.  
A **HELP TO LATIN GRAMMAR**; or, the Form and Use of Words in Latin, with Progressive Exercises. Cr. 8vo. 4s. 6d.
- THE SEVEN KINGS OF ROME**. An Easy Narrative, abridged from the First Book of Livy by the omission of Difficult Passages; being a First Latin Reading Book, with Grammatical Notes and Vocabulary. Fcap. 8vo. 3s. 6d.
- FIRST LATIN STEPS; or, AN INTRODUCTION BY A SERIES OF EXAMPLES TO THE STUDY OF THE LATIN LANGUAGE**. Cr. 8vo. 3s.
- A **COMPLETE LATIN COURSE**, comprising Rules with Examples, Exercises, both Latin and English, on each Rule, and Vocabularies. Cr. 8vo. 2s. 6d.

### Greek.

- BLACKIE**.—**GREEK AND ENGLISH DIALOGUES FOR USE IN SCHOOLS AND COLLEGES**. By JOHN STUART BLACKIE, Emeritus Professor of Greek in the University of Edinburgh. New Edition. Fcap. 8vo. 2s. 6d.
- A **GREEK PRIMER, COLLOQUIAL AND CONSTRUCTIVE**. Cr. 8vo. 2s. 6d.
- BRYANS**.—**GREEK PROSE EXERCISES** based upon Thucydides. By C. BRYANS, M.A. *[In preparation.]*
- GILES**.—See under Latin.
- GOODWIN**.—Works by W. W. GOODWIN, LL.D., D.C.L., Professor of Greek in Harvard University.  
**SYNTAX OF THE MOODS AND TENSES OF THE GREEK VERB**. New Ed., revised and enlarged. 8vo. 14s.
- \***A GREEK GRAMMAR**. Cr. 8vo. 6s.
- \***A GREEK GRAMMAR FOR SCHOOLS**. New Edition. Cr. 8vo. 3s. 6d.
- HADLEY**.—See under Latin.

- HADLEY—ALLEN.**—A GREEK GRAMMAR FOR SCHOOLS AND COLLEGES. By JAMES HADLEY, late Professor in Yale College. Revised by F. DE F. ALLEN, Professor in Harvard College. Cr. 8vo. 6s.
- \***JACKSON.**—FIRST STEPS TO GREEK PROSE COMPOSITION. By BLOMFIELD JACKSON, M.A. 18mo. 1s. 6d. KEY, for Teachers only. 18mo. 3s. 6d.
- \***SECOND STEPS TO GREEK PROSE COMPOSITION**, with Examination Papers. By the same. 18mo. 2s. 6d. KEY, for Teachers only. 18mo. 3s. 6d.
- KYNASTON.**—EXERCISES IN THE COMPOSITION OF GREEK IAMBIC VERSE. By Rev. H. KYNASTON, D.D., Professor of Classics in the University of Durham. With Vocabulary. Ex. fcap. 8vo. 5s. KEY, for Teachers only. Ex. fcap. 8vo. 4s. 6d.
- MACKIE.**—PARALLEL PASSAGES FOR TRANSLATION INTO GREEK AND ENGLISH. With Indexes. By Rev. E. C. MACKIE, M.A., Classical Master at Heversham Grammar School. Gl. 8vo. 4s. 6d.
- MACMILLAN'S GREEK COURSE.**—Edited by Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. Gl. 8vo.
- \***FIRST GREEK GRAMMAR—ACCIDENCE.** By the Editor. 2s.
- \***FIRST GREEK GRAMMAR—SYNTAX.** By the same. 2s.
- ACCIDENCE AND SYNTAX. In one volume. 3s. 6d.
- \***EASY EXERCISES IN GREEK ACCIDENCE.** By H. G. UNDERHILL, M.A., Assistant Master at St. Paul's Preparatory School. 2s.
- \***A SECOND GREEK EXERCISE BOOK.** By Rev. W. A. HEARD, M.A., Headmaster of Fettes College, Edinburgh. 2s. 6d.
- \***EASY EXERCISES IN GREEK SYNTAX.** By Rev. G. H. NALL, M.A., Assistant Master at Westminster School. 2s. 6d.
- MANUAL OF GREEK ACCIDENCE. By the Editor. [In preparation.]
- MANUAL OF GREEK SYNTAX. By the Editor. [In preparation.]
- ELEMENTARY GREEK COMPOSITION. By the Editor. [In preparation.]
- \***MACMILLAN'S GREEK READER.**—STORIES AND LEGENDS. A First Greek Reader, with Notes, Vocabulary, and Exercises. By F. H. COLSON, M.A., Headmaster of Plymouth College. Gl. 8vo. 3s.
- \***MARSHALL.**—A TABLE OF IRREGULAR GREEK VERBS, classified according to the arrangement of Curtius's Greek Grammar. By J. M. MARSHALL, M.A., Headmaster of the Grammar School, Durham. 8vo. 1s.
- MAYOR.**—FIRST GREEK READER. By Prof. JOHN E. B. MAYOR, M.A., Fellow of St. John's College, Cambridge. Fcap. 8vo. 4s. 6d.
- MAYOR.**—GREEK FOR BEGINNERS. By Rev. J. B. MAYOR, M.A., late Professor of Classical Literature in King's College, London. Part I., with Vocabulary, 1s. 6d. Parts II. and III., with Vocabulary and Index. Fcap. 8vo. 3s. 6d. Complete in one Vol. 4s. 6d.
- NALL.**—A SHORT GREEK-ENGLISH DICTIONARY. By Rev. G. H. NALL. [In preparation.]
- PEILE.**—See under Latin.
- RUTHERFORD.**—THE NEW PHRYNICHUS; being a Revised Text of the Ecloga of the Grammarian Phrynichus. With Introduction and Commentary. By the Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. 8vo. 18s.
- STRACHAN—WILKINS.**—See under Latin.
- WHITE.**—FIRST LESSONS IN GREEK. Adapted to GOODWIN'S GREEK GRAMMAR, and designed as an introduction to the ANABASIS OF XENOPHON. By JOHN WILLIAMS WHITE, Assistant Professor of Greek in Harvard University, U.S.A. Cr. 8vo. 3s. 6d.
- WRIGHT.**—ATTIC PRIMER. Arranged for the Use of Beginners. By J. WRIGHT, M.A. Ex. fcap. 8vo. 2s. 6d.

## ANTIQUITIES, ANCIENT HISTORY, AND PHILOSOPHY.

- ARNOLD.**—A HISTORY OF THE EARLY ROMAN EMPIRE. By W. T. ARNOLD, M.A. Cr. 8vo. [In preparation.]
- ARNOLD.**—THE SECOND PUNIC WAR. Being Chapters from THE HISTORY OF ROME by the late THOMAS ARNOLD, D.D., Headmaster of Rugby. Edited, with Notes, by W. T. ARNOLD, M.A. With 8 Maps. Cr. 8vo. 5s.

- \***BEESELY**.—STORIES FROM THE HISTORY OF ROME. By Mrs. BEESELY. Fcap. 8vo. 2s. 6d.
- BLACKIE**.—HORÆ HELLENICÆ. By JOHN STUART BLACKIE, Emeritus Professor of Greek in the University of Edinburgh. 8vo. 12s.
- BURN**.—ROMAN LITERATURE IN RELATION TO ROMAN ART. By Rev. ROBERT BURN, M.A., late Fellow of Trinity College, Cambridge. Illustrated. Ex. cr. 8vo. 14s.
- BURY**.—A HISTORY OF THE LATER ROMAN EMPIRE FROM ARCADIUS TO IRENE, A.D. 395-800. By J. B. BURY, M.A., Fellow of Trinity College, Dublin. 2 vols. 8vo. 32s.
- A SCHOOL HISTORY OF GREECE. By the same. Cr. 8vo. [*In preparation.*]
- BUTCHER**.—SOME ASPECTS OF THE GREEK GENIUS. By S. H. BUTCHER, M.A., Professor of Greek, Edinburgh. 2nd Ed. revised. Cr. 8vo. [*In the Press.*]
- \***CLASSICAL WRITERS**.—Edited by JOHN RICHARD GREEN, M.A., LL.D. Fcap. 8vo. 1s. 6d. each.
- SOPHOCLES. By Prof. LEWIS CAMPBELL, M.A.
- EURIPIDES. By Prof. MAHAFFY, D.D.
- DEMOSTHENES. By Prof. S. H. BUTCHER, M.A.
- VIRGIL. By Prof. NETTLESHIP, M.A.
- LIVY. By Rev. W. W. CAPES, M.A.
- TACITUS. By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A.
- MILTON. By Rev. STOPFORD A. BROOKE, M.A.
- DYER**.—STUDIES OF THE GODS IN GREECE AT CERTAIN SANCTUARIES RECENTLY EXCAVATED. By LOUIS DYER, B.A. Ex. Cr. 8vo. 8s. 6d. net.
- FOWLER**.—THE CITY-STATE OF THE GREEKS AND ROMANS. By W. WARDE FOWLER, M.A. Cr. 8vo. 5s.
- FREEMAN**.—HISTORICAL ESSAYS. By the late EDWARD A. FREEMAN, D.C.L., LL.D. Second Series. [Greek and Roman History.] 8vo. 10s. 6d.
- HISTORY OF THE FEDERAL GOVERNMENT IN GREECE AND ITALY. With a General Introduction. New Edition. Ed. by J. B. BURY, M.A. Ex. Cr. 8vo. 12s. 6d.
- GARDNER**.—SAMOS AND SAMIAN COINS. An Essay. By PERCY GARDNER, Litt.D., Professor of Archæology in the University of Oxford. 8vo. 7s. 6d.
- GEDDES**.—THE PROBLEM OF THE HOMERIC POEMS. By Sir W. D. GEDDES, Principal of the University of Aberdeen. 8vo. 14s.
- GLADSTONE**.—Works by the Rt. Hon. W. E. GLADSTONE, M.P.
- THE TIME AND PLACE OF HOMER. Cr. 8vo. 6s. 6d.
- LANDMARKS OF HOMERIC STUDY. Cr. 8vo. 2s. 6d.
- \*A PRIMER OF HOMER. 18mo. 1s.
- GOW**.—A COMPANION TO SCHOOL CLASSICS. By JAMES GOW, Litt.D., Head Master of the High School, Nottingham. Illustrated. Cr. 8vo. 6s.
- HARRISON-VERRALL**.—MYTHOLOGY AND MONUMENTS OF ANCIENT ATHENS. Translation of a portion of the "Attica" of Pausanias. By MARGARET DE G. VERRALL. With Introductory Essay and Archæological Commentary by JANE E. HARRISON. Illustrated. Cr. 8vo. 16s.
- HOLM**.—HISTORY OF GREECE. By Professor A. HOLM. Translated. 4 vols. [*In preparation.*]
- JEBB**.—Works by R. C. JEBB, Litt.D., Professor of Greek in the University of Cambridge.
- THE ATTIC ORATORS FROM ANTIPHON TO ISAEUS. 2 vols. 2nd Ed. 8vo. 25s.
- \*A PRIMER OF GREEK LITERATURE. 18mo. 1s.
- LECTURES ON GREEK POETRY. Cr. 8vo. [Dec. 1893.]
- KIEPERT**.—MANUAL OF ANCIENT GEOGRAPHY. By Dr. H. KIEPERT. Cr. 8vo. 5s.
- LANCIANI**.—ANCIENT ROME IN THE LIGHT OF RECENT DISCOVERIES. By RODOLFO LANCIANI, Professor of Archæology in the University of Rome. Illustrated. 4to. 24s.
- PAGAN AND CHRISTIAN ROME. By the same. Illustrated. 4to. 24s.
- LEAF**.—COMPANION TO THE ILIAD FOR ENGLISH READERS. By WALTER LEAF, Litt.D. Cr. 8vo. 7s. 6d.
- MAHAFFY**.—Works by J. P. MAHAFFY, D.D., Fellow of Trinity College, Dublin, and Professor of Ancient History in the University of Dublin.



- SOCIAL LIFE IN GREECE; from Homer to Menander. Cr. 8vo. 9s.  
 GREEK LIFE AND THOUGHT; from the Age of Alexander to the Roman Conquest. Cr. 8vo. 12s. 6d.  
 THE GREEK WORLD UNDER ROMAN SWAY. From Plutarch to Polybius. Cr. 8vo. 10s. 6d.  
 PROBLEMS IN GREEK HISTORY. Cr. 8vo. 7s. 6d.  
 RAMBLES AND STUDIES IN GREECE. 4th Ed. Illust. Cr. 8vo. 10s. 6d.  
 A HISTORY OF CLASSICAL GREEK LITERATURE. Cr. 8vo. Vol. I. The Poets. Part I. Epic and Lyric. Part II. Dramatic. Vol. II. Prose Writers. Part I. Herodotus to Plato. Part II. Isocrates to Aristotle. 4s. 6d. each Part.  
 \*A PRIMER OF GREEK ANTIQUITIES. With Illustrations. 18mo. 1s.  
 MAYOR.—BIBLIOGRAPHICAL CLUE TO LATIN LITERATURE. Edited after HÜBNER. By Prof. JOHN E. B. MAYOR. Cr. 8vo. 10s. 6d.  
 NEWTON.—ESSAYS ON ART AND ARCHÆOLOGY. By Sir CHARLES NEWTON, K.C.B., D.C.L. 8vo. 12s. 6d.  
 PATER.—PLATO AND PLATONISM. By WALTER PATER, M.A., Fellow of Brasenose College, Oxford. Ex. Cr. 8vo. 8s. 6d.  
 PHILOLOGY.—THE JOURNAL OF PHILOLOGY. Edited by W. A. WRIGHT, M.A., I. BYWATER, M.A., and H. JACKSON, Litt.D. 4s. 6d. each (half-yearly).  
 SCHMIDT—WHITE. AN INTRODUCTION TO THE RHYTHMIC AND METRIC OF THE CLASSICAL LANGUAGES. By Dr. J. H. H. SCHMIDT. Translated by JOHN WILLIAMS WHITE, Ph.D. 8vo. 10s. 6d.  
 SCHREIBER—ANDERSON.—ATLAS OF CLASSICAL ARCHÆOLOGY. By TH. SCHREIBER, with English Text by Prof. W. C. F. ANDERSON. [*In the Press.*]  
 SCHUCHHARDT.—DR. SCHLIEHMANN'S EXCAVATIONS AT TROY, TIRYNS, MYCENÆ, ORCHOMENOS, ITHACA, presented in the light of recent knowledge. By Dr. CARL SCHUCHHARDT. Translated by EUGENIE SELLERS. Introduction by WALTER LEAF, Litt.D. Illustrated. 8vo. 18s. net.  
 SHUCKBURGH.—A SCHOOL HISTORY OF ROME. By E. S. SHUCKBURGH, M.A. Cr. 8vo. [*In the Press.*]  
 SMITH.—A HANDBOOK ON GREEK PAINTING. By CECIL SMITH. [*In prep.*]  
 \*STEWART.—THE TALE OF TROY. Done into English by AUBREY STEWART. Gl. 8vo. 3s. 6d. [18mo. 1s.]  
 \*TOZER.—A PRIMER OF CLASSICAL GEOGRAPHY. By H. F. TOZER, M.A.  
 WILKINS.—Works by Prof. WILKINS, Litt.D., LL.D.  
 \*A PRIMER OF ROMAN ANTIQUITIES. Illustrated. 18mo. 1s.  
 \*A PRIMER OF ROMAN LITERATURE. 18mo. 1s.  
 WILKINS—ARNOLD.—A MANUAL OF ROMAN ANTIQUITIES. By Prof. A. S. WILKINS, Litt.D., and W. T. ARNOLD, M.A. Cr. 8vo. [*In prep.*]

## MODERN LANGUAGES AND LITERATURE.

English; French; German; Modern Greek; Italian; Spanish.

### ENGLISH.

- \*ABBOTT.—A SHAKESPEARIAN GRAMMAR. An Attempt to Illustrate some of the Differences between Elizabethan and Modern English. By the Rev. E. A. ABBOTT, D.D., formerly Headmaster of the City of London School. Ex. fcap. 8vo. 6s.  
 \*ADDISON.—SELECTIONS FROM "THE SPECTATOR." With Introduction and Notes, by K. DEIGHTON. Gl. 8vo. 2s. 6d.  
 \*BACON.—ESSAYS. With Introduction and Notes, by F. G. SELBY, M.A., Principal and Professor of Logic and Moral Philosophy, Deccan College, Poona. Gl. 8vo. 3s.; sewed, 2s. 6d.  
 \*THE ADVANCEMENT OF LEARNING. By the same. Gl. 8vo. Book I. 2s. Book II. 3s. 6d.  
 BATES.—AN OUTLINE OF THE DEVELOPMENT OF THE EARLY ENGLISH DRAMA. By K. L. BATES. [*In the Press.*]  
 BROOKE.—EARLY ENGLISH LITERATURE. By Rev. STOPFORD A. BROOKE, M.A. 2 vols. 8vo. 20s. net.

- BROWNING.**—A PRIMER ON BROWNING. By F. M. WILSON. Gl. 8vo. 2s. 6d.
- \***BURKE.**—REFLECTIONS ON THE FRENCH REVOLUTION. By F. G. SELBY, M.A. Gl. 8vo. 5s.
- SPEECH ON CONCILIATION WITH AMERICA, ON AMERICAN TAXATION. LETTER TO THE SHERIFF OF BRISTOL. By the same. [*In the Press.*]
- BUTLER.**—HUDIBRAS. With Introduction and Notes, by ALFRED MILNES, M.A. Ex. fcap. 8vo. Part I. 3s. 6d. Parts II. and III. 4s. 6d.
- CAMPBELL.**—SELECTIONS. With Introduction and Notes, by CECIL M. BARROW, M.A., Principal of Victoria College, Palghat. Gl. 8vo. [*In preparation.*]
- CHAUCER.**—A PRIMER OF CHAUCER. By A. W. POLLARD, M.A. 18mo. 1s.
- COLLINS.**—THE STUDY OF ENGLISH LITERATURE: A Plea for its Recognition at the Universities. By J. CHURTON COLLINS, M.A. Cr. 8vo. 4s. 6d.
- COWPER.**—\*THE TASK: an Epistle to Joseph Hill, Esq.; TIROCINIUM, or a Review of the Schools; and THE HISTORY OF JOHN GILPIN. Edited, with Notes, by W. BENHAM B.D. Gl. 8vo. 1s.
- THE TASK. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A. [BOOK IV. *in preparation.*]
- CRAIK.**—ENGLISH PROSE SELECTIONS. With Critical Introductions by various writers, and General Introductions to each Period. Edited by HENRY CRAIK, C.B., LL.D. In 5 vols. Vol. I. 14th to 16th Century. Cr. 8vo. 7s. 6d. [Vol. II. *in the Press.*]
- DRYDEN.**—SELECT PROSE WORKS. Edited, with Introduction and Notes, by Prof. C. D. YONGE. Fcap. 8vo. 2s. 6d.
- SELECT SATIRES. With Introduction and Notes. By J. CHURTON COLLINS, M.A. Gl. 8vo. [Nov. 1893.]
- \***GLOBE READERS.** Edited by A. F. MURISON. Illustrated. Gl. 8vo.
- Primer I. (48 pp.) 3d. Primer II. (48 pp.) 3d. Book I. (132 pp.) 8d. Book II. (136 pp.) 10d. Book III. (232 pp.) 1s. 3d. Book IV. (328 pp.) 1s. 9d. Book V. (408 pp.) 2s. Book VI. (436 pp.) 2s. 6d.
- \***THE SHORTER GLOBE READERS.**—Illustrated. Gl. 8vo.
- Primer I. (48 pp.) 3d. Primer II. (48 pp.) 3d. Book I. (132 pp.) 8d. Book II. (136 pp.) 10d. Book III. (178 pp.) 1s. Book IV. (182 pp.) 1s. Book V. (216 pp.) 1s. 3d. Book VI. (228 pp.) 1s. 6d.
- \***GOLDSMITH.**—THE TRAVELLER, or a Prospect of Society; and THE DESERTED VILLAGE. With Notes, Philological and Explanatory, by J. W. HALES, M.A. Cr. 8vo. 6d.
- \***THE TRAVELLER AND THE DESERTED VILLAGE.** With Introduction and Notes, by A. BARRETT, B.A., Professor of English Literature, Elphinstone College, Bombay. Gl. 8vo. 1s. 9d. Separately, 1s. each, sewed.
- \***THE VICAR OF WAKEFIELD.** With a Memoir of Goldsmith, by Prof. MASSON. Gl. 8vo. 1s.
- SELECT ESSAYS. With Introduction and Notes, by Prof. C. D. YONGE. Fcap. 8vo. 2s. 6d.
- GOW.**—A METHOD OF ENGLISH, for Secondary Schools. Part I. By JAMES GOW, Litt.D. Gl. 8vo. 2s.
- GOYEN.**—THE PRINCIPLES OF ENGLISH COMPOSITION THROUGH ANALYSIS AND SYNTHESIS. By P. GOYEN, Inspector of Schools, New Zealand. Gl. 8vo. [*In the Press.*]
- \***GRAY.**—POEMS. With Introduction and Notes, by JOHN BRADSHAW, LL.D. Gl. 8vo. 1s. 9d.
- \***HALES.**—Works by J. W. HALES, M.A., Professor of English Literature at King's College, London.
- LONGER ENGLISH POEMS. With Notes, Philological and Explanatory, and an Introduction on the Teaching of English. Ex. fcap. 8vo. 4s. 6d.
- SHORTER ENGLISH POEMS. Ex. fcap. 8vo. [*In preparation.*]
- \***HELPS.**—ESSAYS WRITTEN IN THE INTERVALS OF BUSINESS. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A. Gl. 8vo. 1s. 9d.
- \***JOHNSON.**—LIVES OF THE POETS. The Six Chief Lives (Milton, Dryden, Swift, Addison, Pope, Gray), with Macaulay's "Life of Johnson." With Preface and Notes by MATTHEW ARNOLD. Cr. 8vo. 4s. 6d.
- \***LIFE OF MILTON.** With Introduction and Notes, by K. DEIGHTON. Globe 8vo. 1s. 9d.

- KELLNER.**—HISTORICAL OUTLINES OF ENGLISH SYNTAX. By L. KELLNER, Ph.D. Globe 8vo. 6s.
- KINGSLEY.**—WESTWARD HO! By CHARLES KINGSLEY. Abridged Edition for Schools. Gl. 8vo. 1s. 6d.
- \***HERWARD THE WAKE.** By the same. Gl. 8vo. 1s. 6d.
- LAMB.**—TALES FROM SHAKESPEARE. With Introduction and Notes by Rev. A. AINGER, LL.D., Canon of Bristol. 18mo. 2s. 6d. net.
- \***LITERATURE PRIMERS.**—Edited by J. R. GREEN, LL.D. 18mo. 1s. each.
- ENGLISH GRAMMAR. By Rev. R. MORRIS, LL.D.
- ENGLISH GRAMMAR EXERCISES. By R. MORRIS, LL.D., and H. C. BOWEN, M.A.
- EXERCISES ON MORRIS'S PRIMER OF ENGLISH GRAMMAR. By J. WETHERELL, M.A.
- ENGLISH COMPOSITION. By Professor JOHN NICHOL.
- QUESTIONS AND EXERCISES ON ENGLISH COMPOSITION. By Prof. NICHOL and W. S. M'CORMICK.
- ENGLISH LITERATURE. By STOPFORD BROOKE, M.A.
- SHAKSPERE. By Professor DOWDEN.
- CHAUCER. By A. W. POLLARD, M.A.
- SPENSER. By Prof. J. W. HALES.
- THE CHILDREN'S TREASURY OF LYRICAL POETRY. Selected and arranged with Notes by FRANCIS TURNER PALGRAVE. In Two Parts. 1s. each.
- PHILOLOGY. By J. PEILE, Litt.D.
- ROMAN LITERATURE. By Prof. A. S. WILKINS, Litt.D.
- GREEK LITERATURE. By Prof. JEBB, Litt.D.
- HOMER. By the Rt. Hon. W. E. GLADSTONE, M.P.
- A HISTORY OF ENGLISH LITERATURE IN FOUR VOLUMES.** Cr. 8vo.
- EARLY ENGLISH LITERATURE. By STOPFORD BROOKE, M.A. [*In preparation.*]
- ELIZABETHAN LITERATURE. (1560-1665.) By GEORGE SAINTSBURY. 7s. 6d.
- EIGHTEENTH CENTURY LITERATURE. (1660-1780.) By EDMUND GOSSE, M.A. 7s. 6d.
- THE MODERN PERIOD. By Prof. DOWDEN. [*In preparation.*]
- LITTLEDALE.**—ESSAYS ON TENNYSON'S IDYLLS OF THE KING. By H. LITTLEDALE, M.A., Vice-Principal and Professor of English Literature, Baroda College. Cr. 8vo. 4s. 6d.
- MACAULAY.**—\*ESSAY ON LORD CLIVE. With Introduction and Notes by K. DEIGHTON. Gl. 8vo. 2s.
- \*ESSAY ON WARREN HASTINGS. By the same. Gl. 8vo. 2s. 6d.
- ESSAY ON ADDISON. By Prof. J. W. HALES, M.A. Gl. 8vo. [*In preparation.*]
- MACLEAN.**—ZUPITZA'S OLD AND MIDDLE ENGLISH READER. With Notes and Vocabulary by Prof. G. E. MACLEAN. [*In the Press.*]
- \***MACMILLAN'S HISTORY READERS.** (See History, p. 43.)
- \***MACMILLAN'S READING BOOKS.**
- PRIMER. 18mo. (48 pp.) 2d. BOOK I. (96 pp.) 4d. BOOK II. (144 pp.) 5d. BOOK III. (160 pp.) 6d. BOOK IV. (176 pp.) 8d. BOOK V. (380 pp.) 1s. BOOK VI. Cr. 8vo. (430 pp.) 2s.
- Book VI. is fitted for Higher Classes, and as an Introduction to English Literature.
- MACMILLAN'S RECITATION CARDS.** Selections from TENNYSON, KINGSLEY, MATTHEW ARNOLD, CHRISTINA ROSSETTI, DOYLE. Annotated. Cr. 8vo. Nos. 1 to 18, 1d. each; Nos. 19 to 42, 2d. each.
- \***MACMILLAN'S COPY BOOKS.**—1. Large Post 4to. Price 4d. each. 2. Post Oblong. Price 2d. each.
- Nos. 3, 4, 5, 6, 7, 8, 9 may be had with *Goodman's Patent Sliding Copies.* Large Post 4to. Price 6d. each.
- MARTIN.**—\*THE POET'S HOUR: Poetry selected for Children. By FRANCES MARTIN. 18mo. 2s. 6d.
- \***SPRING-TIME WITH THE POETS.** By the same. 18mo. 3s. 6d.
- \***MILTON.**—PARADISE LOST. Books I. and II. With Introduction and Notes, by MICHAEL MACMILLAN, B.A., Professor of English Literature, Elphinstone College, Bombay. Gl. 8vo. 1s. 9d. Or separately, 1s. 3d.; sewed 1s. each.



- \***L'ALLEGRO, IL PENNEROSO, LYCIDAS, ARCADES, SONNETS, &c.** With Introduction and Notes, by W. BELL, M.A., Professor of Philosophy and Logic, Government College, Lahore. Gl. 8vo. 1s. 9d.
- \***COMUS.** By the same. Gl. 8vo. 1s. 3d.
- \***SAMSON AGONISTES.** By H. M. PERCIVAL, M.A., Professor of English Literature, Presidency College, Calcutta. Gl. 8vo. 2s.
- \***INTRODUCTION TO THE STUDY OF MILTON.** By STOPFORD BROOKE, M.A. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)
- MORRIS.**—Works by the Rev. R. MORRIS, LL.D.
- \***A PRIMER OF ENGLISH GRAMMAR.** 18mo. 1s.
- \***ELEMENTARY LESSONS IN HISTORICAL ENGLISH GRAMMAR,** containing Accidence and Word-Formation. 18mo. 2s. 6d.
- \***HISTORICAL OUTLINES OF ENGLISH ACCIDENCE,** with Chapters on the Development of the Language, and on Word-Formation. Ex. fcap. 8vo. 6s.
- NICHOL—M'CORMICK.**—**A SHORT HISTORY OF ENGLISH LITERATURE.** By Prof. JOHN NICHOL and Prof. W. S. M'CORMICK. [*In preparation.*]
- OLIPHANT.**—**THE LITERARY HISTORY OF ENGLAND, 1790–1825.** By Mrs. OLIPHANT. 3 vols. 8vo. 21s.
- OLIPHANT.**—**THE OLD AND MIDDLE ENGLISH.** By T. L. KINGTON OLIPHANT. 2nd Ed. Gl. 8vo. 9s.
- THE NEW ENGLISH.** By the same. 2 vols. Cr. 8vo. 21s.
- PALGRAVE.**—**THE GOLDEN TREASURY OF SONGS AND LYRICS.** Selected by F. T. PALGRAVE. 18mo. 2s. 6d. net.
- \***THE CHILDREN'S TREASURY OF LYRICAL POETRY.** Selected by the same. 18mo. 2s. 6d. net. Also in Two Parts. 1s. each.
- PATMORE.**—**THE CHILDREN'S GARLAND FROM THE BEST POETS.** Selected by COVENTRY PATMORE. 18mo. 2s. 6d. net.
- \***RANSOME.**—**SHORT STUDIES OF SHAKESPEARE'S PLOTS.** By CYRIL RANSOME, M.A., Professor of Modern History and Literature, Yorkshire College, Leeds. Cr. 8vo. 3s. 6d. Also **HAMLET, MACBETH, THE TEMPEST,** 9d. each, sewed.
- \***RYLAND.**—**CHRONOLOGICAL OUTLINES OF ENGLISH LITERATURE.** By F. RYLAND, M.A. Cr. 8vo. 6s.
- SCOTT.**—\***LAY OF THE LAST MINSTREL, and THE LADY OF THE LAKE.** Edited by FRANCIS TURNER PALGRAVE. Gl. 8vo. 1s.
- \***THE LAY OF THE LAST MINSTREL.** With Introduction and Notes, by G. H. STUART, M.A., Principal of Kumbakonam College, and E. H. ELLIOT, B.A. Gl. 8vo. 2s. Canto I. 9d. Cantos I. to III. and IV. to VI. Sewed, 1s. each.
- \***MARMION, and THE LORD OF THE ISLES.** By F. T. PALGRAVE. Gl. 8vo. 1s.
- \***MARMION.** With Introduction and Notes, by MICHAEL MACMILLAN, B.A. Gl. 8vo. 3s.; sewed, 2s. 6d.
- \***THE LADY OF THE LAKE.** By G. H. STUART, M.A. Gl. 8vo. 2s. 6d.; sewed, 2s.
- \***ROKEBY.** With Introduction and Notes, by MICHAEL MACMILLAN, B.A. Gl. 8vo. 3s.; sewed, 2s. 6d.
- SHAKESPEARE.**—\***A SHAKESPEARIAN GRAMMAR.** (*See ABBOTT.*)
- \***A PRIMER OF SHAKESPEARE.** By Prof. DOWDEN. 18mo. 1s.
- \***SHORT STUDIES OF SHAKESPEARE'S PLOTS.** (*See RANSOME.*)
- \***THE TEMPEST.** With Introduction and Notes, by K. DEIGHTON. Gl. 8vo. 1s. 9d.
- \***MUCH ADO ABOUT NOTHING.** By the same. 2s.
- \***A MIDSUMMER NIGHT'S DREAM.** By the same. 1s. 9d.
- \***THE MERCHANT OF VENICE.** By the same. 1s. 9d.
- \***AS YOU LIKE IT.** By the same. 1s. 9d.
- \***TWELFTH NIGHT.** By the same. 1s. 9d.
- \***THE WINTER'S TALE.** By the same. 2s.
- \***KING JOHN.** By the same. 1s. 9d.
- \***RICHARD II.** By the same. 1s. 9d.
- \***HENRY IV.—PART I.** By the same. 2s. 6d.; sewed, 2s.
- \***HENRY IV.—PART II.** By the same. 2s. 6d.; sewed, 2s.
- \***HENRY V.** By the same. 1s. 9d.
- \***RICHARD III.** By C. H. TAWNEY, M.A. 2s. 6d.; sewed, 2s.
- \***CORIOLANUS.** By K. DEIGHTON. 2s. 6d.; sewed, 2s.

- \*ROMEO AND JULIET. By the same. 2s. 6d. ; sewed, 2s.  
 \*JULIUS CÆSAR. By the same. 1s. 9d.  
 \*MACBETH. By the same. 1s. 9d.  
 \*HAMLET. By the same. 2s. 6d. ; sewed, 2s.  
 \*KING LEAR. By the same. 1s. 9d.  
 \*OTHELLO. By the same. 2s.  
 \*ANTONY AND CLEOPATRA. By the same. 2s. 6d. ; sewed, 2s.  
 \*CYMBELINE. By the same. 2s. 6d. ; sewed, 2s.  
 \*SONNENSCHN—MEIKLEJOHN.—THE ENGLISH METHOD OF TEACHING TO READ. By A. SONNENSCHN and J. M. D. MEIKLEJOHN, M.A. Fcap. 8vo. THE NURSERY BOOK, containing all the Two-Letter Words in the Language. 1d. (Also in Large Type on Sheets for School Walls. 5s.) THE FIRST COURSE, consisting of Short Vowels with Single Consonants. 7d. THE SECOND COURSE, with Combinations and Bridges, consisting of Short Vowels with Double Consonants. 7d. THE THIRD AND FOURTH COURSES, consisting of Long Vowels, and all the Double Vowels in the Language. 7d.  
 \*SOUTHEY.—LIFE OF NELSON. With Introduction and Notes, by MICHAEL MACMILLAN, B.A. Gl. 8vo. 3s. ; sewed, 2s. 6d.  
 \*SPENSER.—THE FAIRIE QUEENE. Book I. With Introduction and Notes, by H. M. PERCIVAL, M.A. Gl. 8vo. 3s. ; sewed, 2s. 6d.  
 PRIMER OF SPENSER. By Prof. J. W. HALES. *[In the Press.]*  
 TAYLOR.—WORDS AND PLACES; or, Etymological Illustrations of History, Ethnology, and Geography. By Rev. ISAAC TAYLOR, Litt.D. Gl. 8vo. 6s.  
 TENNYSON.—THE COLLECTED WORKS. In 4 Parts. Cr. 8vo. 2s. 6d. each.  
 \*TENNYSON FOR THE YOUNG. Edited by the Rev. ALFRED AINGER, LL.D., Canon of Bristol. 18mo. 1s. net.  
 \*SELECTIONS FROM TENNYSON. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A. New Ed., enlarged. Gl. 8vo. 3s. 6d. or in two parts. Part I. 2s. 6d. Part II. 2s. 6d.  
 \*ENOCH ARDEN. By W. T. WEBB, M.A. Gl. 8vo. 2s. 6d.  
 \*AYLMER'S FIELD. By W. T. WEBB, M.A. 2s. 6d.  
 \*THE PRINCESS; A MEDLEY. By P. M. WALLACE, M.A. 3s. 6d.  
 \*THE COMING OF ARTHUR, and THE PASSING OF ARTHUR. By F. J. ROWE, M.A. Gl. 8vo. 2s. 6d.  
 \*GARETH AND LYNETTE. By G. C. MACAULAY, M.A. Globe 8vo. 2s. 6d.  
 \*GERAINT AND ENID, and THE MARRIAGE OF GERAINT. By G. C. MACAULAY, M.A. Gl. 8vo. 2s. 6d.  
 \*THE HOLY GRAIL. By G. C. MACAULAY, M.A. Globe 8vo. 2s. 6d.  
 LANCELOT AND ELAINE. By F. J. ROWE, M.A. *[In preparation.]*  
 GUINEVERE. By G. C. MACAULAY, M.A. *[In preparation.]*  
 THERING.—THE ELEMENTS OF GRAMMAR TAUGHT IN ENGLISH. By EDWARD THERING, M.A. With Questions. 4th Ed. 18mo. 2s.  
 \*VAUGHAN.—WORDS FROM THE POETS. By C. M. VAUGHAN. 18mo. 1s.  
 WARD.—THE ENGLISH POETS. Selections, with Critical Introductions by various Writers. Edited by T. H. WARD, M.A. 4 Vols. Vol. I. CHAUCER TO DONNE.—Vol. II. BEN JONSON TO DRYDEN.—Vol. III. ADDISON TO BLAKE.—Vol. IV. WORDSWORTH TO ROSSETTI. 2nd Ed. Cr. 8vo. 7s. 6d. each.  
 WARD.—A HISTORY OF ENGLISH DRAMATIC LITERATURE, TO THE DEATH OF QUEEN ANNE. By A. W. WARD, Litt.D., Principal of Owens College, Manchester. 2 vols. 8vo. *[New Ed. in preparation.]*  
 WOODS.—\*A FIRST POETRY BOOK. By M. A. WOODS. Fcap. 8vo. 2s. 6d.  
 \*A SECOND POETRY BOOK. By the same. 4s. 6d. ; or, Two Parts. 2s. 6d. each.  
 \*A THIRD POETRY BOOK. By the same. 4s. 6d.  
 HYMNS FOR SCHOOL WORSHIP. By the same. 18mo. 1s. 6d.  
 WORDSWORTH.—SELECTIONS. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A. Gl. 8vo. *[In preparation.]*  
 YONGE.—\*A BOOK OF GOLDEN DEEDS. By CHARLOTTE M. YONGE. 18mo. 2s. 6d. net.

## FRENCH.

- BEAUMARCHAIS.—LE BARBIER DE SEVILLE. With Introduction and Notes, by L. P. BLOUET. Fcap. 8vo. 3s. 6d.

- \*BOWEN.—FIRST LESSONS IN FRENCH.** By H. COURTHOPE BOWEN, M.A.  
Ex. fcap. 8vo. 1s.
- BREYMANN.—FIRST FRENCH EXERCISE BOOK.** By HERMANN BREYMANN,  
Ph.D., Professor of Philology in the University of Munich. Ex. fcap. 8vo. 4s. 6d.  
**SECOND FRENCH EXERCISE BOOK.** By the same. Ex. fcap. 8vo. 2s. 6d.
- DELBOS.—NAUTICAL TERMS.** By Prof. DELBOS, H.M.S. "Britannia."
- FASNACHT.—Works by G. E. FASNACHT, late Assistant Master at Westminster.**  
**THE ORGANIC METHOD OF STUDYING LANGUAGES.** Ex. fcap. 8vo. 1.  
French. 3s. 6d.  
**A FRENCH GRAMMAR FOR SCHOOLS.** Cr. 8vo. 3s. 6d.  
**GRAMMAR AND GLOSSARY OF THE FRENCH LANGUAGE OF THE  
SEVENTEENTH CENTURY.** Cr. 8vo. *[In preparation.]*  
**HANDBOOK TO FRENCH LITERATURE.** Cr. 8vo. *[In the Press.]*
- MACMILLAN'S PRIMARY SERIES OF FRENCH READING BOOKS.**—Edited by  
G. E. FASNACHT. Illustrations, Notes, Vocabulary, and Exercises. Gl. 8vo.
- \*FRENCH READINGS FOR CHILDREN.** By G. E. FASNACHT. 1s. 6d.
- \*CORNAZ—NOS ENFANTS ET LEURS AMIS.** By EDITH HARVEY. 1s. 6d.
- \*DE MAISTRE—LA JEUNE SIBERIENNE ET LE LÉPREUX DE LA CITÉ  
D'AOSTE.** By STEPHANE BARLET, B.Sc. 1s. 6d.
- \*FLORIAN—FABLES.** By Rev. CHARLES YELD, M.A., Headmaster of University  
School, Nottingham. 1s. 6d.
- \*LA FONTAINE—A SELECTION OF FABLES.** By L. M. MORIARTY, B.A.,  
Assistant Master at Harrow. 2s. 6d.
- \*MOLESWORTH—FRENCH LIFE IN LETTERS.** By Mrs. MOLESWORTH. 1s. 6d.
- \*PERRAULT—CONTES DE FÉES.** By G. E. FASNACHT. 1s. 6d.
- \*SOUVESTRE—UN PHILOSOPHE SOUS LES TOITS.** By L. M. MORIARTY,  
B.A. *[In the Press.]*
- MACMILLAN'S PROGRESSIVE FRENCH COURSE.**—By G. E. FASNACHT. Ex.  
fcap. 8vo.
- \*FIRST YEAR, Easy Lessons on the Regular Accidence.** 1s.
- \*SECOND YEAR, an Elementary Grammar with Exercises, Notes, and Vocabulary.** 2s.
- \*THIRD YEAR, a Systematic Syntax, and Lessons in Composition.** 2s. 6d.
- THE TEACHER'S COMPANION TO THE ABOVE.** With Copious Notes,  
Hints for Different Renderings, Synonyms, Philological Remarks, etc. By G.  
E. FASNACHT. Ex. fcap. 8vo. Each Year, 4s. 6d.
- \*MACMILLAN'S FRENCH COMPOSITION.**—By G. E. FASNACHT. Part I.  
Elementary. Ex. fcap. 8vo. 2s. 6d. Part II. Advanced. Cr. 8vo. 5s.
- THE TEACHER'S COMPANION TO THE ABOVE.** By G. E. FASNACHT.  
Ex. fcap. 8vo. Part I. 4s. 6d. Part II. 5s. net.
- A SPECIAL VOCABULARY TO MACMILLAN'S SECOND COURSE OF  
FRENCH COMPOSITION.** By the Same. *[In the Press.]*
- MACMILLAN'S PROGRESSIVE FRENCH READERS.** By G. E. FASNACHT. Ex.  
fcap. 8vo.
- \*FIRST YEAR, containing Tales, Historical Extracts, Letters, Dialogues, Ballads,  
Nursery Songs, etc., with Two Vocabulary: (1) in the order of subjects;  
(2) in alphabetical order. With Imitative Exercises.** 2s. 6d.
- \*SECOND YEAR, containing Fiction in Prose and Verse, Historical and Descriptive  
Extracts, Essays, Letters, Dialogues, etc. With Imitative Exercises.** 2s. 6d.
- MACMILLAN'S FOREIGN SCHOOL CLASSICS.** Ed. by G. E. FASNACHT. 18mo.
- \*CORNEILLE—LE CID.** By G. E. FASNACHT. 1s.
- \*DUMAS—LES DEMOISELLES DE ST. CYR.** By VICTOR OGER. 1s. 6d.
- LA FONTAINE'S FABLES.** By L. M. MORIARTY, B.A. *[In preparation.]*
- \*MOLIÈRE—L'AVARE.** By the same. 1s.
- \*MOLIÈRE—LE BOURGEOIS GENTILHOMME.** By the same. 1s. 6d.
- \*MOLIÈRE—LES FEMMES SAVANTES.** By G. E. FASNACHT. 1s.
- \*MOLIÈRE—LE MISANTHROPE.** By the same. 1s.
- \*MOLIÈRE—LE MÉDECIN MALGRÉ LUI.** By the same. 1s.
- \*MOLIÈRE—LES PRÉCIEUSES RIDICULES.** By the same. 1s.
- \*RACINE—BRITANNICUS.** By E. PELLISSIER, M.A. 2s.
- \*FRENCH READINGS FROM ROMAN HISTORY.** Selected from various  
Authors, by C. COLBECK, M.A., Assistant Master at Harrow. 4s. 6d.



- \*SAND, GEORGE—LA MARE AU DIABLE. By W. E. RUSSELL, M.A. Assistant Master at Haileybury. 1s.
- \*SANDEAU, JULES—MADEMOISELLE DE LA SEIGLIÈRE. By H. C. STEEL, Assistant Master at Winchester. 1s. 6d.
- \*VOLTAIRE—CHARLES XII. By G. E. FASNACHT. 3s. 6d.
- \*MASSON.—A COMPENDIOUS DICTIONARY OF THE FRENCH LANGUAGE. Adapted from the Dictionaries of Prof. A. ELWALL. By G. MASSON. Cr. 8vo. 3s. 6d.
- LA LYRE FRANÇAISE. Selected and arranged with Notes. 18mo. 2s. 6d. net.
- MOLIÈRE.—LE MALADE IMAGINAIRE. With Introduction and Notes, by F. TARVER, M.A., Assistant Master at Eton. Fcap. 8vo. 2s. 6d.
- PAYNE.—COMMERCIAL FRENCH. By J. B. PAYNE, King's College School, London. Gl. 8vo. *[In preparation.]*
- \*PELLISSIER.—FRENCH ROOTS AND THEIR FAMILIES. A Synthetic Vocabulary, based upon Derivations. By E. PELLISSIER, M.A., Assistant Master at Clifton College. Gl. 8vo. 6s.
- \*STORM.—FRENCH DIALOGUES. A Systematic Introduction to the Grammar and Idiom of spoken French. By JOH. STORM, LL.D. Intermediate Course. Translated by G. MACDONALD, M.A. Cr. 8vo. 4s. 6d.

## GERMAN.

- \*BEHAGHEL.—A SHORT HISTORICAL GRAMMAR OF THE GERMAN LANGUAGE. By Dr. OTTO BEHAGHEL. Translated by EMIL TRECHMANN, M.A., Ph.D., University of Sydney. Gl. 8vo. 3s. 6d.
- BUCHHEIM.—DEUTSCHE LYRIK. The Golden Treasury of the best German Lyrical Poems. Selected by Dr. BUCHHEIM. 18mo. 2s. 6d. net.
- BALLADEN UND ROMANZEN. Selection of the best German Ballads and Romances. By the same. 18mo. 2s. 6d. net.
- HUSS.—A SYSTEM OF ORAL INSTRUCTION IN GERMAN, by means of Progressive Illustrations and Applications of the leading Rules of Grammar. By H. C. O. HUSS, Ph.D. Cr. 8vo. 5s.
- MACMILLAN'S PRIMARY SERIES OF GERMAN READING BOOKS. Edited by G. E. FASNACHT. With Notes, Vocabulary, and Exercises. Gl. 8vo.
- \*GRIMM—KINDER UND HAUSMÄRCHEN. By G. E. FASNACHT. 2s. 6d.
- \*HAUFF—DIE KARAVANE. By HERMAN HAGER, Ph.D. 3s.
- \*HAUFF—DAS WIRTSCHAUS IM SPESSART. By G. E. FASNACHT. 3s.
- \*SCHMID, CHR. VON—H. VON EICHENFELS. By G. E. FASNACHT. 2s. 6d.
- MACMILLAN'S PROGRESSIVE GERMAN COURSE. By G. E. FASNACHT. Ex. fcap. 8vo.
- \*FIRST YEAR. Easy Lessons and Rules on the Regular Accidence. 1s. 6d.
- \*SECOND YEAR. Conversational Lessons in Systematic Accidence and Elementary Syntax. With Philological Illustrations and Vocabulary. 3s. 6d.
- [THIRD YEAR, in the Press.]*
- THE TEACHER'S COMPANION TO THE ABOVE. With copious Notes, Hints for Different Renderings, Synonyms, Philological Remarks, etc. By G. E. FASNACHT. Ex. fcap. 8vo. Each Year. 4s. 6d.
- MACMILLAN'S GERMAN COMPOSITION. By G. E. FASNACHT. Ex. fcap. 8vo.
- \*I. FIRST COURSE. Parallel German-English Extracts and Parallel English-German Syntax. 2s. 6d.
- THE TEACHER'S COMPANION TO THE ABOVE. By G. E. FASNACHT. FIRST COURSE. Gl. 8vo. 4s. 6d.
- MACMILLAN'S PROGRESSIVE GERMAN READERS. By G. E. FASNACHT. Ex. fcap. 8vo.
- \*FIRST YEAR, containing an Introduction to the German order of Words, with Copious Examples, extracts from German Authors in Prose and Poetry; Notes, and Vocabulary. 2s. 6d.
- MACMILLAN'S FOREIGN SCHOOL CLASSICS.—Edited by G. E. FASNACHT. 18mo.
- \*GOETHE—GÖTZ VON BERLICHINGEN. By H. A. BULL, M.A. 2s.
- \*GOETHE—FAUST. PART I, followed by an Appendix on PART II. By JANE LEE, Lecturer in German Literature at Newnham College, Cambridge. 4s. 6d.

- \*HEINE—SELECTIONS FROM THE REISEBILDER AND OTHER PROSE WORKS. By C. COLBECK, M.A., Assistant Master at Harrow. 2s. 6d.
- \*SCHILLER—SELECTIONS FROM SCHILLER'S LYRICAL POEMS. With a Memoir. By E. J. TURNER, B.A., and E. D. A. MORSHEAD, M.A. 2s. 6d.
- \*SCHILLER—DIE JUNGFRAU VON ORLEANS. By JOSEPH GOSTWICK. 2s. 6d.
- \*SCHILLER—MARIA STUART. By C. SHELDON, D.Litt. 2s. 6d.
- \*SCHILLER—WILHELM TELL. By G. E. FASNACHT. 2s. 6d.
- \*SCHILLER—WALLENSTEIN, DAS LAGER. By H. B. COTTERILL, M.A. 2s.
- \*UHLAND—SELECT BALLADS. Adapted for Beginners. With Vocabulary. By G. E. FASNACHT. 1s.
- \*PYLODET.—NEW GUIDE TO GERMAN CONVERSATION; containing an Alphabetical List of nearly 800 Familiar Words; followed by Exercises, Vocabulary, Familiar Phrases and Dialogues. By L. PYLODET. 18mo. 2s. 6d.
- SIEPMANN.—A GERMAN PRIMER. By OTTO SIEPMANN, Assistant Master in Clifton College. Globe 8vo. *[In preparation.]*
- \*SMITH.—COMMERCIAL GERMAN. By F. C. SMITH, M.A. Gl. 8vo. 3s. 6d.
- WHITNEY.—A COMPENDIOUS GERMAN GRAMMAR. By W. D. WHITNEY, Professor of Sanskrit and Instructor in Modern Languages in Yale College. Cr. 8vo. 4s. 6d.
- A GERMAN READER IN PROSE AND VERSE. By the same. With Notes and Vocabulary. Cr. 8vo. 5s.
- \*WHITNEY—EDGREN.—A COMPENDIOUS GERMAN AND ENGLISH DICTIONARY. By Prof. W. D. WHITNEY and A. H. EDGREN. Cr. 8vo. 5s.
- THE GERMAN-ENGLISH PART, separately, 3s. 6d.

## MODERN GREEK.

- CONSTANTINIDES.—NEO-HELLENICA. Dialogues illustrative of the development of the Greek Language. By Prof. M. CONSTANTINIDES. Cr. 8vo. 6s. net.
- VINCENT—DICKSON.—HANDBOOK TO MODERN GREEK. By Sir EDGAR VINCENT, K.C.M.G., and T. G. DICKSON, M.A. With Appendix on the relation of Modern and Classical Greek by Prof. JEBB. Cr. 8vo. 6s.

## ITALIAN.

- DANTE.—With Translation and Notes, by A. J. BUTLER, M.A.
- THE HELL. Cr. 8vo. 12s. 6d.
- THE PURGATORY. 2nd Ed. Cr. 8vo. 12s. 6d.
- THE PARADISE. 2nd Ed. Cr. 8vo. 12s. 6d.
- THE CONVITO. Cr. 8vo. *[In preparation.]*
- READINGS ON THE PURGATORIO OF DANTE. Chiefly based on the Commentary of Benvenuto Da Imola. By Hon. W. WARREN VERNON, M.A. With Introduction by DEAN CHURCH. 2 vols. Cr. 8vo. 24s.
- THE DIVINE COMEDY. Transl. by C. E. NORTON. I. HELL. II. PURGATORY. III. PARADISE. Cr. 8vo. 6s. each. THE NEW LIFE. Cr. 8vo. 5s.
- THE PURGATORY. Translated by C. L. SHADWELL, M.A. Ex. Cr. 8vo. 10s. net.
- A COMPANION TO DANTE. From the German of G. A. SCARTAZZINI. By A. J. BUTLER, M.A. Cr. 8vo. 10s. 6d.

## SPANISH.

- CALDERON.—FOUR PLAYS OF CALDERON. *El Principe Constante, La Vida es Sueño, El Alcalde de Zalamea, and El Escondido y La Tapada.* With Introduction and Notes. By NORMAN MACCOLL, M.A. Cr. 8vo. 14s.
- DELBOS.—COMMERCIAL SPANISH. By Prof. DELBOS. Gl. 8vo. *[In the Press.]*

## MATHEMATICS.

Arithmetic, Book-keeping, Algebra, Euclid and Pure Geometry, Geometrical Drawing, Mensuration, Trigonometry, Analytical Geometry (Plane and Solid), Problems and Questions in Mathematics, Higher Pure Mathematics, Mechanics (Statics, Dynamics, Hydrostatics, Hydrodynamics: see also Physics), Physics (Sound, Light, Heat, Electricity, Elasticity, Attractions, &c.), Astronomy, Historical.

## ARITHMETIC.

- \***ALDIS.**—THE GREAT GIANT ARITHMOS. A most Elementary Arithmetic for Children. By MARY STEADMAN ALDIS. Illustrated. Gl. 8vo. 2s. 6d.
- \***BRADSHAW.**—A COURSE OF EASY ARITHMETICAL EXAMPLES FOR BEGINNERS. By J. G. BRADSHAW, B.A., Assistant Master at Clifton College. Gl. 8vo. 2s. With Answers, 2s. 6d.
- \***BROOKSMITH.**—ARITHMETIC IN THEORY AND PRACTICE. By J. BROOKSMITH, M.A. Cr. 8vo. 4s. 6d. KEY, for Teachers only. Crown 8vo. 10s. 6d.
- \***BROOKSMITH.**—ARITHMETIC FOR BEGINNERS. By J. and E. J. BROOKSMITH. Gl. 8vo. 1s. 6d. KEY, for Teachers only. Cr. 8vo. 6s. 6d.
- CANDLER.**—HELP TO ARITHMETIC. For the use of Schools. By H. CANDLER, Mathematical Master of Uppingham School. 2nd Ed. Ex. fcap. 8vo. 2s. 6d.
- \***COLLAR.**—NOTES ON THE METRIC SYSTEM. By GEO. COLLAR, B.A., B.Sc. Gl. 8vo. 3d.
- \***DALTON.**—RULES AND EXAMPLES IN ARITHMETIC. By Rev. T. DALTON, M.A., Senior Mathematical Master at Eton. With Answers. 18mo. 2s. 6d.
- \***GOYEN.**—HIGHER ARITHMETIC AND ELEMENTARY MENSURATION. By P. GOYEN. Cr. 8vo. 5s. KEY. Cr. 8vo. 10s. net.
- \***HALL—KNIGHT.**—ARITHMETICAL EXERCISES AND EXAMINATION PAPERS. With an Appendix containing Questions in LOGARITHMS and MENSURATION. By H. S. HALL, M.A., Master of the Military Side, Clifton College, and S. R. KNIGHT, B.A., M.B., Ch.B. With or Without Answers. Gl. 8vo. 2s. 6d.
- HUNTER.**—DECIMAL APPROXIMATIONS. By H. St. J. HUNTER, M.A., Fellow of Jesus College, Cambridge. 18mo. 1s. 6d.
- \***JACKSON.**—COMMERCIAL ARITHMETIC. By S. JACKSON, M.A. Gl. 8vo. 3s. 6d.
- LOCK.**—Works by Rev. J. B. LOCK, M.A., Senior Fellow and Bursar of Gonville and Caius College, Cambridge.
- \***ARITHMETIC FOR SCHOOLS.** With Answers and 1000 additional Examples for Exercise. 4th Ed., revised. Gl. 8vo. 4s. 6d. Or, Part I. 2s. Part II. 3s. KEY, for Teachers only. Cr. 8vo. 10s. 6d.
- \***ARITHMETIC FOR BEGINNERS.** A School Class-Book of Commercial Arithmetic. Gl. 8vo. 2s. 6d. KEY, for Teachers only. Cr. 8vo. 8s. 6d.
- \***A SHILLING BOOK OF ARITHMETIC, FOR ELEMENTARY SCHOOLS.** 18mo. 1s. With Answers. 1s. 6d. [KEY in the Press.]
- LOCK—COLLAR.**—ARITHMETIC FOR THE STANDARDS. By Rev. J. B. LOCK, M.A., and GEO. COLLAR, B.A., B.Sc. Standards I. II. III. and IV., 2d. each; Standards V. VI. and VII., 3d. each. Answers to I. II. III. IV., 3d. each; to V. VI. and VII., 4d. each.
- MACMILLAN'S MENTAL ARITHMETIC.** For the Standards. Containing 6000 Questions and Answers. Standards I. II., 6d.; III. IV., 6d.; V. VI., 6d.
- \***PEDLEY.**—EXERCISES IN ARITHMETIC, containing 7000 Examples. By S. PEDLEY. Cr. 8vo. 5s. Also in Two Parts, 2s. 6d. each.
- SMITH.**—Works by Rev. BARNARD SMITH, M.A.
- \***ARITHMETIC FOR SCHOOLS.** Cr. 8vo. 4s. 6d. KEY, for Teachers. 8s. 6d.
- EXERCISES IN ARITHMETIC.** Cr. 8vo. 2s. With Answers, 2s. 6d. Answers separately, 6d.
- SCHOOL CLASS-BOOK OF ARITHMETIC.** 18mo. 3s. Or separately, in Three Parts, 1s. each. KEYS. Parts I. II. and III., 2s. 6d. each.
- SHILLING BOOK OF ARITHMETIC.** 18mo. Or separately, Part I., 2d.; Part II., 3d.; Part III., 7d. Answers, 6d. KEY, for Teachers only. 18mo. 4s. 6d.
- \***THE SAME,** with Answers. 18mo, cloth. 1s. 6d.
- EXAMINATION PAPERS IN ARITHMETIC.** 18mo. 1s. 6d. The Same, with Answers. 18mo. 2s. Answers, 6d. KEY. 18mo. 4s. 6d.
- THE METRIC SYSTEM OF ARITHMETIC, ITS PRINCIPLES AND APPLICATIONS,** with Numerous Examples. 18mo. 3d.
- A CHART OF THE METRIC SYSTEM,** on a Sheet, size 42 in. by 34 in. on Roller. New Ed. Revised by GEO. COLLAR, B.A., B.Sc. 4s. 6d.
- EASY LESSONS IN ARITHMETIC,** combining Exercises in Reading, Writing, Spelling, and Dictation. Part I. Cr. 8vo. 6d.
- EXAMINATION CARDS IN ARITHMETIC.** With Answers and Hints.



Standards I. and II., in box, 1s. Standards III. IV. and V., in boxes, 1s. each.  
Standard VI. in Two Parts, in boxes, 1s. each.

\***SMITH (BARNARD)—HUDSON.—ARITHMETIC FOR SCHOOLS.** By Rev. BARNARD SMITH, M.A., revised by W. H. H. HUDSON, M.A., Prof. of Mathematics, King's College, London. Cr. 8vo. 4s. 6d.

### BOOK-KEEPING.

\***THORNTON.—FIRST LESSONS IN BOOK-KEEPING.** By J. THORNTON. Cr. 8vo. 2s. 6d. KEY. Oblong 4to. 10s. 6d.

\***PRIMER OF BOOK-KEEPING.** 18mo. 1s. KEY. Demy 8vo. 2s. 6d.

\***EASY EXERCISES IN BOOK-KEEPING.** 18mo. 1s.

\***THE STUDENT'S MANUAL OF BOOK-KEEPING.** [*In the Press.*]

### ALGEBRA.

\***DALTON.—RULES AND EXAMPLES IN ALGEBRA.** By Rev. T. DALTON, Senior Mathematical Master at Eton. Part I. 18mo. 2s. KEY. Cr. 8vo. 7s. 6d. Part II. 18mo. 2s. 6d.

**DUPUIS.—PRINCIPLES OF ELEMENTARY ALGEBRA.** By N. F. DUPUIS, M.A., Professor of Mathematics, University of Queen's College, Kingston, Canada. Cr. 8vo. 6s.

**HALL—KNIGHT.—Works** by H. S. HALL, M.A., Master of the Military Side, Clifton College, and S. R. KNIGHT, B.A., M.B., Ch.B.

\***ALGEBRA FOR BEGINNERS.** Gl. 8vo. 2s. With Answers. 2s. 6d.

\***ELEMENTARY ALGEBRA FOR SCHOOLS.** 6th Ed. Gl. 8vo. 3s. 6d. With Answers, 4s. 6d. Answers, 1s. KEY, for Teachers only. 8s. 6d.

\***ALGEBRAICAL EXERCISES AND EXAMINATION PAPERS.** To accompany ELEMENTARY ALGEBRA. 2nd Ed., revised. With or Without Answers. Gl. 8vo. 2s. 6d.

\***HIGHER ALGEBRA.** 4th Ed. Cr. 8vo. 7s. 6d. KEY. Cr. 8vo. 10s. 6d.

\***JARMAN.—ALGEBRAIC FACTORS.** By J. ABBOT JARMAN. Gl. 8vo. 2s. With Answers, 2s. 6d.

\***JONES—CHEYNE.—ALGEBRAICAL EXERCISES.** Progressively Arranged. By Rev. C. A. JONES and C. H. CHEYNE, M.A., late Mathematical Masters at Westminster School. 18mo. 2s. 6d.

KEY, for Teachers. By Rev. W. FAILES, M.A. Cr. 8vo. 7s. 6d.

**SMITH.—Works** by CHARLES SMITH, M.A., Master of Sidney Sussex College, Cambridge.

\***ELEMENTARY ALGEBRA.** 2nd Ed., revised. Gl. 8vo. 4s. 6d. KEY, for Teachers only. Cr. 8vo. 10s. 6d.

\***A TREATISE ON ALGEBRA.** 4th Ed. Cr. 8vo. 7s. 6d. KEY. Cr. 8vo. 10s. 6d.

**TODHUNTER.—Works** by ISAAC TODHUNTER, F.R.S.

\***ALGEBRA FOR BEGINNERS.** 18mo. 2s. 6d. KEY. Cr. 8vo. 6s. 6d.

\***ALGEBRA FOR COLLEGES AND SCHOOLS.** By ISAAC TODHUNTER, F.R.S. Cr. 8vo. 7s. 6d. KEY, for Teachers. Cr. 8vo. 10s. 6d.

### EUCLID AND PURE GEOMETRY.

**COCKSHOTT—WALTERS.—A TREATISE ON GEOMETRICAL CONICS.** By A. COCKSHOTT, M.A., Assistant Master at Eton, and Rev. F. B. WALTERS, M.A., Principal of King William's College, Isle of Man. Cr. 8vo. 5s.

**CONSTABLE.—GEOMETRICAL EXERCISES FOR BEGINNERS.** By SAMUEL CONSTABLE. Cr. 8vo. 3s. 6d.

**CUTHBERTSON.—EUCLIDIAN GEOMETRY.** By FRANCIS CUTHBERTSON, M.A., LL.D. Ex. fcap. 8vo. 4s. 6d.

**DAY.—PROPERTIES OF CONIC SECTIONS PROVED GEOMETRICALLY.** By Rev. H. G. DAY, M.A. Part I. The Ellipse, with an ample collection of Problems. Cr. 8vo. 3s. 6d.

\***DEAKIN.—RIDER PAPERS ON EUCLID. BOOKS I. AND II.** By RUPERT DEAKIN, M.A. 18mo. 1s.

**DODGSON.—Works** by CHARLES L. DODGSON, M.A., Student and late Mathematical Lecturer, Christ Church, Oxford.

**EUCLID, BOOKS I. AND II.** 6th Ed., with words substituted for the Algebraical Symbols used in the 1st Ed. Cr. 8vo. 2s.

**EUCLID AND HIS MODERN RIVALS.** 2nd Ed. Cr. 8vo. 6s.

- CURIOSA MATHEMATICA.** Part I. A New Theory of Parallels. 3rd Ed. Cr. 8vo. 2s. Part II. Pillow Problems. 2nd Ed. Cr. 8vo. 2s.
- DREW.**—GEOMETRICAL TREATISE ON CONIC SECTIONS. By W. H. DREW, M.A. New Ed., enlarged. Cr. 8vo. 5s.
- DUPUIS.**—ELEMENTARY SYNTHETIC GEOMETRY OF THE POINT, LINE AND CIRCLE IN THE PLANE. By N. F. DUPUIS, M.A., Professor of Mathematics, University of Queen's College, Kingston, Canada. Gl. 8vo. 4s. 6d.
- SYNTHETIC SOLID GEOMETRY.** [*In the Press.*]
- \***HALL—STEVENS.**—A TEXT-BOOK OF EUCLID'S ELEMENTS. By H. S. HALL, M.A., and F. H. STEVENS, M.A., Masters of the Military Side, Clifton College. Gl. 8vo. Book I., 1s.; Books I. and II., 1s. 6d.; Books I.-IV., 3s.; Books III.-IV., 2s.; Books III.-VI., 3s.; Books V.-VI. and XI., 2s. 6d.; Books I.-VI. and XI., 4s. 6d.; Book XI., 1s. KEY to Books I.-IV., 6s. 6d. KEY to VI. and XI., 3s. 6d. KEY to I.-VI. and XI., 8s. 6d.
- HALSTED.**—THE ELEMENTS OF GEOMETRY. By G. B. HALSTED, Professor of Pure and Applied Mathematics in the University of Texas. 8vo. 12s. 6d.
- HAYWARD.**—THE ELEMENTS OF SOLID GEOMETRY. By R. B. HAYWARD, M.A., F.R.S. Gl. 8vo. 3s.
- LACHLAN.**—AN ELEMENTARY TREATISE ON MODERN PURE GEOMETRY. By R. LACHLAN, M.A. 8vo. 9s.
- \***LOCK.**—THE FIRST BOOK OF EUCLID'S ELEMENTS ARRANGED FOR BEGINNERS. By Rev. J. B. LOCK, M.A. Gl. 8vo. 1s. 6d.
- M'CLELLAND.**—A TREATISE ON THE GEOMETRY OF THE CIRCLE, and some extensions to Conic Sections by the Method of Reciprocation. By W. J. M'CLELLAND, M.A. Cr. 8vo. 6s.
- MILNE—DAVIS.**—GEOMETRICAL CONICS. By Rev. J. J. MILNE, M.A., and R. F. DAVIS, M.A. Cr. 8vo. Part I. The Parabola. 2s. Part II. The Central Conic. [*In the Press.*]
- MUKHOPADHYAY.**—GEOMETRICAL CONIC SECTIONS. By ASUTOSH MUKHOPADHYAY, M.A., F.R.S.E. 4s. 6d.
- RAMSEY—RICHARDSON.**—MODERN PLANE GEOMETRY. By A. S. RAMSEY, Fettes College, Edinburgh, and Rev. G. RICHARDSON. [*In the Press.*]
- \***RICHARDSON.**—THE PROGRESSIVE EUCLID. Books I. and II. With Notes, Exercises, and Deductions. Edited by A. T. RICHARDSON, M.A.
- SMITH.**—GEOMETRICAL CONIC SECTIONS. By CHARLES SMITH, M.A., Master of Sidney Sussex College, Cambridge. [*In the Press.*]
- SMITH.**—INTRODUCTORY MODERN GEOMETRY OF POINT, RAY, AND CIRCLE. By W. B. SMITH, A.M., Ph.D., Professor of Mathematics, Missouri University. Cr. 8vo. 5s.
- SYLLABUS OF PLANE GEOMETRY** (corresponding to Euclid, Books I.-VI.)—Prepared by the Association for the Improvement of Geometrical Teaching. Cr. 8vo. Sewed. 1s.
- SYLLABUS OF MODERN PLANE GEOMETRY.**—Prepared by the Association for the Improvement of Geometrical Teaching. Cr. 8vo. Sewed. 1s.
- \***TODHUNTER.**—THE ELEMENTS OF EUCLID. By I. TODHUNTER, F.R.S. 18mo. 3s. 6d. \*Books I. and II. 1s. KEY. Cr. 8vo. 6s. 6d.
- \***WEEKS.**—EXERCISES IN EUCLID, GRADUATED AND SYSTEMATIZED. By W. WEEKS, Lecturer in Geometry, Training College, Exeter. 18mo. 2s.
- WILSON.**—Works by Archdeacon WILSON, M.A., late Headmaster of Clifton College. ELEMENTARY GEOMETRY. BOOKS I.-V. (Corresponding to Euclid, Books I.-VI.) Following the Syllabus of the Geometrical Association. Ex. fcap. 8vo. 4s. 6d.
- SOLID GEOMETRY AND CONIC SECTIONS.** With Appendices on Transversals and Harmonic Division. Ex. fcap. 8vo. 3s. 6d.

## GEOMETRICAL DRAWING.

- EAGLES.**—CONSTRUCTIVE GEOMETRY OF PLANE CURVES. By T. H. EAGLES, M.A., Instructor, Roy. Indian Engineering Coll. Cr. 8vo. 12s.
- EDGAR—PRITCHARD.**—NOTE-BOOK ON PRACTICAL SOLID OR DESCRIPTIVE GEOMETRY. Containing Problems with help for Solutions. By J. H. EDGAR and G. S. PRITCHARD. 4th Ed. Gl. 8vo. 4s. 6d.
- \***KITCHENER.**—A GEOMETRICAL NOTE-BOOK. Containing Easy Problems in Geometrical Drawing. By F. E. KITCHENER, M.A. 4to. 2s.

- MILLAR.**—ELEMENTS OF DESCRIPTIVE GEOMETRY. By J. B. MILLAR, Lecturer on Engineering in the Owens College, Manchester. Cr. 8vo. 6s.  
**PLANT.**—PRACTICAL PLANE AND DESCRIPTIVE GEOMETRY. By E. C. PLANT. [In preparation.]  
**SPANTON.**—PRACTICAL PLANE GEOMETRY. By T. H. SPANTON, Drawing Instructor, *H.M.S. Britannia*. [In the Press.]

### MENSURATION.

- STEVENS.**—ELEMENTARY MENSURATION. With Exercises on the Mensuration of Plane and Solid Figures. By F. H. STEVENS, M.A. Gl. 8vo. [In the Press.]  
**TEBAY.**—ELEMENTARY MENSURATION FOR SCHOOLS. By S. TEBAY. Ex. fcap. 8vo. 3s. 6d.  
**\*TODHUNTER.**—MENSURATION FOR BEGINNERS. By ISAAC TODHUNTER, F.R.S. 18mo. 2s. 6d. KEY. By Rev. FR. L. MCCARTHY. Cr. 8vo. 7s. 6d.

### TRIGONOMETRY.

- BOTTOMLEY.**—FOUR-FIGURE MATHEMATICAL TABLES. Comprising Logarithmic and Trigonometrical Tables, and Tables of Squares, Square Roots, and Reciprocals. By J. T. BOTTOMLEY, M.A., Lecturer in Natural Philosophy in the University of Glasgow. 8vo. 2s. 6d.  
**\*HALL—KNIGHT.**—ELEMENTARY TRIGONOMETRY. By H. S. HALL, M.A., and S. R. KNIGHT, B.A. Gl. 8vo. 4s. 6d. [Dec. 1892]  
**HAYWARD.**—THE ALGEBRA OF CO-PLANAR VECTORS AND TRIGONOMETRY. By R. B. HAYWARD, M.A., F.R.S. Cr. 8vo. 8s. 6d.  
**JOHNSON.**—A TREATISE ON TRIGONOMETRY. By W. E. JOHNSON, M.A. late Mathematical Lecturer at King's College, Cambridge. Cr. 8vo. 8s. 6d.  
**JONES.**—LOGARITHMIC TABLES. By Prof. G. W. JONES, Cornell University. 8vo. 4s. 6d. net.

[Three-digit numbers to four places; four-digit numbers to six places; primes below 20,000 to ten places; mathematical and physical constants; addition-subtraction logarithms; trigonometric logarithms to four and six places; squares, cubes, roots, reciprocals, and prime factors; interpolation coefficients; error and probability tables.]

- KNIGHT.**—HIGHER TRIGONOMETRY. By S. R. KNIGHT, B.A. [In preparation.]  
**\*LEVETT—DAVISON.**—THE ELEMENTS OF PLANE TRIGONOMETRY. By RAWDON LEVETT, M.A., and C. DAVISON, M.A., Assistant Masters at King Edward's School, Birmingham. Gl. 8vo. 6s. 6d.; or, in 2 parts, 3s. 6d. each.

**LOCK.**—Works by Rev. J. B. Lock, M.A.

- \*THE TRIGONOMETRY OF ONE ANGLE.** Gl. 8vo. 2s. 6d.  
**\*TRIGONOMETRY FOR BEGINNERS**, as far as the Solution of Triangles. 3rd Ed. Gl. 8vo. 2s. 6d. KEY, for Teachers. Cr. 8vo. 6s. 6d.  
**\*ELEMENTARY TRIGONOMETRY.** 6th Ed. Gl. 8vo. 4s. 6d. KEY, for Teachers. Cr. 8vo. 8s. 6d.

**HIGHER TRIGONOMETRY.** 5th Ed. 4s. 6d. Both Parts complete in One Volume. 7s. 6d. [KEY in preparation.]

- M'CLELLAND—PRESTON.**—A TREATISE ON SPHERICAL TRIGONOMETRY. By W. J. M'CLELLAND, M.A., and T. PRESTON, M.A. Cr. 8vo. 8s. 6d., or: Part I. To the End of Solution of Triangles, 4s. 6d. Part II., 5s.

**MATTHEWS.**—MANUAL OF LOGARITHMS. By G. F. MATTHEWS, B.A. 8vo. 5s. net.

**PALMER.**—PRACTICAL LOGARITHMS AND TRIGONOMETRY. By J. H. PALMER, Headmaster, R.N., *H.M.S. Cambridge*, Devonport. Gl. 8vo. 4s. 6d.

**SNOWBALL.**—THE ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY. By J. C. SNOWBALL. 14th Ed. Cr. 8vo. 7s. 6d.

**TODHUNTER.**—Works by ISAAC TODHUNTER, F.R.S.

- \*TRIGONOMETRY FOR BEGINNERS.** 18mo. 2s. 6d. KEY. Cr. 8vo. 8s. 6d.  
**PLANE TRIGONOMETRY.** Cr. 8vo. 5s. KEY. Cr. 8vo. 10s. 6d.

**A TREATISE ON SPHERICAL TRIGONOMETRY.** Cr. 8vo. 4s. 6d.

**TODHUNTER—HOGG.**—PLANE TRIGONOMETRY. By ISAAC TODHUNTER. Revised by R. W. HOGG, M.A. Cr. 8vo. 5s. [KEY in preparation.]

**WOLSTENHOLME.**—EXAMPLES FOR PRACTICE IN THE USE OF SEVEN-FIGURE LOGARITHMS. By JOSEPH WOLSTENHOLME, D.Sc., late Professor of Mathematics, Royal Indian Engineering Coll., Cooper's Hill. 8vo. 5s.



## ANALYTICAL GEOMETRY (Plane and Solid).

- DYER.**—EXERCISES IN ANALYTICAL GEOMETRY. By J. M. DYER, M.A., Assistant Master at Eton. Illustrated. Cr. 8vo. 4s. 6d.
- FERRERS.**—AN ELEMENTARY TREATISE ON TRILINEAR CO-ORDINATES, the Method of Reciprocal Polars, and the Theory of Projectors. By the Rev. N. M. FERRERS, D.D., F.R.S., Master of Gonville and Caius College, Cambridge. 4th Ed., revised. Cr. 8vo. 6s. 6d.
- FROST.**—Works by PERCIVAL FROST, D.Sc., F.R.S., Fellow and Mathematical Lecturer at King's College, Cambridge.
- AN ELEMENTARY TREATISE ON CURVE TRACING. 8vo. 12s.
- SOLID GEOMETRY. 3rd Ed. Demy 8vo. 16s.
- HINTS FOR THE SOLUTION OF PROBLEMS in the above. 8vo. 8s. 6d.
- JOHNSON.**—CURVE TRACING IN CARTESIAN CO-ORDINATES. By W. WOOLSEY JOHNSON, Professor of Mathematics at the U.S. Naval Academy, Annapolis, Maryland. Cr. 8vo. 4s. 6d.
- PUCKLE.**—AN ELEMENTARY TREATISE ON CONIC SECTIONS AND ALGEBRAIC GEOMETRY. By G. H. PUCKLE, M.A. 5th Ed. Cr. 8vo. 7s. 6d.
- SMITH.**—Works by CHAS. SMITH, M.A., Master of Sidney Sussex Coll., Cambridge.
- CONIC SECTIONS. 7th Ed. Cr. 8vo. 7s. 6d. KEY. Cr. 8vo. 10s. 6d.
- AN ELEMENTARY TREATISE ON SOLID GEOMETRY. Cr. 8vo. 9s. 6d.
- TODHUNTER.**—Works by ISAAC TODHUNTER, F.R.S.
- PLANE CO-ORDINATE GEOMETRY, as applied to the Straight Line and the Conic Sections. Cr. 8vo. 7s. 6d. KEY. Cr. 8vo. 10s. 6d.
- EXAMPLES OF ANALYTICAL GEOMETRY OF THREE DIMENSIONS. New Ed., revised. Cr. 8vo. 4s.

## PROBLEMS &amp; QUESTIONS IN MATHEMATICS.

- ARMY PRELIMINARY EXAMINATION, PAPERS 1886-1893.** With Answers to the Mathematical Questions. 5th Ed. Cr. 8vo. 3s. 6d.
- BALL.**—MATHEMATICAL RECREATIONS AND PROBLEMS OF PAST AND PRESENT TIMES. By W. W. ROUSE BALL, M.A., Fellow and Lecturer of Trinity College, Cambridge. 2nd Ed. Cr. 8vo. 7s. net.
- CAMBRIDGE SENATE-HOUSE PROBLEMS AND RIDERS, WITH SOLUTIONS—**
- 1875—PROBLEMS AND RIDERS. By A. G. GREENHILL, F.R.S. Cr. 8vo. 8s. 6d.
- 1878—SOLUTIONS OF SENATE-HOUSE PROBLEMS. Edited by J. W. L. GLAISHER, F.R.S., Fellow of Trinity College, Cambridge. Cr. 8vo. 12s.
- CHRISTIE.**—A COLLECTION OF ELEMENTARY TEST-QUESTIONS IN PURE AND MIXED MATHEMATICS. By J. R. CHRISTIE, F.R.S. Cr. 8vo. 8s. 6d.
- CLIFFORD.**—MATHEMATICAL PAPERS. By W. K. CLIFFORD. 8vo. 30s.
- MACMILLAN'S MENTAL ARITHMETIC.** (See page 21.)
- MILNE.**—WEEKLY PROBLEM PAPERS. By Rev. JOHN J. MILNE, M.A. Pott 8vo. 4s. 6d.
- SOLUTIONS TO THE ABOVE. By the same. Cr. 8vo. 10s. 6d.
- COMPANION TO WEEKLY PROBLEM PAPERS. Cr. 8vo. 10s. 6d.
- \***RICHARDSON.**—PROGRESSIVE MATHEMATICAL EXERCISES FOR HOME WORK. By A. T. RICHARDSON, M.A. Gl. 8vo. First Series. 2s. With Answers, 2s. 6d. Second Series. 3s. With Answers, 3s. 6d.
- SANDHURST MATHEMATICAL PAPERS,** for Admission into the Royal Military College, 1881-1889. Edited by E. J. BROOKSMITH, B.A. Cr. 8vo. 3s. 6d.
- THOMAS.**—ENUNCIATIONS IN ARITHMETIC, ALGEBRA, EUCLID, AND TRIGONOMETRY, with Examples and Notes. By P. A. THOMAS, B.A. Gl. 8vo. 2s.
- WOOLWICH MATHEMATICAL PAPERS,** for Admission into the Royal Military Academy, Woolwich, 1880-1890 inclusive. By the same. Cr. 8vo. 6s.
- WOLSTENHOLME.**—MATHEMATICAL PROBLEMS, on Subjects included in the First and Second Divisions of Cambridge Mathematical Tripos. By JOSEPH WOLSTENHOLME, D.Sc. 3rd Ed., greatly enlarged. 8vo. 18s.
- EXAMPLES FOR PRACTICE IN THE USE OF SEVEN-FIGURE LOG-ARITHMS. By the same. 8vo. 5s.

## HIGHER PURE MATHEMATICS.

- AIRY.**—Works by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal.

- ELEMENTARY TREATISE ON PARTIAL DIFFERENTIAL EQUATIONS.  
With Diagrams. 2nd Ed. Cr. 8vo. 5s. 6d.
- ON THE ALGEBRAICAL AND NUMERICAL THEORY OF ERRORS OF  
OBSERVATIONS AND THE COMBINATION OF OBSERVATIONS.  
2nd Ed., revised. Cr. 8vo. 6s. 6d.
- BOOLE.—THE CALCULUS OF FINITE DIFFERENCES. By G. BOOLE. 3rd  
Ed., revised by J. F. MOULTON, Q.C. Cr. 8vo. 10s. 6d.
- EDWARDS.—THE DIFFERENTIAL CALCULUS. By JOSEPH EDWARDS, M.A.  
With Applications and numerous Examples. New Ed. 8vo. 14s.
- THE DIFFERENTIAL CALCULUS FOR SCHOOLS. By the Same. Gl.  
8vo. 4s. 6d.
- THE INTEGRAL CALCULUS. By the same. [In preparation.]
- THE INTEGRAL CALCULUS FOR SCHOOLS. By the same. [In preparation.]
- FORSYTH.—A TREATISE ON DIFFERENTIAL EQUATIONS. By ANDREW  
RUSSELL FORSYTH, F.R.S., Fellow and Assistant Tutor of Trinity College,  
Cambridge. 2nd Ed. 8vo. 14s.
- FROST.—AN ELEMENTARY TREATISE ON CURVE TRACING. By PERCIVAL  
FROST, M.A., D.Sc. 8vo. 12s.
- GRAHAM.—GEOMETRY OF POSITION. By R. H. GRAHAM. Cr. 8vo. 7s. 6d.
- GREENHILL.—DIFFERENTIAL AND INTEGRAL CALCULUS. By A. G.  
GREENHILL, Professor of Mathematics to the Senior Class of Artillery Officers,  
Woolwich. New Ed. Cr. 8vo. 10s. 6d.
- APPLICATIONS OF ELLIPTIC FUNCTIONS. By the same. 8vo. 12s.
- HARKNESS—MORLEY. A TREATISE ON THE THEORY OF FUNCTIONS.  
By J. HARKNESS, M.A., and F. MORLEY, M.A. 8vo. 18s. net.
- HEMMING.—AN ELEMENTARY TREATISE ON THE DIFFERENTIAL AND  
INTEGRAL CALCULUS. By G. W. HEMMING, M.A. 2nd Ed. 8vo. 9s.
- JOHNSON.—Works by W. W. JOHNSON, Professor of Mathematics at the U.S.  
Naval Academy.  
INTEGRAL CALCULUS, an Elementary Treatise. Founded on the Method  
of Rates or Fluxions. 8vo. 9s.
- CURVE TRACING IN CARTESIAN CO-ORDINATES. Cr. 8vo. 4s. 6d.
- A TREATISE ON ORDINARY AND DIFFERENTIAL EQUATIONS. Ex. cr.  
8vo. 15s.
- KELLAND—TAIT.—INTRODUCTION TO QUATERNIONS, with numerous  
examples. By P. KELLAND and P. G. TAIT, Professors in the Department of  
Mathematics in the University of Edinburgh. 2nd Ed. Cr. 8vo. 7s. 6d.
- KEMPE.—HOW TO DRAW A STRAIGHT LINE: a Lecture on Linkages. By A.  
B. KEMPE. Illustrated. Cr. 8vo. 1s. 6d.
- KNOX.—DIFFERENTIAL CALCULUS FOR BEGINNERS. By ALEXANDER  
Knox, M.A. Fcap. 8vo. 3s. 6d.
- RICE—JOHNSON.—AN ELEMENTARY TREATISE ON THE DIFFEREN-  
TIAL CALCULUS. Founded on the Method of Rates or Fluxions. By J. M.  
RICE and W. W. JOHNSON 3rd Ed. 8vo. 18s. Abridged Ed. 9s.
- TODHUNTER.—Works by ISAAC TODHUNTER, F.R.S.  
AN ELEMENTARY TREATISE ON THE THEORY OF EQUATIONS.  
Cr. 8vo. 7s. 6d.
- A TREATISE ON THE DIFFERENTIAL CALCULUS. Cr. 8vo. 10s. 6d.
- KEY. Cr. 8vo. 10s. 6d.
- A TREATISE ON THE INTEGRAL CALCULUS AND ITS APPLICATIONS.  
Cr. 8vo. 10s. 6d. KEY. Cr. 8vo. 10s. 6d.
- A HISTORY OF THE MATHEMATICAL THEORY OF PROBABILITY, from  
the time of Pascal to that of Laplace. 8vo. 18s.
- WELD.—SHORT COURSE IN THE THEORY OF DETERMINANTS. By  
L. G. WELD, M.A. Cr. 8vo. 7s. 6d.

## MECHANICS: Statics, Dynamics, Hydrostatics, Hydrodynamics. (See also Physics.)

- ALEXANDER—THOMSON.—ELEMENTARY APPLIED MECHANICS. By  
Prof. T. ALEXANDER and A. W. THOMSON. Part II. Transverse Stress.  
Cr. 8vo. 10s. 6d.
- BALL.—EXPERIMENTAL MECHANICS. A Course of Lectures delivered at the

- Royal College of Science, Dublin. By Sir R. S. BALL, F.R.S. 2nd Ed. Illustrated. Cr. 8vo. 6s.
- CLIFFORD.**—THE ELEMENTS OF DYNAMIC. An Introduction to the Study of Motion and Rest in Solid and Fluid Bodies. By W. K. CLIFFORD. Part I.—Kinematic. Cr. 8vo. Books I.—III. 7s. 6d.; Book IV. and Appendix, 6s.
- COTTERILL.**—APPLIED MECHANICS: An Elementary General Introduction to the Theory of Structures and Machines. By J. H. COTTERILL, F.R.S., Professor of Applied Mechanics in the Royal Naval College, Greenwich. 3rd Ed. Revised. 8vo. 18s.
- COTTERILL—SLADE.**—LESSONS IN APPLIED MECHANICS. By Prof. J. H. COTTERILL and J. H. SLADE. Fcap. 8vo. 5s. 6d.
- GANGUILLET—KUTTER.**—A GENERAL FORMULA FOR THE UNIFORM FLOW OF WATER IN RIVERS AND OTHER CHANNELS. By E. GANGUILLET and W. R. KUTTER. Translated by R. HERING and J. C. TRAUTWINE. 8vo. 17s.
- GRAHAM.**—GEOMETRY OF POSITION. By R. H. GRAHAM. Cr. 8vo. 7s. 6d.
- \***GREAVES.**—STATICS FOR BEGINNERS. By JOHN GREAVES, M.A., Fellow and Mathematical Lecturer at Christ's College, Cambridge. Gl. 8vo. 3s. 6d.
- A TREATISE ON ELEMENTARY STATICS. By the same. Cr. 8vo. 5s.
- GREENHILL.**—ELEMENTARY HYDROSTATICS. By A. G. GREENHILL, Professor of Mathematics to the Senior Class of Artillery Officers, Woolwich. Gl. 8vo. [Immediately.]
- \***HICKS.**—ELEMENTARY DYNAMICS OF PARTICLES AND SOLIDS. By W. M. HICKS, D.Sc., Principal and Professor of Mathematics and Physics, Firth College, Sheffield. Cr. 8vo. 6s. 6d.
- HOSKINS.**—ELEMENTS OF GRAPHIC STATICS. By L. M. HOSKINS. 8vo. 10s. net.
- KENNEDY.**—THE MECHANICS OF MACHINERY. By A. B. W. KENNEDY, F.R.S. Illustrated. Cr. 8vo. 8s. 6d.
- LANGMAID—GAISFORD.**—(See Engineering, p. 39.)
- LOCK.**—Works by Rev. J. B. LOCK, M.A.
- \***MECHANICS FOR BEGINNERS.** Gl. 8vo. Part I. MECHANICS OF SOLIDS. 2s. 6d. [Part II. MECHANICS OF FLUIDS, *in preparation.*]
- \***ELEMENTARY STATICS.** 2nd Ed. Gl. 8vo. 3s. 6d. KEY. Cr. 8vo. 8s. 6d.
- \***ELEMENTARY DYNAMICS.** 3rd Ed. Gl. 8vo. 3s. 6d. KEY. Cr. 8vo. 8s. 6d.
- \***ELEMENTARY DYNAMICS AND STATICS.** Gl. 8vo. 6s. 6d.
- ELEMENTARY HYDROSTATICS.** Gl. 8vo. [In preparation.]
- MACGREGOR.**—KINEMATICS AND DYNAMICS. An Elementary Treatise. By J. G. MACGREGOR, D.Sc., Munro Professor of Physics in Dalhousie College, Halifax, Nova Scotia. Illustrated. Cr. 8vo. 10s. 6d.
- PARKINSON.**—AN ELEMENTARY TREATISE ON MECHANICS. By S. PARKINSON, D.D., F.R.S., late Tutor and Prælector of St. John's College, Cambridge. 6th Ed., revised. Cr. 8vo. 9s. 6d.
- PIRIE.**—LESSONS ON RIGID DYNAMICS. By Rev. G. PIRIE, M.A., Professor of Mathematics in the University of Aberdeen. Cr. 8vo. 6s.
- ROUTH.**—Works by EDWARD JOHN ROUTH, D.Sc., LL.D., F.R.S., Hon. Fellow of St. Peter's College, Cambridge.
- A TREATISE ON THE DYNAMICS OF THE SYSTEM OF RIGID BODIES. With numerous Examples. Two vols. 8vo. 5th Ed. Vol. I.—Elementary Parts. 14s. Vol. II.—The Advanced Parts. 14s.
- STABILITY OF A GIVEN STATE OF MOTION, PARTICULARLY STEADY MOTION. Adams Prize Essay for 1877. 8vo. 8s. 6d.
- \***SANDERSON.**—HYDROSTATICS FOR BEGINNERS. By F. W. SANDERSON, M.A., Assistant Master at Dulwich College. Gl. 8vo. 4s. 6d.
- SYLLABUS OF ELEMENTARY DYNAMICS.** Part I. Linear Dynamics. With an Appendix on the Meanings of the Symbols in Physical Equations. Prepared by the Association for the Improvement of Geometrical Teaching. 4to. 1s.
- TAIT—STEELE.**—A TREATISE ON DYNAMICS OF A PARTICLE. By Professor TAIT, M.A., and W. J. STEELE, B.A. 6th Ed., revised. Cr. 8vo. 12s.
- TODHUNTER.**—Works by ISAAC TODHUNTER, F.R.S.
- \***MECHANICS FOR BEGINNERS.** 18mo. 4s. 6d. KEY. Cr. 8vo. 6s. 6d.
- A TREATISE ON ANALYTICAL STATICS. 5th Ed. Edited by Prof. J. D. EVERETT, F.R.S. Cr. 8vo. 10s. 6d.



**WEISBACH—HERMANN.**—MECHANICS OF HOISTING MACHINERY. By Dr. J. WEISBACH and Prof. G. HERMANN. Translated by K. P. DAHLSTROM, M.E. *[In the Press.]*

**ZIWET.**—AN ELEMENTARY TREATISE ON THEORETICAL MECHANICS. By Prof. A. ZIWET. 8vo. Part I. KINEMATICS. 8s. 6d. net. Part II. STATICS. *[In the Press.]*

**PHYSICS:** Sound, Light, Heat, Electricity, Elasticity, Attractions, etc. (See also Mechanics.)

**AIRY.**—ON SOUND AND ATMOSPHERIC VIBRATIONS. By Sir G. B. AIRY, K.C.B. With the Mathematical Elements of Music. Cr. 8vo. 9s.

**BARKER.**—PHYSICS. Advanced Course. By Prof. G. F. BARKER. 8vo. 21s.

**CUMMING.**—AN INTRODUCTION TO THE THEORY OF ELECTRICITY. By LINNÆUS CUMMING, M.A., Assistant Master at Rugby. Illustrated. Cr. 8vo. 8s. 6d.

**DANIELL.**—A TEXT-BOOK OF THE PRINCIPLES OF PHYSICS. By ALFRED DANIELL, D.Sc. Illustrated. 2nd Ed., revised and enlarged. 8vo. 21s.

**DAY.**—ELECTRIC LIGHT ARITHMETIC. By R. E. DAY. Pott 8vo. 2s.

**EVERETT.**—ILLUSTRATIONS OF THE C. G. S. SYSTEM OF UNITS WITH TABLES OF PHYSICAL CONSTANTS. By J. D. EVERETT, F.R.S., Professor of Natural Philosophy, Queen's College, Belfast. New Ed. Ex. fcap. 8vo. 5s.

**FESSENDEN.**—PHYSICS FOR PUBLIC SCHOOLS. By C. FESSENDEN, Principal of the Collegiate Institute, Peterboro, Ontario. Illustrated. Fcap. 8vo. 3s.

**GRAY.**—THE THEORY AND PRACTICE OF ABSOLUTE MEASUREMENTS IN ELECTRICITY AND MAGNETISM. By A. GRAY, F.R.S.E., Professor of Physics, University College, Bangor. Two vols. Cr. 8vo. Vol. I. 12s. 6d. Vol. II. In 2 Parts. 25s.

**ABSOLUTE MEASUREMENTS IN ELECTRICITY AND MAGNETISM.** 2nd Ed., revised and greatly enlarged. Fcap. 8vo. 5s. 6d.

**ELECTRIC LIGHTING AND POWER DISTRIBUTION.** *[In preparation.]*

**HANDBOOK OF ELECTRIC LIGHT ENGINEERING.** *[In preparation.]*

**HEAVISIDE.**—ELECTRICAL PAPERS. By O. HEAVISIDE. 2 vols. 8vo. 30s. net.

**HERTZ.**—RESEARCHES ON THE PROPAGATION OF ELECTRICAL FORCE. By Prof. H. HERTZ of Bonn. Translated by D. E. JONES, B.Sc. With Preface by Lord KELVIN, P.R.S. 8vo. *[In the Press.]*

**IBBETSON.**—THE MATHEMATICAL THEORY OF PERFECTLY ELASTIC SOLIDS, with a Short Account of Viscous Fluids. By W. J. IBBETSON, late Senior Scholar of Clare College, Cambridge. 8vo. 21s.

**JACKSON.**—TEXT-BOOK ON ELECTRO-MAGNETISM AND THE CONSTRUCTION OF DYNAMOS. By Prof. D. C. JACKSON. *[In the Press.]*

**NOTES ON ELECTRO-MAGNETS AND THE CONSTRUCTION OF DYNAMOS.** *[In the Press.]*

**JOHNSON.**—NATURE'S STORY BOOKS. SUNSHINE. By AMY JOHNSON, LL.A. Illustrated. Cr. 8vo. 6s.

\***JONES.**—EXAMPLES IN PHYSICS. With Answers and Solutions. By D. E. JONES, B.Sc., late Professor of Physics, University College of Wales, Aberystwith. 2nd Ed., revised and enlarged. Fcap. 8vo. 3s. 6d.

\***ELEMENTARY LESSONS IN HEAT, LIGHT, AND SOUND.** By the same. Gl. 8vo. 2s. 6d.

**LESSONS IN HEAT AND LIGHT.** For Matriculation Students. By the same. Globe 8vo. 3s. 6d.

**KELVIN.**—Works by Lord KELVIN, P.R.S., Professor of Natural Philosophy in the University of Glasgow.

**ELECTROSTATICS AND MAGNETISM, REPRINTS OF PAPERS ON.** 2nd Ed. 8vo. 18s.

**POPULAR LECTURES AND ADDRESSES.** 3 vols. Illustrated. Cr. 8vo. Vol. I. CONSTITUTION OF MATTER. 7s. 6d. Vol. III. NAVIGATION. 7s. 6d.

**LODGE.**—MODERN VIEWS OF ELECTRICITY. By OLIVER J. LODGE, F.R.S., Professor of Physics, University College, Liverpool. Illus. Cr. 8vo. 6s. 6d.

**LOEWY.**—\*QUESTIONS AND EXAMPLES ON EXPERIMENTAL PHYSICS: Sound, Light, Heat, Electricity, and Magnetism. By B. LOEWY, Examiner in Experimental Physics to the College of Preceptors. Fcap. 8vo. 2s.

- \*A GRADUATED COURSE OF NATURAL SCIENCE FOR ELEMENTARY AND TECHNICAL SCHOOLS AND COLLEGES. By the same. Part I. FIRST YEAR'S COURSE. Gl. 8vo. 2s. Part II. 2s. 6d.
- LUPTON.—NUMERICAL TABLES AND CONSTANTS IN ELEMENTARY SCIENCE. By S. LUPTON, M.A. Ex. fcap. 8vo. 2s. 6d.
- McAULAY.—UTILITY OF QUATERNIONS IN PHYSICS. By ALEX. McAULAY. 8vo. 5s. net.
- MACFARLANE.—PHYSICAL ARITHMETIC. By A. MACFARLANE, D.Sc., late Examiner in Mathematics at the University of Edinburgh. Cr. 8vo. 7s. 6d.
- \*MAYER.—SOUND: A Series of Simple Experiments. By A. M. MAYER, Prof. of Physics in the Stevens Institute of Technology. Illustrated. Cr. 8vo. 3s. 6d.
- \*MAYER—BARNARD.—LIGHT: A Series of Simple Experiments. By A. M. MAYER and C. BARNARD. Illustrated. Cr. 8vo. 2s. 6d.
- MOLLOY.—GLEANINGS IN SCIENCE: Popular Lectures. By Rev. GERALD MOLLOY, D.Sc., Rector of the Catholic University of Ireland. 8vo. 7s. 6d.
- NEWTON.—PRINCIPIA. Edited by Prof. Sir W. THOMSON, P.R.S., and Prof. BLACKBURNE. 4to. 31s. 6d.
- THE FIRST THREE SECTIONS OF NEWTON'S PRINCIPIA. With Notes, Illustrations, and Problems. By P. FROST, M.A., D.Sc. 3rd Ed. 8vo. 12s.
- PARKINSON.—A TREATISE ON OPTICS. By S. PARKINSON, D.D., F.R.S., late Tutor of St. John's College, Cambridge. 4th Ed. Cr. 8vo. 10s. 6d.
- PEABODY.—THERMODYNAMICS OF THE STEAM-ENGINE AND OTHER HEAT-ENGINES. By CECIL H. PEABODY. 8vo. 21s.
- PERRY.—STEAM: An Elementary Treatise. By JOHN PERRY, Prof. of Applied Mechanics, Technical College, Finsbury. 18mo. 4s. 6d.
- PICKERING.—ELEMENTS OF PHYSICAL MANIPULATION. By Prof. EDWARD C. PICKERING. Medium 8vo. Part I., 12s. 6d. Part II., 14s.
- PRESTON.—THE THEORY OF LIGHT. By THOMAS PRESTON, M.A. Illustrated. 8vo. 15s. net.
- THE THEORY OF HEAT. By the same. 8vo. [In the Press.]
- RAYLEIGH.—THE THEORY OF SOUND. By LORD RAYLEIGH, F.R.S. 8vo. New Edition. [In the Press.]
- SANDERSON.—ELECTRICITY AND MAGNETISM FOR BEGINNERS. By F. W. SANDERSON. [In the Press.]
- SHANN.—AN ELEMENTARY TREATISE ON HEAT, IN RELATION TO STEAM AND THE STEAM-ENGINE. By G. SHANN, M.A. Cr. 8vo. 4s. 6d.
- SPOTTISWOODE.—POLARISATION OF LIGHT. By the late W. SPOTTISWOODE, F.R.S. Illustrated. Cr. 8vo. 3s. 6d.
- STEWART.—Works by BALFOUR STEWART, F.R.S., late Langworthy Professor of Physics, Owens College, Manchester.
- \*A PRIMER OF PHYSICS. Illustrated. With Questions. 18mo. 1s.
- \*LESSONS IN ELEMENTARY PHYSICS. Illustrated. Fcap. 8vo. 4s. 6d.
- \*QUESTIONS ON THE ABOVE. By Prof. T. H. CORE. Fcap. 8vo. 2s.
- STEWART—GEE.—LESSONS IN ELEMENTARY PRACTICAL PHYSICS. By BALFOUR STEWART, F.R.S., and W. W. HALDANE GEE, B.Sc. Cr. 8vo. Vol. I. GENERAL PHYSICAL PROCESSES. 6s. Vol. II. ELECTRICITY AND MAGNETISM. 7s. 6d. [Vol. III. OPTICS, HEAT, AND SOUND. In the Press.]
- \*PRACTICAL PHYSICS FOR SCHOOLS AND THE JUNIOR STUDENTS OF COLLEGES. Gl. 8vo. Vol. I. ELECTRICITY AND MAGNETISM. 2s. 6d. [Vol. II. OPTICS, HEAT, AND SOUND. In the Press.]
- STOKES.—ON LIGHT. Burnett Lectures. By Sir G. G. STOKES, F.R.S., Lucasian Professor of Mathematics in the University of Cambridge. I. ON THE NATURE OF LIGHT. II. ON LIGHT AS A MEANS OF INVESTIGATION. III. ON THE BENEFICIAL EFFECTS OF LIGHT. 2nd Ed. Cr. 8vo. 7s. 6d.
- STONE.—AN ELEMENTARY TREATISE ON SOUND. By W. H. STONE. Illustrated. Fcap. 8vo. 3s. 6d.
- TAIT.—HEAT. By P. G. TAIT, Professor of Natural Philosophy in the University of Edinburgh. Cr. 8vo. 6s.
- LECTURES ON SOME RECENT ADVANCES IN PHYSICAL SCIENCE. By the same. 3rd Edition. Crown 8vo. 9s.
- TAYLOR.—SOUND AND MUSIC. An Elementary Treatise on the Physical Constitution of Musical Sounds and Harmony, including the Chief Acoustical Discoveries of Prof. Helmholtz. By S. TAYLOR, M.A. Ex. cr. 8vo. 8s. 6d.

- \*THOMPSON.**—ELEMENTARY LESSONS IN ELECTRICITY AND MAGNETISM. By SILVANUS P. THOMPSON, Principal and Professor of Physics in the Technical College, Finsbury. Illustrated. Fcap. 8vo. 4s. 6d.
- THOMSON.**—Works by J. J. THOMSON, Professor of Experimental Physics in the University of Cambridge.
- A TREATISE ON THE MOTION OF VORTEX RINGS. 8vo. 6s.
- APPLICATIONS OF DYNAMICS TO PHYSICS AND CHEMISTRY. Cr. 8vo. 7s. 6d.
- TURNER.**—A COLLECTION OF EXAMPLES ON HEAT AND ELECTRICITY. By H. H. TURNER, Fellow of Trinity College, Cambridge. Cr. 8vo. 2s. 6d.
- WRIGHT.**—LIGHT: A Course of Experimental Optics, chiefly with the Lantern. By LEWIS WRIGHT. Illustrated. New Ed. Cr. 8vo. 7s. 6d.

## ASTRONOMY.

- AIRY.**—Works by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal.
- \*POPULAR ASTRONOMY. Revised by H. H. TURNER, M.A. 18mo. 4s. 6d.
- GRAVITATION: An Elementary Explanation of the Principal Perturbations in the Solar System. 2nd Ed. Cr. 8vo. 7s. 6d.
- CHEYNE.**—AN ELEMENTARY TREATISE ON THE PLANETARY THEORY. By C. H. H. CHEYNE. With Problems. 3rd Ed., revised. Cr. 8vo. 7s. 6d.
- CLARK—SADLER.**—THE STAR GUIDE. By L. CLARK and H. SADLER. Roy. 8vo. 5s.
- CROSSLEY—GLEDHILL—WILSON.**—A HANDBOOK OF DOUBLE STARS. By E. CROSSLEY, J. GLEDHILL, and J. M. WILSON. 8vo. 21s.
- CORRECTIONS TO THE HANDBOOK OF DOUBLE STARS. 8vo. 1s.
- FORBES.**—TRANSIT OF VENUS. By G. FORBES, Professor of Natural Philosophy in the Andersonian University, Glasgow. Illustrated. Cr. 8vo. 3s. 6d.
- GODFRAY.**—Works by HUGH GODFRAY, M.A., Mathematical Lecturer at Pembroke College, Cambridge.
- A TREATISE ON ASTRONOMY. 4th Ed. 8vo. 12s. 6d.
- AN ELEMENTARY TREATISE ON THE LUNAR THEORY. Cr. 8vo. 5s. 6d.
- LOCKYER.**—Works by J. NORMAN LOCKYER, F.R.S.
- \*A PRIMER OF ASTRONOMY. Illustrated. 18mo. 1s.
- \*ELEMENTARY LESSONS IN ASTRONOMY. With Spectra of the Sun, Stars, and Nebulae, and Illus. 36th Thousand. Revised throughout. Fcap. 8vo. 5s. 6d.
- \*QUESTIONS ON THE ABOVE. By J. FORBES ROBERTSON. 18mo. 1s. 6d.
- THE CHEMISTRY OF THE SUN. Illustrated. 8vo. 14s.
- THE METEORITIC HYPOTHESIS OF THE ORIGIN OF COSMICAL SYSTEMS. Illustrated. 8vo. 17s. net.
- STAR-GAZING PAST AND PRESENT. Expanded from Notes with the assistance of G. M. SEABROKE, F.R.A.S. Roy. 8vo. 21s.
- LODGE.**—PIONEERS OF SCIENCE. By OLIVER J. LODGE. Ex. Cr. 8vo. 7s. 6d.
- NEWCOMB.**—POPULAR ASTRONOMY. By S. NEWCOMB, LL.D., Professor U.S. Naval Observatory. Illustrated. 2nd Ed., revised. 8vo. 18s.

## HISTORICAL.

- BALL.**—A SHORT ACCOUNT OF THE HISTORY OF MATHEMATICS. By W. W. ROUSE BALL, M.A. 2nd ed. Cr. 8vo. 10s. net.
- MATHEMATICAL RECREATIONS, AND PROBLEMS OF PAST AND PRESENT TIMES. By the same. Cr. 8vo. 7s. net.
- AN ESSAY ON NEWTON'S PRINCIPIA. By the same. Cr. 8vo. 6s. net.
- CAJORI.**—HISTORY OF MATHEMATICS. By Prof. F. CAJORI. [*In the Press.*]

## NATURAL SCIENCES.

Chemistry; Physical Geography, Geology, and Mineralogy; Biology (*Botany, Zoology, General Biology, Physiology*); Medicine.

### CHEMISTRY.

- ARMSTRONG.**—A MANUAL OF INORGANIC CHEMISTRY. By H. E. ARMSTRONG, F.R.S., Professor of Chemistry, City and Guilds Central Institute. [*In preparation.*]



- BEHRENS.**—MICRO-CHEMICAL METHODS OF ANALYSIS. By Prof. BEHRENS. With Preface by Prof. J. W. JUDD, F.R.S. Cr. 8vo. [*In the Press.*]
- \***COHEN.**—THE OWENS COLLEGE COURSE OF PRACTICAL ORGANIC CHEMISTRY. By JULIUS B. COHEN, Ph.D., Assistant Lecturer on Chemistry, Owens College, Manchester. Fcap. 8vo. 2s. 6d.
- \***DOBBIN-WALKER.**—CHEMICAL THEORY FOR BEGINNERS. By L. DOBBIN, Ph.D., and JAS. WALKER, Ph.D., Assistants in the Chemistry Department, University of Edinburgh. 18mo. 2s. 6d.
- FLEISCHER.**—A SYSTEM OF VOLUMETRIC ANALYSIS. By EMIL FLEISCHER. Translated, with Additions, by M. M. P. MUIR, F.R.S.E. Cr. 8vo. 7s. 6d.
- FRANKLAND.**—AGRICULTURAL CHEMICAL ANALYSIS. (*See Agriculture.*)
- GORDON.**—ELEMENTARY COURSE OF PRACTICAL SCIENCE. By HUGH GORDON, Science Instructor under the London School Board. Pott 8vo. 1s.
- HARTLEY.**—A COURSE OF QUANTITATIVE ANALYSIS FOR STUDENTS. By W. N. HARTLEY, F.R.S., Professor of Chemistry, Royal College of Science, Dublin. Gl. 8vo. 5s.
- HEMPEL.**—METHODS OF GAS ANALYSIS. By Dr. WALTHER HEMPEL. Translated by Dr. L. M. DENNIS. Cr. 8vo. 7s. 6d.
- HIORNS.**—Works by A. H. HIORNS, Principal of the School of Metallurgy, Birmingham and Midland Institute. Gl. 8vo.
- A TEXT-BOOK OF ELEMENTARY METALLURGY. 4s.
- PRACTICAL METALLURGY AND ASSAYING. 6s.
- IRON AND STEEL MANUFACTURE. For Beginners. 3s. 6d.
- MIXED METALS OR METALLIC ALLOYS. 6s.
- METAL COLOURING AND BRONZING. 5s.
- JONES.**—\*THE OWENS COLLEGE JUNIOR COURSE OF PRACTICAL CHEMISTRY. By FRANCIS JONES, F.R.S.E., Chemical Master at the Grammar School, Manchester. Illustrated. Fcap. 8vo. 2s. 6d.
- \***QUESTIONS ON CHEMISTRY.** Inorganic and Organic. By the same. Fcap. 8vo. 3s.
- LANDAUER.**—BLOWPIPE ANALYSIS. By J. LANDAUER. Translated by J. TAYLOR, B.Sc. Revised Edition. Gl. 8vo. 4s. 6d.
- LASSAR-COHN.**—LABORATORY MANUAL OF ORGANIC CHEMISTRY. Translated by Prof. ALEX. SMITH. Cr. 8vo. [*In preparation.*]
- LAURIE.**—(*See Agriculture*, p. 40.)
- LOCKYER.**—THE CHEMISTRY OF THE SUN. By J. NORMAN LOCKYER, F.R.S. Illustrated. 8vo. 14s.
- LUPTON.**—CHEMICAL ARITHMETIC. With 1200 Problems. By S. LUPTON, M.A. 2nd Ed., revised. Fcap. 8vo. 4s. 6d.
- MELDOLA.**—THE CHEMISTRY OF PHOTOGRAPHY. By RAPHAEL MELDOLA, F.R.S., Professor of Chemistry, Technical College, Finsbury. Cr. 8vo. 6s.
- MEYER.**—HISTORY OF CHEMISTRY FROM THE EARLIEST TIMES TO THE PRESENT DAY. By ERNST VON MEYER, Ph.D. Translated by GEORGE MCGOWAN, Ph.D. 8vo. 14s. net.
- MIXTER.**—AN ELEMENTARY TEXT-BOOK OF CHEMISTRY. By W.G. MIXTER, Professor of Chemistry, Yale College. 2nd Ed. Cr. 8vo. 7s. 6d.
- MUIR.**—PRACTICAL CHEMISTRY FOR MEDICAL STUDENTS: First M.B. Course. By M. M. P. MUIR, F.R.S.E., Fellow and Prælector in Chemistry at Gonville and Caius College, Cambridge. Fcap. 8vo. 1s. 6d.
- MUIR-WILSON.**—THE ELEMENTS OF THERMAL CHEMISTRY. By M. M. P. MUIR, F.R.S.E.; assisted by D. M. WILSON. 8vo. 12s. 6d.
- OSTWALD.**—OUTLINES OF GENERAL CHEMISTRY: Physical and Theoretical. By Prof. W. OSTWALD. Trans. by JAS. WALKER, D.Sc. 8vo. 10s. net.
- RAMSAY.**—EXPERIMENTAL PROOFS OF CHEMICAL THEORY FOR BEGINNERS. By WILLIAM RAMSAY, F.R.S., Professor of Chemistry, University College, London. New Ed. 18mo. 2s. 6d.
- REMSEN.**—Works by IRA REMSEN, Prof. of Chemistry, Johns Hopkins University.
- \***THE ELEMENTS OF CHEMISTRY.** For Beginners. Fcap. 8vo. 2s. 6d.
- AN INTRODUCTION TO THE STUDY OF CHEMISTRY (INORGANIC CHEMISTRY). Cr. 8vo. 6s. 6d.
- COMPOUNDS OF CARBON: an Introduction to the Study of Organic Chemistry. Cr. 8vo. 6s. 6d.
- A TEXT-BOOK OF INORGANIC CHEMISTRY. 8vo. 16s.

- ROS COE.**—Works by Sir HENRY E. ROS COE, F.R.S., formerly Professor of Chemistry, Owens College, Manchester.
- \***A PRIMER OF CHEMISTRY.** Illustrated. With Questions. 18mo. 1s.
- \***INORGANIC CHEMISTRY FOR BEGINNERS.** Assisted by J. LUNT, B.Sc. Gl. 8vo. 2s. 6d.
- \***LESSONS IN ELEMENTARY CHEMISTRY, INORGANIC AND ORGANIC.** With Illustrations and Chromolitho of the Solar Spectrum, and of the Alkalies and Alkaline Earths. New Ed., 1892. Fcap. 8vo. 4s. 6d.
- ROS COE—SCHORLEMMER.**—A TREATISE ON INORGANIC AND ORGANIC CHEMISTRY. By Sir HENRY ROS COE, F.R.S., and Prof. C. SCHORLEMMER, F.R.S. 8vo.
- Vols. I. and II.—**INORGANIC CHEMISTRY.** Vol. I.—The Non-Metallic Elements. 2nd Ed. 21s. Vol. II. Two Parts, 18s. each.
- Vol. III.—**ORGANIC CHEMISTRY. THE CHEMISTRY OF THE HYDRO-CARBONS** and their Derivatives. Parts I. II. IV. and VI. 21s. each. Parts III. and V. 18s. each.
- ROS COE—SCHUSTER.**—**SPECTRUM ANALYSIS.** By Sir HENRY ROS COE, F.R.S. 4th Ed., revised by the Author and A. SCHUSTER, F.R.S., Professor of Applied Mathematics in the Owens College, Manchester. 8vo. 21s.
- SCHORLEMMER.**—**RISE AND DEVELOPMENT OF ORGANIC CHEMISTRY.** By Prof. SCHORLEMMER, N. E. Edited by Prof. A. H. SMITHELLS. [*In the Press.*]
- SCHULTZ—JULIUS.**—**SYSTEMATIC SURVEY OF THE ORGANIC COLOURING MATTERS.** By Dr. G. SCHULTZ and P. JULIUS. Translated and Edited by ARTHUR G. GREEN, F.I.C., F.C.S., Examiner in City and Guilds of London Institute. Royal 8vo. [*In the Press.*]
- \***THORPE.**—**A SERIES OF CHEMICAL PROBLEMS.** With Key. By T. E. THORPE, F.R.S., Professor of Chemistry, Royal College of Science. New Ed. Fcap. 8vo. 2s.
- ESSAYS IN HISTORICAL CHEMISTRY.** By the same. 8vo. [*In the Press.*]
- THORPE—RÜCKER.**—**A TREATISE ON CHEMICAL PHYSICS.** By Prof. T. E. THORPE and Prof. A. W. RÜCKER. 8vo. [*In preparation.*]
- \***TURPIN.**—**ORGANIC CHEMISTRY FOR BEGINNERS.** By G. S. TURPIN, M.A. Gl. 8vo. [*In the Press.*]
- WURTZ.**—**A HISTORY OF CHEMICAL THEORY.** By AD. WURTZ. Translated by HENRY WATTS, F.R.S. Crown 8vo. 6s.
- WYNNE.**—**COAL TAR PRODUCTS.** By W. P. WYNNE, Royal College of Science. [*In preparation.*]

## PHYSICAL GEOGRAPHY, GEOLOGY, AND MINERALOGY.

- BLANFORD.**—**THE RUDIMENTS OF PHYSICAL GEOGRAPHY FOR INDIAN SCHOOLS;** with Glossary. By H. F. BLANFORD, F.G.S. Cr. 8vo. 2s. 6d.
- FERREL.**—**A POPULAR TREATISE ON THE WINDS.** By W. FERREL, M.A., Member of the American National Academy of Sciences. 8vo. 17s. net.
- FISHER.**—**PHYSICS OF THE EARTH'S CRUST.** By Rev. OSMOND FISHER, M.A., F.G.S., Hon. Fellow of King's College, London. 2nd Ed., enlarged. 8vo. 12s.
- GEE.**—**SHORT STUDIES IN EARTH KNOWLEDGE.** By WILLIAM GEE. Gl. 8vo. Illustrated. [*In the Press.*]
- GEIKIE.**—Works by Sir ARCHIBALD GEIKIE, F.R.S., Director-General of the Geological Survey of the United Kingdom.
- \***A PRIMER OF PHYSICAL GEOGRAPHY.** Illus. With Questions. 18mo. 1s.
- \***ELEMENTARY LESSONS IN PHYSICAL GEOGRAPHY.** Illustrated. Fcap. 8vo. 4s. 6d. \***QUESTIONS ON THE SAME.** 1s. 6d.
- \***A PRIMER OF GEOLOGY.** Illustrated. 18mo. 1s.
- \***CLASS-BOOK OF GEOLOGY.** Illustrated. Cr. 8vo. 4s. 6d.
- TEXT-BOOK OF GEOLOGY.** Illustrated. 3rd Ed. (1893). 8vo. 28s.
- OUTLINES OF FIELD GEOLOGY.** Illustrated. Gl. 8vo. 3s. 6d.
- THE SCENERY AND GEOLOGY OF SCOTLAND, VIEWED IN CONNEXION WITH ITS PHYSICAL GEOLOGY.** Illustrated. Cr. 8vo. 12s. 6d.
- HUXLEY.**—**PHYSIOGRAPHY.** An Introduction to the Study of Nature. By the Right Hon. T. H. HUXLEY, F.R.S. Illustrated. Cr. 8vo. 6s.

- KELVIN.**—POPULAR LECTURES AND ADDRESSES. By Lord KELVIN, P.R.S. Vol. II. CONTRIBUTIONS TO GEOLOGY. [*In the Press.*]
- LESSING.**—TABLES FOR THE DETERMINATION OF THE ROCK-FORMING MINERALS. Compiled by F. L. LOEWINSON-LESSING, Professor of Geology at the University of Dorpat. Translated from the Russian by J. W. GREGORY, B.Sc., F.G.S., of the British Museum. With a Glossary added by Prof. G. A. J. COLE, F.G.S. 8vo. 4s. 6d. net.
- LOCKYER.**—OUTLINES OF PHYSIOGRAPHY—THE MOVEMENTS OF THE EARTH. By J. NORMAN LOCKYER, F.R.S., Examiner in Physiography for the Science and Art Department. Illustrated. Cr. 8vo. Sewed, 1s. 6d.
- LOUIS.**—HANDBOOK OF GOLD MILLING. By HENRY LOUIS. [*In the Press.*]
- \*MARR—HARKER.** PHYSIOGRAPHY FOR BEGINNERS. By J. E. MARR, F.R.S., and A. HARKER, M.A. Gl. 8vo. [*In the Press.*]
- MIERS.**—A TREATISE ON MINERALOGY. By H. A. MIERS, of the British Museum. 8vo. [*In preparation.*]
- MIERS—CROSSKEY.**—(See Hygiene, p. 41.)
- WILLIAMS.**—ELEMENTS OF CRYSTALLOGRAPHY, for students of Chemistry, Physics, and Mineralogy. By G. H. WILLIAMS, Ph.D. Cr. 8vo. 6s.

## BIOLOGY.

(Botany, Zoology, General Biology, Physiology.)

## Botany.

- ALLEN.**—ON THE COLOURS OF FLOWERS, as Illustrated in the British Flora. By GRANT ALLEN. Illustrated. Cr. 8vo. 3s. 6d.
- BALFOUR—WARD.**—A GENERAL TEXT-BOOK OF BOTANY. By Prof. I. B. BALFOUR, F.R.S., University of Edinburgh, and Prof. H. MARSHALL WARD, F.R.S., Roy. Indian Engineering Coll. [*In preparation.*]
- \*BETTANY.**—FIRST LESSONS IN PRACTICAL BOTANY. By G. T. BETTANY. 18mo. 1s.
- \*BOWER.**—Works by F. O. BOWER, D.Sc., F.R.S., Professor of Botany, University of Glasgow.
- A COURSE OF PRACTICAL INSTRUCTION IN BOTANY. Cr. 8vo. 10s. 6d.
- THREE MONTHS COURSE OF PRACTICAL BOTANY. [*In the Press.*]
- CHURCH—VINES.**—MANUAL OF VEGETABLE PHYSIOLOGY. By Prof. A. H. CHURCH, F.R.S., and S. H. VINES. Illustrated. Cr. 8vo. [*In prep.*]
- GOODALE.**—PHYSIOLOGICAL BOTANY. I. Outlines of the Histology of Phanogamous Plants. II. Vegetable Physiology. By G. L. GOODALE, M.A., M.D., Professor of Botany in Harvard University. 8vo. 10s. 6d.
- GRAY.**—STRUCTURAL BOTANY, OR ORGANOGRAPHY ON THE BASIS OF MORPHOLOGY. By Prof. ASA GRAY, LL.D. 8vo. 10s. 6d.
- HARTIG.**—TEXT-BOOK OF THE DISEASES OF TREES. (See Agriculture, p. 40.)
- HOOKE.**—Works by Sir JOSEPH HOOKER, F.R.S., &c.
- \*PRIMER OF BOTANY. Illustrated. 18mo. 1s.
- THE STUDENT'S FLORA OF THE BRITISH ISLANDS. 3rd Ed., revised. Gl. 8vo. 10s. 6d.
- LUBBOCK.**—FLOWERS, FRUITS, AND LEAVES. By the Right Hon. Sir J. LUBBOCK, F.R.S. Illustrated. 2nd Ed. Cr. 8vo. 4s. 6d.
- MÜLLER.**—THE FERTILISATION OF FLOWERS. By HERMANN MÜLLER. Translated by D'ARCY W. THOMPSON, B.A., Professor of Biology in University College, Dundee. Preface by CHARLES DARWIN. Illustrated. 8vo. 21s.
- NISBET.**—BRITISH FOREST TREES. (See Agriculture, p. 40.)
- OLIVER.**—\*LESSONS IN ELEMENTARY BOTANY. By DANIEL OLIVER, F.R.S., late Professor of Botany in University College, London. Fcap. 8vo. 4s. 6d.
- FIRST BOOK OF INDIAN BOTANY. By the same. Ex. fcap. 8vo. 6s. 6d.
- SMITH.**—DISEASES OF FIELD AND GARDEN CROPS. (See Agriculture, p. 40.)
- WARD.**—TIMBER AND SOME OF ITS DISEASES. (See Agriculture, p. 40.)

## Zoology.

- BADENOCH.**—THE ROMANCE OF THE INSECT WORLD. By L. N. BADENOCH. Illustr. Cr. 8vo. [*In the Press.*]
- BALFOUR.**—A TREATISE ON COMPARATIVE EMBRYOLOGY. By F. M. BALFOUR, F.R.S. Illustrated. 2 vols. 8vo. Vol. I. 18s. Vol. II. 21s.



- BERNARD**—THE APODIDAE. By H. M. BERNARD, M.A., LL.D. Cr. Svo. 7s. 6d.
- BUCKTON**—MONOGRAPH OF THE BRITISH CICADÆ, OR TETTIGIDÆ. By G. B. BUCKTON. 2 vols. Svo. 33s. 6d. each, net.
- COUES**—HANDBOOK OF FIELD AND GENERAL ORNITHOLOGY. By Prof. ELLIOTT COUES, M.A. Illustrated. Svo. 10s. net.
- FLOWER**—GADOW.—AN INTRODUCTION TO THE OSTEOLOGY OF THE MAMMALIA. By Sir W. H. FLOWER, F.R.S., Director of the Natural History Museum. Illus. 3rd Ed., revised with the help of HANS GADOW, Ph.D. Cr. Svo. 10s. 6d.
- FOSTER**—BALFOUR.—THE ELEMENTS OF EMBRYOLOGY. By Prof. MICHAEL FOSTER, M.D., F.R.S., and the late F. M. BALFOUR, F.R.S., 2nd Ed., revised by A. SEDGWICK, M.A., Fellow and Assistant Lecturer of Trinity College, Cambridge, and W. HEAPE, M.A. Illustrated. Cr. Svo. 10s. 6d.
- GÜNTHER**—GUIDE TO BRITISH FISHES. By Dr. A. GÜNTHER. Cr. Svo. [In the Press.]
- HERDMAN**—BRITISH MARINE FAUNA. Vol. I. By Prof. W. A. HERDMAN. Cr. Svo. [In the Press.]
- LANG**—TEXT-BOOK OF COMPARATIVE ANATOMY. By Dr. ARNOLD LANG, Professor of Zoology in the University of Zurich. Transl. by H. M. and M. BERNARD. Introduction by Prof. HÆCKEL. 2 vols. Illustrated. Svo. Part I 17s. net. [Part II. in the Press.]
- LUBBOCK**—THE ORIGIN AND METAMORPHOSES OF INSECTS. By the Right Hon. Sir JOHN LUBBOCK, F.R.S., D.C.L. Illus. Cr. Svo. 3s. 6d.
- MEYRICK**—BRITISH LEPIDOPTERA. By L. MEYRICK. [In preparation.]
- MIALI**—AQUATIC INSECTS. By Prof. L. C. MIALI. [In preparation.]
- MIVART**—LESSONS IN ELEMENTARY ANATOMY. By St. G. MIVART, F.R.S., Lecturer on Comparative Anatomy at St. Mary's Hospital. Fcap. Svo. 6s. 6d.
- PARKER**—A COURSE OF INSTRUCTION IN ZOOTOMY (VERTEBRATA). By T. JEFFERY PARKER, F.R.S., Professor of Biology in the University of Otago, New Zealand. Illustrated. Cr. Svo. 8s. 6d.
- PARKER**—HASWELL.—A TEXT-BOOK OF ZOOLOGY. By Prof. T. J. PARKER, F.R.S., and Prof. HASWELL. Svo. [In preparation.]
- SEDGWICK**—TREATISE ON EMBRYOLOGY. By ADAM SEDGWICK, F.R.S., Fellow and Lecturer of Trinity College, Cambridge. Svo. [In preparation.]
- SHUFELDT**—THE MYOLOGY OF THE RAVEN (*Corvus corax sinuatus*). A Guide to the Study of the Muscular System in Birds. By R. W. SHUFELDT. Illustrated. Svo. 13s. net.
- WIEDERSHEIM**—ELEMENTS OF THE COMPARATIVE ANATOMY OF VERTEBRATES. By Prof. R. WIEDERSHEIM. Adapted by W. NEWTON PARKER, Professor of Biology, University College, Cardiff. Svo. 12s. 6d.
- HUMAN ANATOMY**. Translated by H. M. BERNARD and G. B. HOWES. [In prep.]

### General Biology.

- BALL**—ARE THE EFFECTS OF USE AND DISUSE INHERITED? By W. PLATT BALL. Cr. Svo. 3s. 6d.
- BATESON**—MATERIALS FOR THE STUDY OF VARIATION IN ANIMALS. Part I. Discontinuous Variation. By W. BATESON. Svo. Illus. [In the Press.]
- CALDERWOOD**—EVOLUTION AND MAN'S PLACE IN NATURE. By Prof. H. CALDERWOOD, LL.D. Cr. Svo. 7s. 6d.
- EIMER**—ORGANIC EVOLUTION as the Result of the Inheritance of Acquired Characters according to the Laws of Organic Growth. By Dr. G. H. T. EIMER. Transl. by J. T. CUNNINGHAM, F.R.S.E. Svo. 12s. 6d.
- HOWES**—AN ATLAS OF PRACTICAL ELEMENTARY BIOLOGY. By G. B. HOWES, Assistant Professor of Zoology, Royal College of Science. 4to. 14s.
- \***HUXLEY**—INTRODUCTORY PRIMER OF SCIENCE. By Prof. T. H. HUXLEY, F.R.S. 18mo. 1s.
- HUXLEY**—MARTIN.—A COURSE OF PRACTICAL INSTRUCTION IN ELEMENTARY BIOLOGY. By Prof. T. H. HUXLEY, F.R.S., assisted by H. N. MARTIN, F.R.S., Professor of Biology in the Johns Hopkins University. New Ed., revised by G. B. HOWES, Assistant Professor, Royal College of Science, and D. H. SCOTT, D.Sc. Cr. Svo. 10s. 6d.
- LUBBOCK**—ON BRITISH WILD FLOWERS CONSIDERED IN RELATION TO INSECTS. By the Right Hon. Sir J. LUBBOCK, F.R.S. Illustrated. Cr. Svo. 4s. 6d.

- PARKER.**—LESSONS IN ELEMENTARY BIOLOGY. By Prof. T. JEFFERY PARKER, F.R.S. Illustrated. 2nd Ed. Cr. 8vo. 10s. 6d.  
**VARIGNY.**—EXPERIMENTAL EVOLUTION. By H. DE VARIGNY. Cr. 8vo. 5s.  
**WALLACE.**—Works by ALFRED RUSSEL WALLACE, F.R.S., LL.D.  
**DARWINISM:** An Exposition of the Theory of Natural Selection. Cr. 8vo. 9s.  
**NATURAL SELECTION: AND TROPICAL NATURE.** New Ed. Cr. 8vo. 6s.  
**ISLAND LIFE.** New Ed. Cr. 8vo. 6s.

### Physiology.

- FEARNLEY.**—A MANUAL OF ELEMENTARY PRACTICAL HISTOLOGY. By WILLIAM FEARNLEY. Illustrated. Cr. 8vo. 7s. 6d.  
**FOSTER.**—Works by MICHAEL FOSTER, M.D., F.R.S., Professor of Physiology in the University of Cambridge.  
 \*A PRIMER OF PHYSIOLOGY. Illustrated. 18mo. 1s.  
 A TEXT-BOOK OF PHYSIOLOGY. Illustrated. 5th Ed., largely revised. 8vo. Part I. Blood—The Tissues of Movement, The Vascular Mechanism. 10s. 6d. Part II. The Tissues of Chemical Action, with their Respective Mechanisms—Nutrition. 10s. 6d. Part III. The Central Nervous System. 7s. 6d. Part IV. The Senses and some Special Muscular Mechanisms. The Tissues and Mechanisms of Reproduction. 10s. 6d. APPENDIX—THE CHEMICAL BASIS OF THE ANIMAL BODY. By A. S. LEA, M.A. 7s. 6d.  
**FOSTER—LANGLEY.**—A COURSE OF ELEMENTARY PRACTICAL PHYSIOLOGY AND HISTOLOGY. By Prof. MICHAEL FOSTER, and J. N. LANGLEY, F.R.S., Fellow of Trinity College, Cambridge. 6th Ed. Cr. 8vo. 7s. 6d.  
**FOSTER—SHORE.**—PHYSIOLOGY FOR BEGINNERS. By MICHAEL FOSTER, M.D., F.R.S., and L. E. SHORE, M.A., M.D. Gl. 8vo. [In the Press.]  
**GAMGEE.**—A TEXT-BOOK OF THE PHYSIOLOGICAL CHEMISTRY OF THE ANIMAL BODY. By A. GAMGEE, M.D., F.R.S. 8vo. Vol. I. 18s. [Vol. II. in the Press.]  
 \***HUXLEY.**—LESSONS IN ELEMENTARY PHYSIOLOGY. By Prof. T. H. HUXLEY, F.R.S. Illust. Fcap. 8vo. 4s. 6d.  
 \***QUESTIONS ON THE ABOVE.** By T. ALCOCK, M.D. 18mo. 1s. 6d.

### MEDICINE.

- BLYTH.**—(See Hygiene, p. 41.)  
**BRUNTON.**—Works by T. LAUDER BRUNTON, M.D., F.R.S., Examiner in Materia Medica in the University of London, in the Victoria University, and in the Royal College of Physicians, London.  
 A TEXT-BOOK OF PHARMACOLOGY, THERAPEUTICS, AND MATERIA MEDICA. Adapted to the United States Pharmacopœia by F. H. WILLIAMS, M.D., Boston, Mass. 3rd Ed. Adapted to the New British Pharmacopœia, 1885, and additions, 1891. 8vo. 21s. Or in 2 vols. 22s. 6d. Supplement. 1s.  
**TABLES OF MATERIA MEDICA:** A Companion to the Materia Medica Museum. Illustrated. Cheaper Issue. 8vo. 5s.  
 AN INTRODUCTION TO MODERN THERAPEUTICS. 8vo. 3s. 6d. net.  
**GRIFFITHS.**—LESSONS ON PRESCRIPTIONS AND THE ART OF PRESCRIBING. By W. H. GRIFFITHS. Adapted to the Pharmacopœia, 1885. 18mo. 3s. 6d.  
**HAMILTON.**—A TEXT-BOOK OF PATHOLOGY, SYSTEMATIC AND PRACTICAL. By D. J. HAMILTON, F.R.S.E., Professor of Pathological Anatomy, University of Aberdeen. Illustrated. 8vo. Vol. I. 25s. [Vol. II. in the Press.]  
**KLEIN.**—Works by E. KLEIN, F.R.S., Lecturer on General Anatomy and Physiology in the Medical School of St. Bartholomew's Hospital, London.  
**MICRO-ORGANISMS AND DISEASE.** An Introduction into the Study of Specific Micro-Organisms. Illustrated. 3rd Ed., revised. Cr. 8vo. 6s.  
**THE BACTERIA IN ASIATIC CHOLERA.** Cr. 8vo. 5s.  
**VON KAHLDEN.**—HANDBOOK OF HISTOLOGICAL METHODS. By Dr. VON KAHLDEN. Translated by H. MORLEY FLETCHER, M.D. 8vo. Being a Companion to Ziegler's "Pathological Anatomy." [In preparation.]  
**WHITE.**—A TEXT-BOOK OF GENERAL THERAPEUTICS. By W. HALE WHITE, M.D., Senior Assistant Physician to and Lecturer in Materia Medica at Guy's Hospital. Illustrated. Cr. 8vo. 8s. 6d.  
**WILLOUGHBY.**—(See Hygiene, p. 41.)

**ZIEGLER—MACALISTER.**—TEXT-BOOK OF PATHOLOGICAL ANATOMY AND PATHOGENESIS. By Prof. E. ZIEGLER. Translated and Edited by DONALD MACALISTER, M.A., M.D., Fellow and Medical Lecturer of St. John's College, Cambridge. Illustrated. 8vo.

Part I.—GENERAL PATHOLOGICAL ANATOMY. 2nd Ed. 12s. 6d.

Part II.—SPECIAL PATHOLOGICAL ANATOMY. Sections I.—VIII. 2nd Ed. 12s. 6d. Sections IX.—XII. 12s. 6d.

## HUMAN SCIENCES.

Mental and Moral Philosophy ; Political Economy ; Law and Politics ;  
Anthropology ; Education.

### MENTAL AND MORAL PHILOSOPHY.

**BALDWIN.**—HANDBOOK OF PSYCHOLOGY : SENSES AND INTELLECT

By Prof. J. M. BALDWIN, M.A., LL.D. 2nd Ed., revised. 8vo. 12s. 6d.

FEELING AND WILL. By the same. 8vo. 12s. 6d.

ELEMENTS OF PSYCHOLOGY. Cr. 8vo. 7s. 6d.

**BOOLE.**—THE MATHEMATICAL ANALYSIS OF LOGIC. Being an Essay towards a Calculus of Deductive Reasoning. By GEORGE BOOLE. 8vo. 5s.

**CALDERWOOD.**—HANDBOOK OF MORAL PHILOSOPHY. By Rev. HENRY CALDERWOOD, LL.D., Professor of Moral Philosophy in the University of Edinburgh. 14th Ed., largely rewritten. Cr. 8vo. 6s.

**CLIFFORD.**—SEEING AND THINKING. By the late Prof. W. K. CLIFFORD, F.R.S. With Diagrams. Cr. 8vo. 3s. 6d.

**HILL.**—GENETIC PHILOSOPHY. By DAVID J. HILL.

[In the Press.

**HÖFFDING.**—OUTLINES OF PSYCHOLOGY. By Prof. H. HÖFFDING. Translated by M. E. LOWNDES. Cr. 8vo. 6s.

**JAMES.**—THE PRINCIPLES OF PSYCHOLOGY. By WM. JAMES, Professor of Psychology in Harvard University. 2 vols. 8vo. 25s. net.

A TEXT-BOOK OF PSYCHOLOGY. By the same. Cr. 8vo. 7s. net.

**JARDINE.**—THE ELEMENTS OF THE PSYCHOLOGY OF COGNITION. By Rev. ROBERT JARDINE, D.Sc. 3rd Ed., revised. Cr. 8vo. 6s. 6d.

**JEVONS.**—Works by W. STANLEY JEVONS, F.R.S.

\*A PRIMER OF LOGIC. 18mo. 1s.

\*ELEMENTARY LESSONS IN LOGIC, Deductive and Inductive, with Copious Questions and Examples, and a Vocabulary. Fcap. 8vo. 3s. 6d.

THE PRINCIPLES OF SCIENCE. Cr. 8vo. 12s. 6d.

STUDIES IN DEDUCTIVE LOGIC. 2nd Ed. Cr. 8vo. 6s.

PURE LOGIC: AND OTHER MINOR WORKS. Edited by R. ADAMSON, M.A., LL.D., Professor of Logic at Owens College, Manchester, and HARRIET A. JEVONS. With a Preface by Prof. ADAMSON. 8vo. 10s. 6d.

**KANT—MAX MÜLLER.**—CRITIQUE OF PURE REASON. By IMMANUEL KANT. 2 vols. 8vo. 16s. each. Vol. I. HISTORICAL INTRODUCTION, by LUDWIG NOIRÉ ; Vol. II. CRITIQUE OF PURE REASON, translated by F. MAX MÜLLER.

**KANT—MAHAFFY—BERNARD.**—KANT'S CRITICAL PHILOSOPHY FOR ENGLISH READERS. By J. P. MAHAFFY, D.D., Professor of Ancient History in the University of Dublin, and JOHN H. BERNARD, B.D., Fellow of Trinity College, Dublin. A new and complete Edition in 2 vols. Cr. 8vo.

Vol. I. THE KRITIK OF PURE REASON EXPLAINED AND DEFENDED. 7s. 6d.

Vol. II. THE PROLEGOMENA. Translated with Notes and Appendices. 6s.

**KANT.**—KRITIK OF JUDGMENT. Translated with Introduction and Notes by J. H. BERNARD, D.D. 8vo. 10s. net.

**KEYNES.**—FORMAL LOGIC, Studies and Exercises in. By J. N. KEYNES, D.Sc. 2nd Ed., revised and enlarged. Cr. 8vo. 10s. 6d.

**MCCOSH.**—Works by JAMES MCCOSH, D.D., President of Princeton College.

PSYCHOLOGY. Cr. 8vo. I. THE COGNITIVE POWERS. 6s. 6d. II. THE MOTIVE POWERS. 6s. 6d.

FIRST AND FUNDAMENTAL TRUTHS: a Treatise on Metaphysics. 8vo. 9s.

THE PREVAILING TYPES OF PHILOSOPHY. CAN THEY LOGICALLY REACH REALITY? 8vo. 3s. 6d.



- MARSHALL.**—PAIN, PLEASURE, AND ÆSTHETICS. An Essay by H. R. MARSHALL, M.A. [In the Press.]
- MAURICE.**—MORAL AND METAPHYSICAL PHILOSOPHY. By F. D. MAURICE, M.A., late Professor of Moral Philosophy in the University of Cambridge. 4th Ed. 2 vols. 8vo. 16s.
- \***RAY.**—A TEXT-BOOK OF DEDUCTIVE LOGIC FOR THE USE OF STUDENTS. By P. K. RAY, D.Sc., Professor of Logic and Philosophy, Presidency College, Calcutta. 4th Ed. Globe 8vo. 4s. 6d.
- SIDGWICK.**—Works by HENRY SIDGWICK, LL.D., D.C.L., Knightbridge Professor of Moral Philosophy in the University of Cambridge.
- THE METHODS OF ETHICS. 4th Ed. 8vo. 14s.
- OUTLINES OF THE HISTORY OF ETHICS. 3rd Ed. Cr. 8vo. 3s. 6d.
- VENN.**—Works by JOHN VENN, F.R.S., Examiner in Moral Philosophy in the University of London.
- THE LOGIC OF CHANCE. An Essay on the Foundations and Province of the Theory of Probability. 3rd Ed., rewritten and enlarged. Cr. 8vo. 10s. 6d.
- SYMBOLIC LOGIC. Cr. 8vo. 10s. 6d.
- THE PRINCIPLES OF EMPIRICAL OR INDUCTIVE LOGIC. 8vo. 18s.
- WILLIAMS.**—REVIEW OF THE SYSTEM OF ETHICS FOUNDED ON THE THEORY OF EVOLUTION. By C. M. WILLIAMS. Ex. Cr. 8vo. 12s. net.
- WINDELBAND.**—HISTORY OF PHILOSOPHY. By Dr. W. WINDELBAND. Translated by Prof. J. A. TUFTS. [In the Press.]

## POLITICAL ECONOMY.

- BASTABLE.**—PUBLIC FINANCE. By C. F. BASTABLE, Professor of Political Economy in the University of Dublin. 8vo. 12s. 6d. net.
- BÖHM-BAWERK.**—CAPITAL AND INTEREST. Translated by WILLIAM SMART, M.A. 8vo. 12s. net.
- THE POSITIVE THEORY OF CAPITAL. By the same. 8vo. 12s. net.
- CAIRNES.**—THE CHARACTER AND LOGICAL METHOD OF POLITICAL ECONOMY. By J. E. CAIRNES. Cr. 8vo. 6s.
- SOME LEADING PRINCIPLES OF POLITICAL ECONOMY NEWLY EXPOUNDED. By the same. 8vo. 14s.
- CLARE.**—THE ABC OF THE FOREIGN EXCHANGES. By GEORGE CLARE. Crown 8vo. 3s. net.
- COMMONS.**—DISTRIBUTION OF WEALTH. By Prof. J. R. COMMONS. [In the Press.]
- COSSA.**—INTRODUCTION TO THE STUDY OF POLITICAL ECONOMY. By LUIGI COSSA, Professor in the Royal University of Pavia. Translated by LOUIS DYER, M.A. Cr. 8vo. 8s. 6d. net.
- \***FAWCETT.**—POLITICAL ECONOMY FOR BEGINNERS, WITH QUESTIONS. By Mrs. HENRY FAWCETT. 7th Ed. 18mo. 2s. 6d.
- FAWCETT.**—A MANUAL OF POLITICAL ECONOMY. By the Right Hon. HENRY FAWCETT, F.R.S. 7th Ed., revised. Cr. 8vo. 12s.
- AN EXPLANATORY DIGEST of above. By C. A. WATERS, B.A. Cr. 8vo. 2s. 6d.
- GILMAN.**—PROFIT-SHARING BETWEEN EMPLOYER AND EMPLOYEE. By N. P. GILMAN. Cr. 8vo. 7s. 6d.
- SOCIALISM AND THE AMERICAN SPIRIT. By the Same. Cr. 8vo. 6s. 6d.
- GUNTON.**—WEALTH AND PROGRESS: An examination of the Wages Question and its Economic Relation to Social Reform. By GEORGE GUNTON. Cr. 8vo. 6s.
- HOWELL.**—THE CONFLICTS OF CAPITAL AND LABOUR HISTORICALLY AND ECONOMICALLY CONSIDERED. Being a History and Review of the Trade Unions of Great Britain. By GEORGE HOWELL, M.P. 2nd Ed., revised. Cr. 8vo. 7s. 6d.
- JEVONS.**—Works by W. STANLEY JEVONS, F.R.S.
- \*PRIMER OF POLITICAL ECONOMY. 18mo. 1s.
- THE THEORY OF POLITICAL ECONOMY. 3rd Ed., revised. 8vo. 10s. 6d.
- KEYNES.**—THE SCOPE AND METHOD OF POLITICAL ECONOMY. By J. N. KEYNES, D.Sc. 7s. net.
- MARSHALL.**—PRINCIPLES OF ECONOMICS. By ALFRED MARSHALL, M.A., Professor of Political Economy in the University of Cambridge. 2 vols. 8vo. Vol. I. 2nd Ed. 12s. 6d. net.

- ELEMENTS OF ECONOMICS OF INDUSTRY.** By the same. New Ed., 1892. Cr. 8vo. 3s. 6d.
- PALGRAVE.**—A DICTIONARY OF POLITICAL ECONOMY. By various Writers. Edited by R. H. INGLIS PALGRAVE, F.R.S. 3s. 6d. each, net. No. 1. *July* 1891.
- SIDGWICK.**—THE PRINCIPLES OF POLITICAL ECONOMY. By HENRY SIDGWICK, LL.D., D.C.L., Knightbridge Professor of Moral Philosophy in the University of Cambridge. 2nd Ed., revised. 8vo. 16s.
- SMART.**—AN INTRODUCTION TO THE THEORY OF VALUE. By WILLIAM SMART, M.A. Crown 8vo. 3s. net.
- THOMPSON.**—THE THEORY OF WAGES. By H. M. THOMPSON. Cr. 8vo. 3s. 6d.
- WALKER.**—Works by FRANCIS A. WALKER, M.A.  
**FIRST LESSONS IN POLITICAL ECONOMY.** Cr. 8vo. 5s.  
**A BRIEF TEXT-BOOK OF POLITICAL ECONOMY.** Cr. 8vo. 6s. 6d.  
**POLITICAL ECONOMY.** 2nd Ed., revised and enlarged. 8vo. 12s. 6d.  
**THE WAGES QUESTION.** Ex. Cr. 8vo. 8s. 6d. net.
- MONEY.** Ex. Cr. 8vo. 8s. 6d. net.
- WICKSTEED.**—ALPHABET OF ECONOMIC SCIENCE. By P. H. WICKSTEED, M.A. Part I. Elements of the Theory of Value or Worth. Gl. 8vo. 2s. 6d.
- WIESER.**—NATURAL VALUE. By Prof. F. von WIESER. Translated by W. MALLOCK. Edited by W. SMART, M.A. 8vo. [*In the Press.*]

## LAW AND POLITICS.

- BALL.**—THE STUDENT'S GUIDE TO THE BAR. By W. W. ROUSE BALL, M.A., Fellow of Trinity College, Cambridge. 4th Ed., revised. Cr. 8vo. 2s. 6d.
- BOUTMY.**—STUDIES IN CONSTITUTIONAL LAW. By EMILE BOUTMY. Translated by Mrs. DICEY, with Preface by Prof. A. V. DICEY. Cr. 8vo. 6s.
- THE ENGLISH CONSTITUTION.** By the same. Translated by Mrs. EADEN, with Introduction by Sir F. POLLOCK, Bart. Cr. 8vo. 6s.
- \*BUCKLAND.**—OUR NATIONAL INSTITUTIONS. By A. BUCKLAND. 18mo. 1s.
- CHERRY.**—LECTURES ON THE GROWTH OF CRIMINAL LAW IN ANCIENT COMMUNITIES. By R. R. CHERRY, LL.D., Reid Professor of Constitutional and Criminal Law in the University of Dublin. 8vo. 5s. net.
- DICEY.**—INTRODUCTION TO THE STUDY OF THE LAW OF THE CONSTITUTION. By A. V. DICEY, B.C.L., Vinerian Professor of English Law in the University of Oxford. 3rd Ed. 8vo. 12s. 6d.
- HOLMES.**—THE COMMON LAW. By O. W. HOLMES, Jun. Demy 8vo. 12s.
- JENKS.**—THE GOVERNMENT OF VICTORIA. By EDWARD JENKS, B.A., LL.B., late Professor of Law in the University of Melbourne. 14s.
- MUNRO.**—COMMERCIAL LAW. (*See Commerce*, p. 41.)
- PHILLIMORE.**—PRIVATE LAW AMONG THE ROMANS. From the Pandects. By J. G. PHILLIMORE, Q.C. 8vo. 16s.
- POLLOCK.**—ESSAYS IN JURISPRUDENCE AND ETHICS. By Sir FREDERICK POLLOCK, Bart. 8vo. 10s. 6d.
- INTRODUCTION TO THE HISTORY OF THE SCIENCE OF POLITICS.** By the same. Cr. 8vo. 2s. 6d.
- SIDGWICK.**—THE ELEMENTS OF POLITICS. By HENRY SIDGWICK, LL.D. 8vo. 14s. net.
- STEPHEN.**—Works by Sir JAMES FITZJAMES STEPHEN, Bart.  
**A DIGEST OF THE LAW OF EVIDENCE.** 5th Ed. Cr. 8vo. 6s.  
**A DIGEST OF THE CRIMINAL LAW: CRIMES AND PUNISHMENTS.** 4th Ed., revised. 8vo. 16s.  
**A DIGEST OF THE LAW OF CRIMINAL PROCEDURE IN INDICTABLE OFFENCES.** By Sir J. F. STEPHEN, Bart., and H. STEPHEN. 8vo. 12s. 6d.  
**A HISTORY OF THE CRIMINAL LAW OF ENGLAND.** 3 vols. 8vo. 48s.  
**A GENERAL VIEW OF THE CRIMINAL LAW OF ENGLAND.** 8vo. 14s.
- \*WYATT.**—THE ENGLISH CITIZEN, HIS LIFE AND DUTIES. By C. H. WYATT, Clerk to the Manchester School Board. Gl. 8vo. 2s.

## ANTHROPOLOGY.

- TYLOR.**—ANTHROPOLOGY. By E. B. TYLOR, F.R.S., Reader in Anthropology in the University of Oxford. Illustrated. Cr. 8vo. 7s. 6d.

## EDUCATION.

- ARNOLD.**—REPORTS ON ELEMENTARY SCHOOLS. 1852-1882. By MATTHEW ARNOLD. Edited by Lord SANDFORD. Cr. 8vo. 2s. 6d.  
**HIGHER SCHOOLS AND UNIVERSITIES IN GERMANY.** By the same. Crown 8vo. 6s.  
**A FRENCH ETON, AND HIGHER SCHOOLS AND UNIVERSITIES IN FRANCE.** By the same. Cr. 8vo. 6s.  
**BALL.**—THE STUDENT'S GUIDE TO THE BAR. (*See* Law, p. 38.)  
**\*BLAKISTON.**—THE TEACHER. Hints on School Management. By J. R. BLAKISTON, H.M.I.S. Cr. 8vo. 2s. 6d.  
**CALDERWOOD.**—ON TEACHING. By Prof. HENRY CALDERWOOD. New Ed. Ex. fcap. 8vo. 2s. 6d.  
**FEARON.**—SCHOOL INSPECTION. By D. R. FEARON. 6th Ed. Cr. 8vo. 2s. 6d.  
**FITCH.**—NOTES ON AMERICAN SCHOOLS AND TRAINING COLLEGES. By J. G. FITCH, M.A., LL.D. Gl. 8vo. 2s. 6d.  
**GEIKIE.**—THE TEACHING OF GEOGRAPHY. (*See* Geography, p. 41.)  
**GLADSTONE.**—SPELLING REFORM FROM A NATIONAL POINT OF VIEW. By J. H. GLADSTONE. Cr. 8vo. 1s. 6d.  
**HERTEL.**—OVERPRESSURE IN HIGH SCHOOLS IN DENMARK. By Dr. HERTEL. Introd. by Sir J. CRICHTON-BROWNE, F.R.S. Cr. 8vo. 3s. 6d.  
**RECORD OF TECHNICAL AND SECONDARY EDUCATION.** Bi-Monthly. 8vo. Sewed, 2s. net. Part I. Nov. 1891.

## TECHNICAL KNOWLEDGE.

Civil and Mechanical Engineering; Military and Naval Science;  
 Agriculture; Domestic Economy; Hygiene; Commerce; Manual Training.

### CIVIL AND MECHANICAL ENGINEERING.

- ALEXANDER—THOMSON.**—ELEMENTARY APPLIED MECHANICS. (*See* p. 26.)  
**CHALMERS.**—GRAPHICAL DETERMINATION OF FORCES IN ENGINEERING STRUCTURES. By J. B. CHALMERS, C.E. Illustrated. 8vo. 24s.  
**COTTERILL.**—APPLIED MECHANICS. (*See* p. 27.)  
**COTTERILL—SLADE.**—LESSONS IN APPLIED MECHANICS. (*See* p. 27.)  
**GRAHAM.**—GEOMETRY OF POSITION. (*See* p. 27.)  
**KENNEDY.**—THE MECHANICS OF MACHINERY. (*See* p. 27.)  
**LANGMAID—GAISFORD.**—ELEMENTARY LESSONS IN STEAM MACHINERY AND IN MARINE STEAM ENGINES. By T. LANGMAID, Chief Engineer R.N., and H. GAISFORD, R.N. [*Shortly.*]  
**PEABODY.**—THERMODYNAMICS OF THE STEAM-ENGINE AND OTHER HEAT-ENGINES. (*See* p. 29.)  
**SHANN.**—AN ELEMENTARY TREATISE ON HEAT IN RELATION TO STEAM AND THE STEAM-ENGINE. (*See* p. 29.)  
**YOUNG.**—SIMPLE PRACTICAL METHODS OF CALCULATING STRAINS ON GIRDERS, ARCHES, AND TRUSSES. By E. W. YOUNG, C.E. 8vo. 7s. 6d.  
**WIESBACH—HERMANN.**—THE MECHANICS OF HOISTING MACHINERY. By Dr. JULIUS WIESBACH and Professor GUSTAV HERMANN. Translated by K.P. DAHLSTROM. [*In the Press.*]

### MILITARY AND NAVAL SCIENCE.

- ARMY PRELIMINARY EXAMINATION PAPERS, 1886-1893.** (*See* Mathematics.)  
**FLAGG.**—A PRIMER OF NAVIGATION. By A. T. FLAGG. 18mo. [*In preparation.*]  
**KELVIN.**—POPULAR LECTURES AND ADDRESSES. By Lord KELVIN, P.R.S. 3 vols. Illustrated. Cr. 8vo. Vol. III. Navigation. 7s. 6d.  
**MATTHEWS.**—MANUAL OF LOGARITHMS. (*See* Mathematics, p. 24.)  
**MAURICE.**—WAR. By Col. G. F. MAURICE, C.B., R.A. 8vo. 5s. net.  
**MERCUR.**—ELEMENTS OF THE ART OF WAR. Prepared for the use of Cadets of the United States Military Academy. By JAMES MERCUR. 8vo. 17s.  
**PALMER.**—TEXT-BOOK OF PRACTICAL LOGARITHMS AND TRIGONOMETRY. (*See* Mathematics, p. 24.)  
**ROBINSON.**—TREATISE ON MARINE SURVEYING. For younger Naval Officers. By Rev. J. L. ROBINSON. Cr. 8vo. 7s. 6d.



- SANDHURST MATHEMATICAL PAPERS.** (*See Mathematics*, p. 25.)  
**SHORTLAND.**—**NAUTICAL SURVEYING.** By Vice-Adm. **SHORTLAND.** 8vo. 21s.  
**WOLSELEY.**—**Works by General Viscount WOLSELEY, G.C.M.G.**  
**THE SOLDIER'S POCKET-BOOK FOR FIELD SERVICE.** 16mo. Roan. 5s  
**FIELD POCKET-BOOK FOR THE AUXILIARY FORCES.** 16mo. 1s. 6d.  
**WOOLWICH MATHEMATICAL PAPERS.** (*See Mathematics*, p. 25.)

## AGRICULTURE AND FORESTRY.

- FRANKLAND.**—**AGRICULTURAL CHEMICAL ANALYSIS.** By P. F. **FRANKLAND**, F.R.S., Prof. of Chemistry, University College, Dundee. Cr. 8vo. 7s. 6d.  
**HARTIG.**—**TEXT-BOOK OF THE DISEASES OF TREES.** By Dr. **ROBERT HARTIG.** Translated by Wm. **SOMERVILLE**, B.S., D.C.E., Professor of Agriculture and Forestry, Durham College of Science. 8vo. [*In the Press.*]  
**LASLETT.**—**TIMBER AND TIMBER TREES, NATIVE AND FOREIGN.** By **THOMAS LASLETT.** [*New Edition in Preparation.*]  
**LAURIE.**—**THE FOOD OF PLANTS.** By A. P. **LAURIE**, M.A. 18mo. 1s.  
**MUIR.**—**MANUAL OF DAIRY-WORK.** By Professor **JAMES MUIR**, Yorkshire College, Leeds. 18mo. 1s.  
**NICHOLLS.**—**A TEXT-BOOK OF TROPICAL AGRICULTURE.** By H. A. **ALFORD NICHOLLS**, M.D. Illustrated. Crown 8vo. 6s.  
**NISBET.**—**BRITISH FOREST TREES AND THEIR AGRICULTURAL CHARACTERISTICS AND TREATMENT.** By **JOHN NISBET**, D.C.E., of the Indian Forest Service. Cr. 8vo. 6s. net.  
**SOMERVILLE.**—**INSECTS IN RELATION TO AGRICULTURE.** By Dr. W. **SOMERVILLE.** 18mo. [*In preparation.*]  
**SMITH.**—**DISEASES OF FIELD AND GARDEN CROPS**, chiefly such as are caused by Fungi. By **WORTHINGTON G. SMITH**, F.L.S. Fcap. 8vo. 4s. 6d.  
**TANNER.**—**\*ELEMENTARY LESSONS IN THE SCIENCE OF AGRICULTURAL PRACTICE.** By **HENRY TANNER**, F.C.S., M.R.A.C., Examiner in Agriculture under the Science and Art Department. Fcap. 8vo. 3s. 6d.  
**\*FIRST PRINCIPLES OF AGRICULTURE.** By the same. 18mo. 1s.  
**\*THE PRINCIPLES OF AGRICULTURE.** For use in Elementary Schools. By the same. Ex. fcap. 8vo. I. The Alphabet. 6d. II. Further Steps. 1s. III. Elementary School Readings for the Third Stage. 1s.  
**WARD.**—**TIMBER AND SOME OF ITS DISEASES.** By H. **MARSHALL WARD**, F.R.S., Prof. of Botany, Roy. Ind. Engin. Coll., Cooper's Hill. Cr. 8vo. 6s.  
**WRIGHT.**—**A PRIMER OF PRACTICAL HORTICULTURE.** By J. **WRIGHT**, F.R.H.S. 18mo. 1s.

## DOMESTIC ECONOMY.

- \*BARKER.**—**FIRST LESSONS IN THE PRINCIPLES OF COOKING.** By **LADY BARKER.** 18mo. 1s.  
**\*BARNETT-O'NEILL.**—**A PRIMER OF DOMESTIC ECONOMY.** By E. A. **BARNETT** and H. C. **O'NEILL.** 18mo. 1s.  
**\*COOKERY BOOK.**—**THE MIDDLE-CLASS COOKERY BOOK.** Edited by the Manchester School of Domestic Cookery. Fcap. 8vo. 1s. 6d.  
**CRAVEN.**—**A GUIDE TO DISTRICT NURSES.** By **MRS. CRAVEN.** Cr. 8vo. 2s. 6d.  
**\*GRAND'HOMME.**—**CUTTING-OUT AND DRESSMAKING.** From the French of **Mdlle. E. GRAND'HOMME.** With Diagrams. 18mo. 1s.  
**\*GRENFELL.**—**DRESSMAKING.** A Technical Manual for Teachers. By **MRS. HENRY GRENFELL.** With Diagrams. 18mo. 1s.  
**JEX-BLAKE.**—**THE CARE OF INFANTS.** A Manual for Mothers and Nurses. By **SOPHIA JEX-BLAKE**, M.D. 18mo. 1s.  
**ROSEVEAR.**—**MANUAL OF NEEDLEWORK.** By E. **ROSEVEAR**, Lecturer on Needlework, Training College, Stockwell. Cr. 8vo. 6s.  
**\*TEGETMEIER.**—**HOUSEHOLD MANAGEMENT AND COOKERY.** Compiled for the London School Board. By W. B. **TEGETMEIER.** 18mo. 1s.  
**\*WRIGHT.**—**THE SCHOOL COOKERY-BOOK.** Compiled and Edited by C. E. **GUTHRIE WRIGHT**, Hon. Sec. to the Edinburgh School of Cookery. 18mo. 1s.

## HYGIENE.

- \*BERNERS.**—**FIRST LESSONS ON HEALTH.** By J. **BERNERS.** 18mo. 1s.

- BLYTH.**—A MANUAL OF PUBLIC HEALTH. By A. WYNTER BLYTH. M.R.C.S. 8vo. 17s. net.  
**LECTURES ON SANITARY LAW.** By the same. 8vo. 8s. 6d. net.  
**MIERS—CROSSKEY.**—THE SOIL IN RELATION TO HEALTH. By H. A. MIERS, M.A., F.G.S., F.C.S., and R. CROSSKEY, M.A., D.P.H. Cr. 8vo. 3s. 6d.  
**REYNOLDS.**—A PRIMER OF HYGIENE. By E. S. REYNOLDS, M.D., Victoria University Extension Lecturer in Hygiene. 18mo. *[In preparation.]*  
**WILLOUGHBY.**—HANDBOOK OF PUBLIC HEALTH AND DEMOGRAPHY. By Dr. E. F. WILLOUGHBY. Fcap. 8vo. 4s. 6d.

## COMMERCE.

- MACMILLAN'S ELEMENTARY COMMERCIAL CLASS BOOKS.** Edited by JAMES GOW, Litt.D., Headmaster of the High School, Nottingham. Globe 8vo.  
**\*THE HISTORY OF COMMERCE IN EUROPE.** By H. DE B. GIBBINS, M.A. 8s. 6d.  
**\*COMMERCIAL ARITHMETIC.** By S. JACKSON, M.A. 3s. 6d.  
**THE STUDENT'S MANUAL OF BOOKKEEPING.** By J. THORNTON. *[In preparation.]*  
**COMMERCIAL GEOGRAPHY.** By E. C. K. GONNER, M.A., Professor of Political Economy in University College, Liverpool. *[In preparation.]*  
**\*INTRODUCTION TO COMMERCIAL GERMAN.** By F. C. SMITH, B.A., formerly Scholar of Magdalene College, Cambridge. 3s. 6d.  
**COMMERCIAL FRENCH.** By JAMES B. PAYNE, King's College School, London. *[In preparation.]*  
**COMMERCIAL SPANISH.** By Prof. DELBOS, Instructor, H.M.S. *Britannia*, Dartmouth. *[In the Press.]*  
**\*COMMERCIAL LAW.** By J. E. C. MUNRO, LL.D., late Professor of Law and Political Economy in the Owens College, Manchester. 3s. 6d.

## TECHNOLOGY.

- BENSON.**—ELEMENTARY HANDICRAFT AND DESIGN. By W. A. S. BENSON. Illustrated. Cr. 8vo. 5s. net.  
**BURDETT.**—BOOT AND SHOE MANUFACTURE. By C. W. B. BURDETT. Illustrated. Cr. 8vo. *[In the Press.]*  
**\*DEGERDON.**—THE GRAMMAR OF WOODWORK. By W. E. DEGERDON, Head Instructor, Whitechapel Craft School. 4to. 2s. sewed; 3s. cloth.  
**LETHABY.**—CAST IRON AND LEAD WORK. By W. R. LETHABY. Illustrated. Cr. 8vo. *[In preparation.]*  
**LOUIS.**—GOLD MILLING. By H. LOUIS. Cr. 8vo. *[In the Press.]*  
**VICKERMAN.**—WOOLLEN SPINNING. By CHARLES VICKERMAN. Illustrated. Cr. 8vo. *[In the Press.]*

## GEOGRAPHY.

(See also PHYSICAL GEOGRAPHY, p. 32.)

- BARTHOLOMEW.**—\*THE ELEMENTARY SCHOOL ATLAS. By JOHN BARTHOLOMEW, F.R.G.S. 4to. 1s.  
**\*MACMILLAN'S SCHOOL ATLAS, PHYSICAL AND POLITICAL.** 80 Maps and Index. By the same. Royal 4to. 8s. 6d. Half-morocco, 10s. 6d.  
**THE LIBRARY REFERENCE ATLAS OF THE WORLD.** By the same. 84 Maps and Index to 100,000 places. Half-morocco. Gilt edges. Folio. £2:12:6 net. Also in parts, 5s. each, net. Index, 7s. 6d. net.  
**\*CLARKE.**—CLASS-BOOK OF GEOGRAPHY. By C. B. CLARKE, F.R.S. With 18 Maps. Fcap. 8vo. 3s.; sewed, 2s. 6d.  
**\*GREEN.**—A SHORT GEOGRAPHY OF THE BRITISH ISLANDS. By JOHN RICHARD GREEN, LL.D., and A. S. GREEN. With Maps. Fcap. 8vo. 3s. 6d.  
**\*GROVE.**—A PRIMER OF GEOGRAPHY. By Sir GEORGE GROVE. 18mo. 1s.  
**KIEPERT.**—A MANUAL OF ANCIENT GEOGRAPHY. By Dr. H. KIEPERT. Cr. 8vo. 5s.  
**MACMILLAN'S GEOGRAPHICAL SERIES.**—Edited by Sir ARCHIBALD GEIKIE, F.R.S., Director-General of the Geological Survey of the United Kingdom.  
**\*THE TEACHING OF GEOGRAPHY.** A Practical Handbook for the Use of Teachers. By Sir ARCHIBALD GEIKIE, F.R.S. Cr. 8vo. 2s.

- \*MAPS AND MAP-DRAWING. By W. A. ELDERTON. 18mo. 1s.  
 \*GEOGRAPHY OF THE BRITISH ISLES. By Sir A. GEIKIE, F.R.S. 18mo. 1s.  
 \*AN ELEMENTARY CLASS-BOOK OF GENERAL GEOGRAPHY. By H. R. MILL, D.Sc. Illustrated. Cr. 8vo. 3s. 6d.  
 \*GEOGRAPHY OF EUROPE. By J. SIME, M.A. Illustrated. Gl. 8vo. 3s.  
 \*ELEMENTARY GEOGRAPHY OF INDIA, BURMA, AND CEYLON. By H. F. BLANFORD, F.G.S. Gl. 8vo. 2s. 6d.  
 GEOGRAPHY OF NORTH AMERICA. By Prof. N. S. SHALER. [*In preparation.*]  
 \*ELEMENTARY GEOGRAPHY OF THE BRITISH COLONIES. By G. M. DAWSON, LL.D., and A. SUTHERLAND. Globe 8vo. 3s.  
 STRACHEY.—LECTURES ON GEOGRAPHY. By General RICHARD STRACHEY, R.E. Cr. 8vo. 4s. 6d.  
 SUTHERLAND.—GEOGRAPHY OF VICTORIA. By A. SUTHERLAND. Pott. 8vo. 1s.  
 GENERAL GEOGRAPHY. For use in Elementary Schools in Victoria. By the same. [*In the Press.*]  
 \*TOZER.—A PRIMER OF CLASSICAL GEOGRAPHY. By H. F. TOZER, M.A. 18mo 1s.

## HISTORY.

- ARNOLD.—THE SECOND PUNIC WAR. (*See Classics, p. 11.*)  
 ARNOLD.—A HISTORY OF THE EARLY ROMAN EMPIRE. (*See p. 11.*)  
 \*BEESLY.—STORIES FROM THE HISTORY OF ROME. (*See p. 12.*)  
 BRYCE.—THE HOLY ROMAN EMPIRE. By JAMES BRYCE, M.P., D.C.L., Cr. 8vo. 7s. 6d. Library Edition. 8vo. 14s.  
 \*BUCKLEY.—A HISTORY OF ENGLAND FOR BEGINNERS. By ARABELLA B. BUCKLEY. With Maps and Tables. Gl. 8vo. 3s.  
 BURY.—A HISTORY OF THE LATER ROMAN EMPIRE FROM ARCADIUS TO IRENE. (*See Classics, p. 12.*)  
 CASSEL.—MANUAL OF JEWISH HISTORY AND LITERATURE. By Dr. D. CASSEL. Translated by Mrs. HENRY LUCAS. Fcap. 8vo. 2s. 6d.  
 ENGLISH STATESMEN, TWELVE. Cr. 8vo. 2s. 6d. each.  
 WILLIAM THE CONQUEROR. By EDWARD A. FREEMAN, D.C.L., LL.D.  
 HENRY II. By Mrs. J. R. GREEN.  
 EDWARD I. By Prof. T. F. TOUT.  
 HENRY VII. By JAMES GAIRDNER.  
 CARDINAL WOLSEY. By Bishop CREIGHTON.  
 ELIZABETH. By E. S. BEESLY.  
 OLIVER CROMWELL. By FREDERIC HARRISON.  
 WILLIAM III. By H. D. TRAILL.  
 WALPOLE. By JOHN MORLEY.  
 CHATHAM. By JOHN MORLEY.  
 PITT. By Earl of ROSEBURY.  
 PEELE. By J. R. THURSFIELD.  
 FISKE.—Works by JOHN FISKE, formerly Lecturer on Philosophy at Harvard University.  
 THE CRITICAL PERIOD IN AMERICAN HISTORY, 1783-1789. 10s. 6d.  
 THE BEGINNINGS OF NEW ENGLAND. Cr. 8vo. 7s. 6d.  
 THE AMERICAN REVOLUTION. 2 vols. Cr. 8vo. 18s.  
 THE DISCOVERY OF AMERICA. 2 vols. Cr. 8vo. 18s.  
 FREEMAN.—Works by the late EDWARD A. FREEMAN, D.C.L.  
 \*OLD ENGLISH HISTORY. With Maps. Ex. fcap. 8vo. 6s.  
 METHODS OF HISTORICAL STUDY. 8vo. 10s. 6d.  
 THE CHIEF PERIODS OF EUROPEAN HISTORY. 8vo. 10s. 6d.  
 HISTORICAL ESSAYS. 8vo. First Series. 10s. 6d. Second Series. 10s. 6d. Third Series. 12s. Fourth Series. 12s. 6d.  
 THE GROWTH OF THE ENGLISH CONSTITUTION FROM THE EARLIEST TIMES. 5th Ed. Cr. 8vo. 5s.  
 WESTERN EUROPE IN THE FIFTH CENTURY. 8vo. [*In the Press.*]  
 WESTERN EUROPE IN THE EIGHTH CENTURY. 8vo. [*In the Press.*]  
 GREEN.—Works by JOHN RICHARD GREEN, LL.D.  
 \*A SHORT HISTORY OF THE ENGLISH PEOPLE. Cr. 8vo. 8s. 6d.  
 \*Also in Four Parts. With Analysis. Crown 8vo. 3s. each. Part I. 607-1265.



- Part II. 1204-1553. Part III. 1540-1689. Part IV. 1660-1873. Illustrated Edition. 8vo. Monthly parts, 1s. net. Part I. Oct. 1891. Vols. I. II. and III. 12s. each net.
- HISTORY OF THE ENGLISH PEOPLE.** In four vols. 8vo. 16s. each.
- Vol. I.—Early England, 449-1071; Foreign Kings, 1071-1214; The Charter, 1214-1291; The Parliament, 1307-1461. 8 Maps.
- Vol. II.—The Monarchy, 1461-1540; The Reformation, 1540-1603.
- Vol. III.—Puritan England, 1603-1660; The Revolution, 1660-1683. 4 Maps.
- Vol. IV.—The Revolution, 1688-1760; Modern England, 1760-1815.
- THE MAKING OF ENGLAND (449-829).** With Maps. 8vo. 16s.
- THE CONQUEST OF ENGLAND (758-1071).** With Maps and Portrait. 8vo. 18s.
- \***ANALYSIS OF ENGLISH HISTORY**, based on Green's "Short History of the English People." By C. W. A. TAIT, M.A. Crown 8vo. 3s. 6d.
- \***READINGS IN ENGLISH HISTORY.** Selected by J. R. GREEN. Three Parts. Gl. 8vo. 1s. 6d. each. I. Hengist to Cressy. II. Cressy to Cromwell. III. Cromwell to Balaklava.
- GREEN.—THE ENGLISH TOWN IN THE FIFTEENTH CENTURY.** By A. S. GREEN. [In the Press.]
- GUEST.—LECTURES ON THE HISTORY OF ENGLAND.** By M. J. GUEST. With Maps. Cr. 8vo. 6s.
- HISTORICAL COURSE FOR SCHOOLS.**—Edited by E. A. FREEMAN. 18mo.
- GENERAL SKETCH OF EUROPEAN HISTORY.** By E. A. FREEMAN. 3s. 6d.
- HISTORY OF ENGLAND.** By EDITH THOMPSON. 2s. 6d.
- HISTORY OF SCOTLAND.** By MARGARET MACARTHUR. 2s.
- HISTORY OF FRANCE.** By CHARLOTTE M. YONGE. 3s. 6d.
- HISTORY OF GERMANY.** By J. SIME, M.A. 3s.
- HISTORY OF ITALY.** By Rev. W. HUNT, M.A. 3s. 6d.
- HISTORY OF AMERICA.** By JOHN A. DOYLE. 4s. 6d.
- HISTORY OF EUROPEAN COLONIES.** By E. J. PAYNE, M.A. 4s. 6d.
- \***HISTORY PRIMERS.**—Edited by JOHN RICHARD GREEN, LL.D. 18mo. 1s. each.
- ROME. By Bishop CREIGHTON.
- GREECE. By C. A. FYFFE, M.A., late Fellow of University College, Oxford.
- EUROPE. By E. A. FREEMAN, D.C.L.
- FRANCE. By CHARLOTTE M. YONGE.
- ROMAN ANTIQUITIES. By Prof. WILKINS, Litt.D. Illustrated.
- GREEK ANTIQUITIES. By Rev. J. P. MAHAFFY, D.D. Illustrated.
- GEOGRAPHY. By Sir G. GROVE, D.C.L. Maps.
- CLASSICAL GEOGRAPHY. By H. F. TOZER, M.A.
- ENGLAND. By ARABELLA B. BUCKLEY.
- ANALYSIS OF ENGLISH HISTORY.** By Prof. T. F. TOUT, M.A.
- INDIAN HISTORY: ASIATIC AND EUROPEAN.** By J. TALBOYS WHEELER.
- HOLE.—A GENEALOGICAL STEMMA OF THE KINGS OF ENGLAND AND FRANCE.** By Rev. C. HOLE. On Sheet. 1s.
- JENNINGS.—CHRONOLOGICAL TABLES OF ANCIENT HISTORY.** By Rev. A. C. JENNINGS. 8vo. 5s.
- LABBERTON.—NEW HISTORICAL ATLAS AND GENERAL HISTORY.** By R. H. LABBERTON. 4to. 15s.
- LETHBRIDGE.—A SHORT MANUAL OF THE HISTORY OF INDIA.** With an Account of INDIA as it is. By Sir ROPER LETHBRIDGE. Cr. 8vo. 5s.
- A HISTORY OF INDIA.** New Edition. Cr. 8vo. 2s.
- \***MACMILLAN'S HISTORY READERS.** Adapted to the New Code, 1893. Gl. 8vo. Book I. (*Shortly.*) Book II. 10d. Book III. 1s. Book IV. 1s. 3d. Book V. 1s. 6d. Book VI. 1s. 6d. Book VII. 1s. 6d.
- MAHAFFY.—GREEK LIFE AND THOUGHT FROM THE AGE OF ALEXANDER TO THE ROMAN CONQUEST.** (*See Classics*, p. 13.)
- THE GREEK WORLD UNDER ROMAN SWAY.** (*See Classics*, p. 13.)
- PROBLEMS IN GREEK HISTORY.** (*See Classics*, p. 13.)
- MARRIOTT.—THE MAKERS OF MODERN ITALY: MAZZINI, CAVOUR, GARIBALDI.** By J. A. R. MARRIOTT, M.A. Cr. 8vo. 1s. 6d.
- MICHELET.—A SUMMARY OF MODERN HISTORY.** By M. MICHELET. Translated by M. C. M. SIMPSON. Gl. 8vo. 4s. 6d.
- NORGATE.—ENGLAND UNDER THE ANGEVIN KINGS.** By KATE NORGATE. With Maps and Plans. 2 vols. 8vo. 32s.

- OTTÉ.**—SCANDINAVIAN HISTORY. By E. C. OTTÉ. With Maps. Gl. 8vo. 6s.  
**RHOADES.**—HISTORY OF THE UNITED STATES. 1850-1880. By J. F. RHOADES. 2 vols. 8vo. 24s.  
**SHUCKBURGH.**—A SCHOOL HISTORY OF ROME. (See p. 13.)  
**SEELEY.**—THE EXPANSION OF ENGLAND. By J. R. SEELEY, M.A., Regius Professor of Modern History in the University of Cambridge. Cr. 8vo. 4s. 6d.  
**OUR COLONIAL EXPANSION.** Extracts from the above. Cr. 8vo. Sewed. 1s.  
**SEWELL-YONGE.**—EUROPEAN HISTORY. Selections from the Best Authorities. Edited by E. M. SEWELL and C. M. YONGE. Cr. 8vo. First Series, 1003-1154. 6s. Second Series, 1088-1228. 6s.  
**\*TAIT.**—ANALYSIS OF ENGLISH HISTORY. (See under Green, p. 43.)  
**WHEELER.**—Works by J. TALBOYS WHEELER.  
**\*A PRIMER OF INDIAN HISTORY.** 18mo. 1s.  
**\*COLLEGE HISTORY OF INDIA.** With Maps. Cr. 8vo. 3s.; sewed, 2s. 6d.  
**A SHORT HISTORY OF INDIA AND OF THE FRONTIER STATES OF AFGHANISTAN, NEPAUL, AND BURMA.** With Maps. Cr. 8vo. 12s.  
**YONGE.**—Works by CHARLOTTE M. YONGE.  
**CAMEOS FROM ENGLISH HISTORY.** Ex. fcap. 8vo. 5s. each. (1) From Rollo to Edward II. (2) The Wars in France. (3) The Wars of the Roses. (4) Reformation Times. (5) England and Spain. (6) Forty Years of Stewart Rule (1603-1643). (7) Rebellion and Restoration (1642-1678).  
**THE VICTORIAN HALF CENTURY.** Cr. 8vo. 1s. 6d.; sewed, 1s.

## ART.

- \*ANDERSON.**—LINEAR PERSPECTIVE AND MODEL DRAWING. With Questions and Exercises. By LAURENCE ANDERSON. Illustrated. 8vo. 2s.  
**BENSON.**—See Technology, p. 41.  
**COLLIER.**—A PRIMER OF ART. By Hon. JOHN COLLIER. 18mo. 1s.  
**COOK.**—THE NATIONAL GALLERY, A POPULAR HANDBOOK TO. By E. T. Cook, with preface by Mr. RUSKIN, and Selections from his Writings. 4th Ed., 1893. Cr. 8vo. Half-mor., 14s.  
**DELAMOTTE.**—A BEGINNER'S DRAWING BOOK. By P. H. DELAMOTTE, F.S.A. Progressively arranged. Cr. 8vo. 3s. 6d.  
**ELLIS.**—SKETCHING FROM NATURE. A Handbook. By TRISTRAM J. ELLIS. Illustrated by H. STACY MARKS, R.A., and the Author. Cr. 8vo. 3s. 6d.  
**GROVE.**—A DICTIONARY OF MUSIC AND MUSICIANS. 1450-1889. Edited by Sir GEORGE GROVE. 4 vols. 8vo. 21s. each. INDEX. 7s. 6d.  
**HUNT.**—TALKS ABOUT ART. By WILLIAM HUNT. Cr. 8vo. 3s. 6d.  
**HUTCHINSON.**—SOME HINTS ON LEARNING TO DRAW. Containing Examples from Leighton, Watts, Poynter, etc. By G. W. C. HUTCHINSON, Art Master at Clifton College. Sup. Roy. 8vo. 8s. 6d.  
**LETHABY.**—See under Technology, p. 41.  
**MELDOLA.**—THE CHEMISTRY OF PHOTOGRAPHY. By RAPHAEL MELDOLA, F.R.S., Professor of Chemistry in the Technical College, Finsbury. Cr. 8vo. 6s.  
**TAYLOR.**—A PRIMER OF PIANOFORTE-PLAYING. By F. TAYLOR. 18mo. 1s.  
**TAYLOR.**—A SYSTEM OF SIGHT-SINGING FROM THE ESTABLISHED MUSICAL NOTATION; based on the Principle of Tonic Relation. By SEDLEY TAYLOR, M.A. 8vo. 5s. net.  
**TAYLOR.**—DRAWING AND DESIGN. By E. R. TAYLOR, Principal of the Birmingham School of Art. Illustrated. Oblong Cr. 8vo.  
**TYRWHITT.**—OUR SKETCHING CLUB. Letters and Studies on Landscape Art. By Rev. R. St. JOHN TYRWHITT. With reproductions of the Lessons and Woodcuts in Mr. Ruskin's "Elements of Drawing." Cr. 8vo. 7s. 6d.

## DIVINITY.

The Bible; History of the Christian Church; The Church of England; The Fathers; Hymnology.

### THE BIBLE.

*History of the Bible.*—THE ENGLISH BIBLE; A Critical History of the various English Translations. By Prof. JOHN EADIE. 2 vols. 8vo. 28s.

- THE BIBLE IN THE CHURCH. By Right Rev. B. F. WESTCOTT, Bishop of Durham. 10th Ed. 18mo. 4s. 6d.
- Biblical History*.—BIBLE LESSONS. By Rev. E. A. ABBOTT. Cr. 8vo. 4s. 6d.
- SIDE-LIGHTS UPON BIBLE HISTORY. By Mrs. SYDNEY BUXTON. Cr. 8vo. 5s.
- STORIES FROM THE BIBLE. By Rev. A. J. CHURCH. Illustrated. Cr. 8vo. 2 parts. 3s. 6d. each.
- \*BIBLE READINGS SELECTED FROM THE PENTATEUCH AND THE BOOK OF JOSHUA. By Rev. J. A. CROSS. Gl. 8vo. 2s. 6d.
- \*THE CHILDREN'S TREASURY OF BIBLE STORIES. By Mrs. H. GASKOIN. 18mo. 1s. each. Part I. OLD TESTAMENT. Part II. NEW TESTAMENT. Part III. THE APOSTLES.
- \*A CLASS-BOOK OF OLD TESTAMENT HISTORY. By Rev. G. F. MACLEAR, D.D. 18mo. 4s. 6d.
- \*A CLASS-BOOK OF NEW TESTAMENT HISTORY. 18mo. 5s. 6d.
- \*A SHILLING BOOK OF OLD TESTAMENT HISTORY. 18mo. 1s.
- \*A SHILLING BOOK OF NEW TESTAMENT HISTORY. 18mo. 1s.
- \*SCRIPTURE READINGS FOR SCHOOLS AND FAMILIES. By C. M. YONGE. Globe 8vo. 1s. 6d. each; also with comments, 3s. 6d. each. GENESIS TO DEUTERONOMY. JOSHUA TO SOLOMON. KINGS AND THE PROPHETS. THE GOSPEL TIMES. APOSTOLIC TIMES.
- The Old Testament*.—THE PATRIARCHS AND LAWGIVERS OF THE OLD TESTAMENT. By F. D. MAURICE. Cr. 8vo. 3s. 6d.
- THE PROPHETS AND KINGS OF THE OLD TESTAMENT. By the same. Cr. 8vo. 3s. 6d.
- THE CANON OF THE OLD TESTAMENT. By Rev. H. E. RYLE, B.D., Hulsean Professor of Divinity in the University of Cambridge. Cr. 8vo. 6s.
- THE EARLY NARRATIVES OF GENESIS. By the same. Cr. 8vo. 3s. net.
- THE DIVINE LIBRARY OF THE OLD TESTAMENT. By A. F. KIRKPATRICK, M.A., Professor of Hebrew in the University of Cambridge. Cr. 8vo. 3s. net.
- The Pentateuch*.—AN HISTORICO-CRITICAL INQUIRY INTO THE ORIGIN AND COMPOSITION OF THE PENTATEUCH AND BOOK OF JOSHUA. By Prof. A. KUENEN. Trans. by P. H. WICKSTEED, M.A. 8vo. 14s.
- The Psalms*.—THE PSALMS CHRONOLOGICALLY ARRANGED. By FOUR FRIENDS. Cr. 8vo. 5s. net.
- GOLDEN TREASURY PSALTER. Student's Edition of above. 18mo. 2s. 6d. net.
- THE PSALMS, WITH INTRODUCTION AND NOTES. By A. C. JENNINGS, M.A., and W. H. LOWE, M.A. 2 vols. Cr. 8vo. 10s. 6d. each.
- INTRODUCTION TO THE STUDY AND USE OF THE PSALMS. By Rev. J. F. THRUPE. 2nd Ed. 2 vols. 8vo. 21s.
- Isaiah*.—ISAIAH XL.—LXVI. With the Shorter Prophecies allied to it. Edited by MATTHEW ARNOLD. Cr. 8vo. 5s.
- ISAIAH OF JERUSALEM. In the Authorised English Version, with Introduction and Notes. By the same. Cr. 8vo. 4s. 6d.
- A BIBLE-READING FOR SCHOOLS,—THE GREAT PROPHECY OF ISRAEL'S RESTORATION (Isaiah, Chapters xl.—lxvi.) Arranged and Edited for Young Learners. By the same. 18mo. 1s.
- COMMENTARY ON THE BOOK OF ISAIAH: CRITICAL, HISTORICAL, AND PROPHETICAL; with Translation. By T. R. BIRKS. 8vo. 12s. 6d.
- THE BOOK OF ISAIAH CHRONOLOGICALLY ARRANGED. By T. K. CHEYNE. Cr. 8vo. 7s. 6d.
- Zechariah*.—THE HEBREW STUDENT'S COMMENTARY ON ZECHARIAH, HEBREW AND LXX. By W. H. LOWE, M.A. 8vo. 10s. 6d.
- The Minor Prophets*.—DOCTRINE OF THE PROPHETS. By Prof. A. F. KIRKPATRICK. Cr. 8vo. 6s.
- The New Testament*.—THE NEW TESTAMENT. Essay on the Right Estimation of MS. Evidence in the Text of the New Testament. By T. R. BIRKS. Cr. 8vo. 3s. 6d.
- THE MESSAGES OF THE BOOKS. Discourses and Notes on the Books of the New Testament. By Archd. FARRAR. 8vo. 14s.
- GREEK-ENGLISH LEXICON TO THE NEW TESTAMENT. By W. J. HICKIE, M.A. Pott. 8vo. 3s.
- THE CLASSICAL ELEMENT IN THE NEW TESTAMENT. Considered as a proof of its Genuineness. By C. H. HOOLE. 8vo. 10s. 6d.



- ON A FRESH REVISION OF THE ENGLISH NEW TESTAMENT. By Bishop LIGHTFOOT. Cr. 8vo. 7s. 6d.
- THE UNITY OF THE NEW TESTAMENT. By F. D. MAURICE. 2 vols. Cr. 8vo. 12s.
- A GENERAL SURVEY OF THE HISTORY OF THE CANON OF THE NEW TESTAMENT DURING THE FIRST FOUR CENTURIES. By Bishop WESTCOTT. Cr. 8vo. 10s. 6d.
- THE NEW TESTAMENT IN THE ORIGINAL GREEK. The Text revised by Bishop WESTCOTT, D.D., and Prof. F. J. A. HORT, D.D. 2 vols. Cr. 8vo. 10s. 6d. each. Vol. I. Text. Vol. II. Introduction and Appendix.
- SCHOOL EDITION OF THE ABOVE. 18mo, 4s. 6d.; 18mo, roan, 5s. 6d.; morocco, gilt edges, 6s. 6d.
- The Gospels.*—THE COMMON TRADITION OF THE SYNOPTIC GOSPELS, in the Text of the Revised Version. By Rev. E. A. ABBOTT and W. G. RUSHBROOKE. Cr. 8vo. 3s. 6d.
- SYNOPTICON: AN EXPOSITION OF THE COMMON MATTER OF THE SYNOPTIC GOSPELS. By W. G. RUSHBROOKE. Printed in Colours. In six Parts, and Appendix. 4to. Part I. 3s. 6d. Parts II. and III. 7s. Parts IV. V. and VI., with Indices, 10s. 6d. Appendices, 10s. 6d. Complete in 1 vol. 35s. "Indispensable to a Theological Student."—*The Cambridge Guide.*
- ESSAYS ON THE WORK ENTITLED "SUPERNATURAL RELIGION." A discussion of the authenticity of the Gospels. By Bishop LIGHTFOOT. 2nd Ed. 8vo. 10s. 6d.
- INTRODUCTION TO THE STUDY OF THE FOUR GOSPELS. By Bishop WESTCOTT. Cr. 8vo. 10s. 6d.
- THE COMPOSITION OF THE FOUR GOSPELS. By Rev. A. WRIGHT. Cr. 8vo. 5s.
- THE SYNOPTIC PROBLEM FOR ENGLISH READERS. By A. J. JOLLEY. 3s. net.
- THE AKHMIM FRAGMENT OF THE APOCRYPHAL GOSPEL OF ST. PETER. With an Introduction, Notes, and Indices. By H. B. SWETE, D.D., Litt.D. 8vo. 5s. net.
- The Gospel according to St. Matthew.*—\*THE GREEK TEXT, with Introduction and Notes by Rev. A. SLOMAN. Fcap. 8vo. 2s. 6d.
- CHOICE NOTES ON ST. MATTHEW. Drawn from Old and New Sources. Cr. 8vo. 4s. 6d. (St. Matthew and St. Mark in 1 vol. 9s.)
- The Gospel according to St. Mark.*—\*SCHOOL READINGS IN THE GREEK TESTAMENT. Being the Outlines of the Life of our Lord as given by St. Mark, with additions from the Text of the other Evangelists. Edited, with Notes and Vocabulary, by Rev. A. CALVERT, M.A. Fcap. 8vo. 2s. 6d.
- THE GREEK TEXT, with Introduction and Notes. By Rev. J. O. F. MURRAY, M.A. [In preparation.]
- CHOICE NOTES ON ST. MARK. Cr. 8vo. 4s. 6d.
- The Gospel according to St. Luke.*—\*THE GREEK TEXT, with Introduction and Notes by Rev. J. BOND, M.A. Fcap. 8vo. 2s. 6d.
- CHOICE NOTES ON ST. LUKE. Cr. 8vo. 4s. 6d.
- THE GOSPEL OF THE KINGDOM OF HEAVEN. A Course of Lectures on the Gospel of St. Luke. By F. D. MAURICE. Cr. 8vo. 3s. 6d.
- The Gospel according to St. John.*—THE GOSPEL OF ST. JOHN. By F. D. MAURICE. Cr. 8vo. 3s. 6d.
- CHOICE NOTES ON ST. JOHN. Cr. 8vo. 4s. 6d.
- The Acts of the Apostles.*—\*THE GREEK TEXT, with Notes by T. E. PAGE, M.A. Fcap. 8vo. 3s. 6d.
- THE CHURCH OF THE FIRST DAYS: THE CHURCH OF JERUSALEM, THE CHURCH OF THE GENTILES, THE CHURCH OF THE WORLD. Lectures on the Acts of the Apostles. By Very Rev. C. J. VAUGHAN. Cr. 8vo. 10s. 6d.
- THE OLD SYRIAC ELEMENT IN THE TEXT OF THE CODEX BEZAE. By Rev. F. H. CHASE. 8vo. 7s. 6d. net.
- The Epistles of St. Paul.*—THE EPISTLE TO THE ROMANS. The Greek Text, with English Notes. By the Very Rev. C. J. VAUGHAN. 7th Ed. Cr. 8vo. 7s. 6d.
- THE EPISTLES TO THE CORINTHIANS. Greek Text, with Commentary. By Rev. W. KAY. 8vo. 9s.

- THE EPISTLE TO THE GALATIANS. A Revised Text, with Introduction, Notes, and Dissertations. By Bishop LIGHTFOOT. 10th Ed. 8vo. 12s.
- THE EPISTLE TO THE PHILIPPIANS. A Revised Text, with Introduction, Notes, and Dissertations. By the same. 8vo. 12s.
- THE EPISTLE TO THE PHILIPPIANS. With Translation, Paraphrase, and Notes for English Readers. By Very Rev. C. J. VAUGHAN. Cr. 8vo. 5s.
- THE EPISTLE TO THE COLOSSIANS AND TO PHILEMON. A Revised Text, with Introductions, etc. By Bishop LIGHTFOOT. 9th Ed. 8vo. 12s.
- THE EPISTLES TO THE EPHESIANS, THE COLOSSIANS, AND PHILEMON. With Introduction and Notes. By Rev. J. LL. DAVIES. 8vo. 7s. 6d.
- THE FIRST EPISTLE TO THE THESSALONIANS. By Very Rev. C. J. VAUGHAN. 8vo. Sewed, 1s. 6d.
- THE EPISTLES TO THE THESSALONIANS. Commentary on the Greek Text. By Prof. JOHN EADIE. 8vo. 12s.
- INTRODUCTORY LECTURES ON THE EPISTLES TO THE ROMANS AND TO THE EPHESIANS. By the late Prof. HORT. Cr. 8vo. [*In preparation.*]
- The Epistle of St. James.*—THE GREEK TEXT, with Introduction and Notes. By Rev. JOSEPH B. MAYOR. 8vo. 14s.
- The Epistles of St. John.*—THE EPISTLES OF ST. JOHN. By F. D. MAURICE. Cr. 8vo. 3s. 6d.
- THE GREEK TEXT, with Notes. By Bishop WESTCOTT. 2nd Ed. 8vo. 12s. 6d.
- The Epistle to the Hebrews.*—GREEK AND ENGLISH. Edited by Rev. F. RENDALL. Cr. 8vo. 6s.
- ENGLISH TEXT, with Commentary. By the same. Cr. 8vo. 7s. 6d.
- THE GREEK TEXT, with Notes. By Very Rev. C. J. VAUGHAN. Cr. 8vo. 7s. 6d.
- THE GREEK TEXT, with Notes and Essays. By Bishop WESTCOTT. 8vo. 14s.
- Revelation.*—LECTURES ON THE APOCALYPSE. By F. D. MAURICE. Cr. 8vo. 3s. 6d.
- THE REVELATION OF ST. JOHN. By Prof. W. MILLIGAN. Cr. 8vo. 7s. 6d.
- LECTURES ON THE APOCALYPSE. By the same. Cr. 8vo. 5s.
- DISCUSSIONS ON THE APOCALYPSE. By the same. Crown 8vo. 5s.
- LECTURES ON THE REVELATION OF ST. JOHN. By Very Rev. C. J. VAUGHAN. 5th Ed. Cr. 8vo. 10s. 6d.

---

WRIGHT.—THE BIBLE WORD-BOOK. By W. ALDIS WRIGHT. Cr. 8vo. 7s. 6d.

## HISTORY OF THE CHRISTIAN CHURCH.

- CUNNINGHAM.—THE GROWTH OF THE CHURCH IN ITS ORGANISATION AND INSTITUTIONS. By Rev. JOHN CUNNINGHAM. 8vo. 9s.
- CUNNINGHAM.—THE CHURCHES OF ASIA: A METHODICAL SKETCH OF THE SECOND CENTURY. By Rev. WILLIAM CUNNINGHAM. Cr. 8vo. 6s.
- DALE.—THE SYNOD OF ELVIRA, AND CHRISTIAN LIFE IN THE FOURTH CENTURY. By A. W. W. DALE. Cr. 8vo. 10s. 6d.
- HARDWICK.—Works by Archdeacon HARDWICK.
- A HISTORY OF THE CHRISTIAN CHURCH: MIDDLE AGE. Edited by Bishop STUBBS. Cr. 8vo. 10s. 6d.
- A HISTORY OF THE CHRISTIAN CHURCH DURING THE REFORMATION. 9th Ed., revised by Bishop STUBBS. Cr. 8vo. 10s. 6d.
- HORT.—Works by the late Prof. F. J. A. HORT, D.D.
- TWO DISSERTATIONS. I. ON ΜΟΝΟΤΕΝΗΣ ΘΕΟΣ IN SCRIPTURE AND TRADITION. II. ON THE "CONSTANTINOPOLITAN" CREED AND OTHER CREEDS OF THE FOURTH CENTURY. 8vo. 7s. 6d.
- LECTURES ON JUDAISTIC CHRISTIANITY. Cr. 8vo. [*In the Press.*]
- LECTURES ON EARLY CHURCH HISTORY. Cr. 8vo. [*In the Press.*]
- KILLEN.—ECCLESIASTICAL HISTORY OF IRELAND, from the earliest date to the present time. By W. D. KILLEN. 2 vols. 8vo. 25s.
- SIMPSON.—AN EPITOME OF THE HISTORY OF THE CHRISTIAN CHURCH. By Rev. W. SIMPSON. 7th Ed. Fcap. 8vo. 3s. 6d.
- VAUGHAN.—THE CHURCH OF THE FIRST DAYS: THE CHURCH OF JERUSALEM, THE CHURCH OF THE GENTILES, THE CHURCH OF THE WORLD. By Very Rev. C. J. VAUGHAN. Cr. 8vo. 10s. 6d.

## THE CHURCH OF ENGLAND.

**BENHAM.**—A COMPANION TO THE LECTIONARY. By Rev. W. BENHAM, B.D. Cr. Svo. 4s. 6d.

CATECHISM AND CONFIRMATION. Pott Svo. 1s.

**COLENSO.**—THE COMMUNION SERVICE FROM THE BOOK OF COMMON PRAYER. With Select Readings from the Writings of the Rev. F. D. MAURICE. Edited by Bishop COLENSO. 6th Ed. 16mo. 2s. 6d.

**MACLEAR.**—Works by Rev. G. F. MACLEAR, D.D.

\*A CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND. 18mo. 1s. 6d.

\*A FIRST CLASS-BOOK OF THE ABOVE. 18mo. 6d.

THE ORDER OF CONFIRMATION. With Prayers and Devotions. 32mo. 6d.

FIRST COMMUNION. With Prayers and Devotions for the newly Confirmed. 32mo. 6d.

\*A MANUAL OF INSTRUCTION FOR CONFIRMATION AND FIRST COMMUNION. With Prayers and Devotions. 32mo. 2s.

\*AN INTRODUCTION TO THE CREEDS. 18mo. 3s. 6d.

AN INTRODUCTION TO THE THIRTY-NINE ARTICLES. [*In the Press.*]

**PROCTER.**—A HISTORY OF THE BOOK OF COMMON PRAYER. By Rev. F. PROCTER. 18th Ed. Cr. Svo. 10s. 6d.

\***PROCTER—MACLEAR.**—AN ELEMENTARY INTRODUCTION TO THE BOOK OF COMMON PRAYER. By Rev. F. PROCTER and Rev. G. F. MACLEAR, D.D. 18mo. 2s. 6d.

**VAUGHAN.**—TWELVE DISCOURSES ON SUBJECTS CONNECTED WITH THE LITURGY AND WORSHIP OF THE CHURCH OF ENGLAND. By Very Rev. C. J. VAUGHAN. Fcap. 8vo. 6s.

NOTES FOR LECTURES ON CONFIRMATION. With suitable Prayers. By the same. 18mo. 1s. 6d.

## THE FATHERS.

**CUNNINGHAM.**—THE EPISTLE OF ST. BARNABAS. The Greek Text, the Latin Version, and a new English Translation and Commentary. By Rev. W. CUNNINGHAM. Cr. Svo. 7s. 6d.

**DONALDSON.**—THE APOSTOLICAL FATHERS. A Critical Account of their Genuine Writings, and of their Doctrines. By Prof. JAMES DONALDSON. 2nd Ed. Cr. Svo. 7s. 6d.

**GWATKIN.**—SELECTIONS FROM THE EARLY CHRISTIAN WRITERS. By Rev. Prof. GWATKIN. Cr. Svo.

EARLY HISTORY OF THE CHRISTIAN CHURCH. By the same. [*In prep.*]

**LIGHTFOOT.**—Works by Bishop LIGHTFOOT.

THE APOSTOLIC FATHERS. Revised Texts, with Introductions, Notes, Dissertations, and Translations. Part I. ST. CLEMENT OF ROME. 2 vols. 8vo. 32s.

THE APOSTOLIC FATHERS. Part II. ST. IGNATIUS to ST. POLYCARP. 3 vols. 2nd Ed. 8vo. 48s.

ABRIDGED EDITION. With short Introductions, Greek Text, and Translation. 8vo. 16s.

## HYMNOLOGY.

**PALGRAVE.**—ORIGINAL HYMNS. By Prof. F. T. PALGRAVE. 18mo. 1s. 6d.

**SELBORNE.**—THE BOOK OF PRAISE. By EARL OF SELBORNE. 18mo. 2s. 6d. net.

A HYMNAL. A. 32mo. 6d. B. 18mo, larger type. 1s. C. Fine Paper 1s. 6d. Edited, with Music, by JOHN HULLAH. 18mo. 3s. 6d.

**WOODS.**—HYMNS FOR SCHOOL WORSHIP. By M. A. WOODS. 18mo. 1s. 6d.





## Date Due

۷۸۲۵۴۱

NOV 24 1995



BOSTON COLLEGE

BOSTON COLLEGE



3 9031 01550250 3

150025

QA533 .H22

MATH. DEPT.

**BOSTON COLLEGE LIBRARY**  
UNIVERSITY HEIGHTS  
CHESTNUT HILL, MASS.

Books may be kept for two weeks and may be renewed for the same period, unless reserved.

Two cents a day is charged for each book kept overtime.

If you cannot find what you want, ask the Librarian who will be glad to help you.

The borrower is responsible for books drawn on his card and for all fines accruing on the same.

